



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07D 233/54, A61K 31/495, 31/415, C07D 241/08, 295/18, 401/06, 401/14, 403/06, 403/12	A1	(11) International Publication Number: WO 96/30343 (43) International Publication Date: 3 October 1996 (03.10.96)																					
(21) International Application Number: PCT/US96/04019 (22) International Filing Date: 25 March 1996 (25.03.96) (30) Priority Data: <table border="0"> <tr> <td>08/412,829</td> <td>29 March 1995 (29.03.95)</td> <td>US</td> </tr> <tr> <td>08/470,690</td> <td>6 June 1995 (06.06.95)</td> <td>US</td> </tr> <tr> <td>08/600,728</td> <td>1 March 1996 (01.03.96)</td> <td>US</td> </tr> </table> (60) Parent Applications or Grants (63) Related by Continuation <table border="0"> <tr> <td>US</td> <td>08/600,728 (CON)</td> </tr> <tr> <td>Filed on</td> <td>1 March 1996 (01.03.96)</td> </tr> <tr> <td>US</td> <td>08/412,829 (CON)</td> </tr> <tr> <td>Filed on</td> <td>29 March 1995 (29.03.95)</td> </tr> <tr> <td>US</td> <td>08/470,690 (CON)</td> </tr> <tr> <td>Filed on</td> <td>6 June 1995 (06.06.95)</td> </tr> </table> (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		08/412,829	29 March 1995 (29.03.95)	US	08/470,690	6 June 1995 (06.06.95)	US	08/600,728	1 March 1996 (01.03.96)	US	US	08/600,728 (CON)	Filed on	1 March 1996 (01.03.96)	US	08/412,829 (CON)	Filed on	29 March 1995 (29.03.95)	US	08/470,690 (CON)	Filed on	6 June 1995 (06.06.95)	(72) Inventors; and (75) Inventors/Applicants (for US only): ANTHONY, Neville, J. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). CICCARONE, Terrence, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). GOMEZ, Robert, P. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HUTCHINSON, John, H. [CA/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). WILLIAMS, Theresa, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). DINSMORE, Christopher, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). STOKKER, Gerald, E. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (81) Designated States: AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IS, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
08/412,829	29 March 1995 (29.03.95)	US																					
08/470,690	6 June 1995 (06.06.95)	US																					
08/600,728	1 March 1996 (01.03.96)	US																					
US	08/600,728 (CON)																						
Filed on	1 March 1996 (01.03.96)																						
US	08/412,829 (CON)																						
Filed on	29 March 1995 (29.03.95)																						
US	08/470,690 (CON)																						
Filed on	6 June 1995 (06.06.95)																						
(54) Title: INHIBITORS OF FARNESYL-PROTEIN TRANSFERASE (57) Abstract <p>The present invention is directed to compounds which inhibit farnesyl-protein transferase (FTase) and the farnesylation of the oncogene protein Ras. The invention is further directed to chemotherapeutic compositions containing the compounds of this invention and methods for inhibiting farnesyl-protein transferase and the farnesylation of the oncogene protein Ras.</p>																							

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

TITLE OF THE INVENTION

INHIBITORS OF FARNESYL-PROTEIN TRANSFERASE

RELATED APPLICATION

5 The present patent application is a continuation-in-part application of copending application Serial No. 08/470,690, filed June 6, 1995, which is a continuation-in-part application of copending application Serial No. 08/412,829, filed March 29, 1995.

10 BACKGROUND OF THE INVENTION

 The Ras proteins (Ha-Ras, Ki4a-Ras, Ki4b-Ras and N-Ras) are part of a signalling pathway that links cell surface growth factor receptors to nuclear signals initiating cellular proliferation. Biological and biochemical studies of Ras action indicate that Ras functions like a
15 G-regulatory protein. In the inactive state, Ras is bound to GDP. Upon growth factor receptor activation Ras is induced to exchange GDP for GTP and undergoes a conformational change. The GTP-bound form of Ras propagates the growth stimulatory signal until the signal is terminated by the intrinsic GTPase activity of Ras, which returns the
20 protein to its inactive GDP bound form (D.R. Lowy and D.M. Willumsen, *Ann. Rev. Biochem.* 62:851-891 (1993)). Mutated *ras* genes (Ha-*ras*, Ki4a-*ras*, Ki4b-*ras* and N-*ras*) are found in many human cancers, including colorectal carcinoma, exocrine pancreatic carcinoma, and myeloid leukemias. The protein products of these genes are
25 defective in their GTPase activity and constitutively transmit a growth stimulatory signal.

 Ras must be localized to the plasma membrane for both normal and oncogenic functions. At least 3 post-translational modifications are involved with Ras membrane localization, and all 3
30 modifications occur at the C-terminus of Ras. The Ras C-terminus contains a sequence motif termed a "CAAX" or "Cys-Aaa¹-Aaa²-Xaa" box (Cys is cysteine, Aaa is an aliphatic amino acid, the Xaa is any amino acid) (Willumsen *et al.*, *Nature* 310:583-586 (1984)). Depending on the specific sequence, this motif serves as a signal sequence for the

- 2 -

enzymes farnesyl-protein transferase or geranylgeranyl-protein transferase, which catalyze the alkylation of the cysteine residue of the CAAX motif with a C₁₅ or C₂₀ isoprenoid, respectively. (S. Clarke., *Ann. Rev. Biochem.* 61:355-386 (1992); W.R. Schafer and J. Rine, *Ann. Rev. Genetics* 30:209-237 (1992)). The Ras protein is one of several proteins that are known to undergo post-translational farnesylation. Other farnesylated proteins include the Ras-related GTP-binding proteins such as Rho, fungal mating factors, the nuclear lamins, and the gamma subunit of transducin. James, et al., *J. Biol. Chem.* 269, 14182 (1994) have identified a peroxisome associated protein Pxf which is also farnesylated. James, et al., have also suggested that there are farnesylated proteins of unknown structure and function in addition to those listed above.

Inhibition of farnesyl-protein transferase has been shown to block the growth of Ras-transformed cells in soft agar and to modify other aspects of their transformed phenotype. It has also been demonstrated that certain inhibitors of farnesyl-protein transferase selectively block the processing of the Ras oncoprotein intracellularly (N.E. Kohl *et al.*, *Science*, 260:1934-1937 (1993) and G.L. James *et al.*, *Science*, 260:1937-1942 (1993). Recently, it has been shown that an inhibitor of farnesyl-protein transferase blocks the growth of *ras*-dependent tumors in nude mice (N.E. Kohl *et al.*, *Proc. Natl. Acad. Sci U.S.A.*, 91:9141-9145 (1994) and induces regression of mammary and salivary carcinomas in *ras* transgenic mice (N.E. Kohl *et al.*, *Nature Medicine*, 1:792-797 (1995).

Indirect inhibition of farnesyl-protein transferase *in vivo* has been demonstrated with lovastatin (Merck & Co., Rahway, NJ) and compactin (Hancock *et al.*, *ibid*; Casey *et al.*, *ibid*; Schafer *et al.*, *Science* 245:379 (1989)). These drugs inhibit HMG-CoA reductase, the rate limiting enzyme for the production of polyisoprenoids including farnesyl pyrophosphate. Farnesyl-protein transferase utilizes farnesyl pyrophosphate to covalently modify the Cys thiol group of the Ras CAAX box with a farnesyl group (Reiss *et al.*, *Cell*, 62:81-88 (1990); Schaber *et al.*, *J. Biol. Chem.*, 265:14701-14704 (1990); Schafer *et al.*,

- 3 -

Science, 249:1133-1139 (1990); Manne *et al.*, *Proc. Natl. Acad. Sci USA*, 87:7541-7545 (1990)). Inhibition of farnesyl pyrophosphate biosynthesis by inhibiting HMG-CoA reductase blocks Ras membrane localization in cultured cells. However, direct inhibition of farnesyl-protein transferase would be more specific and attended by fewer side effects than would occur with the required dose of a general inhibitor of isoprene biosynthesis.

Inhibitors of farnesyl-protein transferase (FPTase) have been described in two general classes. The first are analogs of farnesyl diphosphate (FPP), while the second class of inhibitors is related to the protein substrates (e.g., Ras) for the enzyme. The peptide derived inhibitors that have been described are generally cysteine containing molecules that are related to the CAAX motif that is the signal for protein prenylation. (Schaber *et al.*, *ibid*; Reiss *et al.*, *ibid*; Reiss *et al.*, *PNAS*, 88:732-736 (1991)). Such inhibitors may inhibit protein prenylation while serving as alternate substrates for the farnesyl-protein transferase enzyme, or may be purely competitive inhibitors (U.S. Patent 5,141,851, University of Texas; N.E. Kohl *et al.*, *Science*, 260:1934-1937 (1993); Graham, *et al.*, *J. Med. Chem.*, 37, 725 (1994)). In general, deletion of the thiol from a CAAX derivative has been shown to dramatically reduce the inhibitory potency of the compound. However, the thiol group potentially places limitations on the therapeutic application of FPTase inhibitors with respect to pharmacokinetics, pharmacodynamics and toxicity. Therefore, a functional replacement for the thiol is desirable.

It has recently been reported that farnesyl-protein transferase inhibitors are inhibitors of proliferation of vascular smooth muscle cells and are therefore useful in the prevention and therapy of arteriosclerosis and diabetic disturbance of blood vessels (JP H7-112930).

It has recently been disclosed that certain tricyclic compounds which optionally incorporate a piperidine moiety are inhibitors of FPTase (WO 95/10514, WO 95/10515 and WO 95/10516).

- 4 -

Imidazole-containing inhibitors of farnesyl protein transferase have also been disclosed (WO 95/09001 and EP 0 675 112 A1).

It is, therefore, an object of this invention to develop peptidomimetic compounds that do not have a thiol moiety, and that will inhibit farnesyl-protein transferase and thus, the post-translational farnesylation of proteins. It is a further object of this invention to develop chemotherapeutic compositions containing the compounds of this invention and methods for producing the compounds of this invention.

10




SUMMARY OF THE INVENTION

The present invention comprises peptidomimetic piperazine-containing compounds which inhibit the farnesyl-protein transferase. The instant compounds lack a thiol moiety and thus offer unique advantages in terms of improved pharmacokinetic behavior in animals, prevention of thiol-dependent chemical reactions, such as rapid autoxidation and disulfide formation with endogenous thiols, and reduced systemic toxicity. Further contained in this invention are chemotherapeutic compositions containing these farnesyl transferase inhibitors and methods for their production.

20

The compounds of this invention are illustrated by the formulae A, B and C:


- 5 -



- 6 -

DETAILED DESCRIPTION OF THE INVENTION

The compounds of this invention are useful in the inhibition of farnesyl-protein transferase and the farnesylation of the oncogene protein Ras. In a first embodiment of this invention, the inhibitors of farnesyl-protein transferase are illustrated by the formula A:



wherein:


R^{1a} and R^{1b} are independently selected from:

- 10 a) hydrogen,
- b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂N-C(O)-, CN, NO₂, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-,
- 15 c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocyclic, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂N-C(O)-, CN, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, and R¹¹OC(O)-NR¹⁰-;
- 20

R² and R³ are independently selected from: H; unsubstituted or substituted C₁-8 alkyl, unsubstituted or substituted C₂-8 alkenyl,

25 unsubstituted or substituted C₂-8 alkynyl, unsubstituted or substituted aryl,

unsubstituted or substituted heterocycle,



- 7 -

wherein the substituted group is substituted with one or more of:

1) aryl or heterocycle, unsubstituted or substituted with:






- a) C₁-4 alkyl,
- b) (CH₂)_pOR⁶,
- c) (CH₂)_pNR⁶R⁷,
- d) halogen,
- e) CN,

2) C₃-6 cycloalkyl,


3) OR⁶,

4) SR^{6a}, S(O)R^{6a}, SO₂R^{6a},


5) —NR⁶R⁷,



11) —SO₂—NR⁶R⁷,



- 8 -





5 R^2 and R^3 are attached to the same C atom and are combined to form $-(\text{CH}_2)_u-$ wherein one of the carbon atoms is optionally replaced by a moiety selected from: O, $\text{S}(\text{O})_m$, $-\text{NC}(\text{O})-$, and $-\text{N}(\text{COR}^{10})-$;

R^4 and R^5 are independently selected from H and CH_3 ;

10 and any two of R^2 , R^3 , R^4 and R^5 are optionally attached to the same carbon atom;

15 R^6 , R^7 and R^{7a} are independently selected from: H; C_{1-4} alkyl, C_{3-6} cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, unsubstituted or substituted with:

- a) C_{1-4} alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,




20

R^6 and R^7 may be joined in a ring;

- 9 -

R^7 and R^{7a} may be joined in a ring;

R^{6a} is selected from: C_{1-4} alkyl, C_{3-6} cycloalkyl, heterocycle, aryl, unsubstituted or substituted with:

- 5 a) C_{1-4} alkoxy,
 b) aryl or heterocycle,
 c) halogen,
 d) HO,
 e) 
 f) $-\text{SO}_2\text{R}^{11}$, or
 g) $\text{N}(\text{R}^{10})_2$;
- 10

R^8 is independently selected from:

- 15 a) hydrogen,
 b) aryl, heterocycle, $C_3\text{-}C_{10}$ cycloalkyl, $C_2\text{-}C_6$ alkenyl, $C_2\text{-}C_6$ alkynyl, perfluoroalkyl, F, Cl, Br, $\text{R}^{10}\text{O}-$, $\text{R}^{11}\text{S}(\text{O})_m-$, $\text{R}^{10}\text{C}(\text{O})\text{NR}^{10}-$, $(\text{R}^{10})_2\text{NC}(\text{O})-$, $\text{R}^{10}_2\text{N-C}(\text{NR}^{10})-$, CN, NO_2 , $\text{R}^{10}\text{C}(\text{O})-$, $\text{R}^{10}\text{OC}(\text{O})-$, N_3 , $-\text{N}(\text{R}^{10})_2$, or $\text{R}^{11}\text{OC}(\text{O})\text{NR}^{10}-$, and
 c) $C_1\text{-}C_6$ alkyl unsubstituted or substituted by aryl,
 cyanophenyl, heterocycle, $C_3\text{-}C_{10}$ cycloalkyl, $C_2\text{-}C_6$ alkenyl, $C_2\text{-}C_6$ alkynyl, perfluoroalkyl, F, Cl, Br, $\text{R}^{10}\text{O}-$, $\text{R}^{11}\text{S}(\text{O})_m-$, $\text{R}^{10}\text{C}(\text{O})\text{NH}-$, $(\text{R}^{10})_2\text{NC}(\text{O})-$, $\text{R}^{10}_2\text{N-C}(\text{NR}^{10})-$, CN, $\text{R}^{10}\text{C}(\text{O})-$, $\text{R}^{10}\text{OC}(\text{O})-$, N_3 , $-\text{N}(\text{R}^{10})_2$, or $\text{R}^{10}\text{OC}(\text{O})\text{NH}-$;
- 20
- 25

R^9 is selected from:

- 30 a) hydrogen,
 b) $C_2\text{-}C_6$ alkenyl, $C_2\text{-}C_6$ alkynyl, perfluoroalkyl, F, Cl, Br, $\text{R}^{10}\text{O}-$, $\text{R}^{11}\text{S}(\text{O})_m-$, $\text{R}^{10}\text{C}(\text{O})\text{NR}^{10}-$, $(\text{R}^{10})_2\text{NC}(\text{O})-$, $\text{R}^{10}_2\text{N-C}(\text{NR}^{10})-$, CN, NO_2 , $\text{R}^{10}\text{C}(\text{O})-$, $\text{R}^{10}\text{OC}(\text{O})-$, N_3 , $-\text{N}(\text{R}^{10})_2$, or $\text{R}^{11}\text{OC}(\text{O})\text{NR}^{10}-$, and

- 10 -

- c) C₁-C₆ alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

5

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

10

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR¹⁰-, -NR¹⁰C(O)-, O, -N(R¹⁰)-, -S(O)₂N(R¹⁰)-, -N(R¹⁰)S(O)₂-, or S(O)_m;

15 V is selected from:

- a) hydrogen,
- b) heterocycle,
- c) aryl,
- d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C₂-C₂₀ alkenyl,

20

provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

25 W is a heterocycle;

X is -CH₂-, -C(=O)-, or -S(=O)_m-;

30

Y is unsubstituted or substituted aryl or unsubstituted or substituted heterocycle, wherein the substituted aryl or substituted heterocycle is substituted with one or more of:

- 1) C₁-4 alkyl, unsubstituted or substituted with:
 - a) C₁-4 alkoxy,

- 11 -


- 5 b) NR^6R^7 ,
 c) C₃₋₆ cycloalkyl,
 d) aryl or heterocycle,
 e) HO,
 f) $-\text{S}(\text{O})_m\text{R}^{6a}$, or
 g) $-\text{C}(\text{O})\text{NR}^6\text{R}^7$,
 10 2) aryl or heterocycle,
 3) halogen,
 4) OR^6 ,
 5) NR^6R^7 ,
 6) CN,
 7) NO_2 ,
 8) CF_3 ;
 15 9) $-\text{S}(\text{O})_m\text{R}^{6a}$,
 10) $-\text{C}(\text{O})\text{NR}^6\text{R}^7$, or
 11) C_{3-C6} cycloalkyl;

 m is 0, 1 or 2;
 n is 0, 1, 2, 3 or 4;
 20 p is 0, 1, 2, 3 or 4;
 q is 1 or 2;
 r is 0 to 5, provided that r is 0 when V is hydrogen;
 s is 0 or 1;
 t is 0 or 1; and
 25 u is 4 or 5;

or the pharmaceutically acceptable salts thereof.

30 In a second embodiment of this invention, the inhibitors of
 farnesyl-protein transferase are illustrated by the formula B:

- 12 -




wherein:

R^{1a} and R^{1b} are independently selected from:

- 5 a) hydrogen,
- b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, CN(R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
- 10 R¹¹OC(O)NR¹⁰-,
- c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocyclic, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-,
- 15 R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, and R¹¹OC(O)-NR¹⁰-;

20 R² and R³ are independently selected from: H; unsubstituted or substituted C₁-8 alkyl, unsubstituted or substituted C₂-8 alkenyl, unsubstituted or substituted C₂-8 alkynyl, unsubstituted or substituted aryl,

unsubstituted or substituted heterocycle,











wherein the substituted group is substituted with one or more of:

- 1) aryl or heterocycle, unsubstituted or substituted with:
 - 25 a) C₁-4 alkyl,
 - b) (CH₂)_pOR⁶,

- 13 -

5

- c) $(\text{CH}_2)_p\text{NR}^6\text{R}^7$,
 d) halogen,
 e) CN,
 2) C₃₋₆ cycloalkyl,
 3) OR^6 ,
 4) SR^{6a} , S(O)R^{6a} , SO_2R^{6a} ,



10

- 14 -



R^2 and R^3 are attached to the same C atom and are combined to form
 - $(CH_2)_u$ - wherein one of the carbon atoms is optionally replaced by a
 5 moiety selected from: O, $S(O)_m$, $-NC(O)-$, and $-N(COR^{10})-$;



R^4 is selected from H and CH_3 ;

10 and any two of R^2 , R^3 and R^4 are optionally attached to the same
 carbon atom;


R^6 , R^7 and R^{7a} are independently selected from: H; C1-4 alkyl, C3-6
 cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl,
 heteroarylsulfonyl, unsubstituted or substituted with:

15

- a) C1-4 alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,



20




R^6 and R^7 may be joined in a ring;
 R^7 and R^{7a} may be joined in a ring;

- 15 -

R^{6a} is selected from: C₁₋₄ alkyl, C₃₋₆ cycloalkyl, heterocycle, aryl, unsubstituted or substituted with:

- a) C₁₋₄ alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,



- f) $-\text{SO}_2\text{R}^{11}$, or
- g) $\text{N}(\text{R}^{10})_2$;

R⁸ is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and
- c) C₁₋₆ alkyl unsubstituted or substituted by aryl, cyanophenyl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NH-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹⁰OC(O)NH-;

R⁹ is selected from:

- a) hydrogen,
- b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and

- 16 -

- c) C₁-C₆ alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

5

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

10

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR¹⁰-, -NR¹⁰C(O)-, O, -N(R¹⁰)-, -S(O)₂N(R¹⁰)-, -N(R¹⁰)S(O)₂-, or S(O)_m;

15

G is H₂ or O;

V is selected from:

20

- a) hydrogen,
- b) heterocycle,
- c) aryl,
- d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C₂-C₂₀ alkenyl,

25

provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

W is a heterocycle;

30

X is -CH₂-, -C(=O)-, or -S(=O)_m-;

Z is

a unsubstituted or substituted group selected from aryl, heteroaryl, arylmethyl, heteroarylmethyl, arylsulfonyl,

- 17 -

heteroarylsulfonyl, wherein the substituted group is substituted with one or more of the following:

1) C₁₋₄ alkyl, unsubstituted or substituted with:

- a) C₁₋₄ alkoxy,
- b) NR⁶R⁷,
- c) C₃₋₆ cycloalkyl,
- d) aryl or heterocycle,
- e) HO,
- f) -S(O)_mR^{6a}, or
- g) -C(O)NR⁶R⁷,

2) aryl or heterocycle,

3) halogen,

4) OR⁶,

5) NR⁶R⁷,

6) CN,

7) NO₂,

8) CF₃;

9) -S(O)_mR^{6a},

10) -C(O)NR⁶R⁷, or

11) C₃₋₆ cycloalkyl;

m is 0, 1 or 2;

n is 0, 1, 2, 3 or 4;

p is 0, 1, 2, 3 or 4;

25 q is 1 or 2;

r is 0 to 5, provided that r is 0 when V is hydrogen;

s is 0 or 1;


t is 0 or 1; and

u is 4 or 5;

30 or the pharmaceutically acceptable salts thereof.

In a third embodiment of this invention, the inhibitors of farnesyl-protein transferase are illustrated by the formula C:

- 18 -




wherein:

R^{1a} and R^{1b} are independently selected from:

- 5 a) hydrogen,
- b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-,
- 10 c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocyclic, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, and R¹¹OC(O)NR¹⁰-;
- 15

R² and R³ are independently selected from: H; unsubstituted or substituted C₁-8 alkyl, unsubstituted or substituted C₂-8 alkenyl, unsubstituted or substituted C₂-8 alkynyl, unsubstituted or substituted aryl,



unsubstituted or substituted heterocycle,

wherein the substituted group is substituted with one or more of:







- 1) aryl or heterocycle, unsubstituted or substituted with:
 - 25 a) C₁-4 alkyl,
 - b) (CH₂)_pOR⁶,
 - c) (CH₂)_pNR⁶R⁷,
 - d) halogen,

- 19 -

e) CN,

2) C₃-6 cycloalkyl,3) OR⁶,4) SR^{6a}, S(O)R^{6a}, SO₂R^{6a},

5

5) $\text{—NR}^6\text{R}^7$ 11) $\text{—SO}_2\text{—NR}^6\text{R}^7$ 

- 20 -





R^2 and R^3 are attached to the same C atom and are combined to form
 - $(CH_2)_u$ - wherein one of the carbon atoms is optionally replaced by a
 5 moiety selected from: O, S(O)_m, -NC(O)-, and -N(COR¹⁰)-;

R^4 is selected from H and CH₃;

10 and any two of R^2 , R^3 and R^4 are optionally attached to the same
 carbon atom;

R^6 , R^7 and R^{7a} are independently selected from: H; C₁₋₄ alkyl, C₃₋₆
 cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl,
 heteroarylsulfonyl, unsubstituted or substituted with:

- 15 a) C₁₋₄ alkoxy,
 b) aryl or heterocycle,
 c) halogen,
 d) HO,




20 g) $N(R^{10})_2$; or

R^6 and R^7 may be joined in a ring;
 R^7 and R^{7a} may be joined in a ring;

- 21 -

R^{6a} is selected from: C₁₋₄ alkyl, C₃₋₆ cycloalkyl, heterocycle, aryl, unsubstituted or substituted with:

- a) C₁₋₄ alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,



- f) $-\text{SO}_2\text{R}^{11}$, or
- g) $\text{N}(\text{R}^{10})_2$;

R⁸ is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and
- c) C₁₋₆ alkyl unsubstituted or substituted by aryl, cyanophenyl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NH-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹⁰OC(O)NH-;

R⁹ is selected from:

- a) hydrogen,
- b) C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and

- 22 -

- c) C₁-C₆ alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

5 R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

10 A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR¹⁰-, -NR¹⁰C(O)-, O, -N(R¹⁰)-, -S(O)₂N(R¹⁰)-, -N(R¹⁰)S(O)₂-, or S(O)_m;

15 G is O;

V is selected from:

- 20 a) hydrogen,
b) heterocycle,
c) aryl,
d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
e) C₂-C₂₀ alkenyl,
25 provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

W is a heterocycle;

30 X is -CH₂-, -C(=O)-, or -S(=O)_m-;

Z is a unsubstituted or substituted group selected from aryl, heteroaryl, arylmethyl, heteroarylmethyl, arylsulfonyl,

- 23 -

heteroarylsulfonyl, wherein the substituted group is substituted with one or more of the following:


- 1) C₁₋₄ alkyl, unsubstituted or substituted with:
 - a) C₁₋₄ alkoxy,
 - b) NR⁶R⁷,
 - c) C₃₋₆ cycloalkyl,
 - d) aryl or heterocycle,
 - e) HO,
 - f) -S(O)_mR^{6a}, or
 - g) -C(O)NR⁶R⁷,
- 2) aryl or heterocycle,
- 3) halogen,
- 4) OR⁶,
- 5) NR⁶R⁷,
- 6) CN,
- 7) NO₂,
- 8) CF₃;
- 9) -S(O)_mR^{6a},
- 10) -C(O)NR⁶R⁷, or
- 11) C₃₋₆ cycloalkyl;

- m is 0, 1 or 2;
- n is 0, 1, 2, 3 or 4;
- p is 0, 1, 2, 3 or 4;
- q is 1 or 2;
- r is 0 to 5, provided that r is 0 when V is hydrogen;
- s is 1;
- t is 0 or 1; and
- u is 4 or 5;

or the pharmaceutically acceptable salts thereof.

A preferred embodiment of the compounds of this invention is illustrated by the following formula:

- 24 -



wherein:

5 R^{1a} is independently selected from: hydrogen or C₁-C₆ alkyl;

R^{1b} is independently selected from:

- 10 a) hydrogen,
 b) aryl, heterocycle, cycloalkyl, R¹⁰O-, -N(R¹⁰)₂ or C₂-C₆ alkenyl,
 c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocycle, cycloalkyl, alkenyl, R¹⁰O- and -N(R¹⁰)₂;

15 R³, R⁴ and R⁵ are independently selected from H and CH₃;

R² is H; $\begin{array}{c} \text{NR}^6\text{R}^7 \\ | \\ \text{O} \end{array}$; or C₁-5 alkyl, unbranched or branched, unsubstituted or substituted with one or more of:

- 20 1) aryl,
 2) heterocycle,
 3) OR⁶,
 4) SR^{6a}, SO₂R^{6a}, or
 5) $\begin{array}{c} \text{NR}^6\text{R}^7 \\ | \\ \text{O} \end{array}$;

25 and any two of R², R³, R⁴, and R⁵ are optionally attached to the same carbon atom;

- 25 -

R⁶, R⁷ and R^{7a} are independently selected from:

H; C₁₋₄ alkyl, C₃₋₆ cycloalkyl, aryl, heterocycle,
unsubstituted or substituted with:

- 5 a) C₁₋₄ alkoxy,
 b) halogen, or
 c) aryl or heterocycle;

R^{6a} is selected from:

- 10 C₁₋₄ alkyl or C₃₋₆ cycloalkyl,
 unsubstituted or substituted with:
 a) C₁₋₄ alkoxy,
 b) halogen, or
 c) aryl or heterocycle;

15 R⁸ is independently selected from:

- a) hydrogen,
 b) C_{1-C6} alkyl, C_{2-C6} alkenyl, C_{2-C6} alkynyl, C_{1-C6}
 perfluoroalkyl, F, Cl, R¹⁰O-, R¹⁰C(O)NR¹⁰-, CN, NO₂,
 (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, -N(R¹⁰)₂, or
20 R¹¹OC(O)NR¹⁰-, and
 c) C_{1-C6} alkyl substituted by C_{1-C6} perfluoroalkyl, R¹⁰O-,
 R¹⁰C(O)NR¹⁰-, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-,
 R¹⁰OC(O)-, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

25 R⁹ is selected from:

- a) hydrogen,
 b) C_{2-C6} alkenyl, C_{2-C6} alkynyl, C_{1-C6} perfluoroalkyl, F,
 Cl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, CN, NO₂,
 (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, -N(R¹⁰)₂, or
30 R¹¹OC(O)NR¹⁰-, and
 c) C_{1-C6} alkyl unsubstituted or substituted by C_{1-C6}
 perfluoroalkyl, F, Cl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-,
 CN, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-,
 -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

- 26 -

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and aryl;

5 R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR¹⁰-, O, -N(R¹⁰)-, or S(O)_m;

10 V is selected from:

- a) hydrogen,
- b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,
- 15 c) aryl,
- d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C₂-C₂₀ alkenyl, and

20 provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or
25 isoquinolinyl;

X is -CH₂- or -C(=O)-;

30 Y is mono- or bicyclic aryl, or mono- or bicyclic heterocycle, unsubstituted or substituted with one or more of:


- a) C₁₋₄ alkyl,
- b) C₁₋₄ alkoxy,
- c) halogen, or
- d) NR⁶R⁷;

- 27 -

m is 0, 1 or 2;
 n is 0, 1, 2, 3 or 4;
 p is 0, 1, 2, 3 or 4;
 5 r is 0 to 5, provided that r is 0 when V is hydrogen;
 s is 0 or 1; and
 t is 0 or 1;

or the pharmaceutically acceptable salts thereof.

In a second preferred embodiment of this invention, the inhibitors of farnesyl-protein transferase are illustrated by the formula B:



wherein:


R^{1a} is independently selected from: hydrogen or C₁-C₆ alkyl;


20 R^{1b} is independently selected from:

- a) hydrogen,
b) aryl, heterocycle, cycloalkyl, $R^{10}O-$, $-N(R^{10})_2$ or C_2-C_6 alkenyl,
c) unsubstituted or substituted C_1-C_6 alkyl wherein the substituent on the substituted C_1-C_6 alkyl is selected from unsubstituted or substituted aryl, heterocycle, cycloalkyl, alkenyl, $R^{10}O-$ and $-N(R^{10})_2$;

- 28 -

R³ and R⁴ are independently selected from H and CH₃;

R² is H; ; or C₁-5 alkyl, unbranched or branched, unsubstituted or substituted with one or more of:

- 1) aryl,
- 2) heterocycle,
- 3) OR⁶,
- 4) SR^{6a}, SO₂R^{6a}, or
- 5) ;

and any two of R², R³, R⁴, and R⁵ are optionally attached to the same carbon atom;

R⁶, R⁷ and R^{7a} are independently selected from:

H; C₁-4 alkyl, C₃-6 cycloalkyl, aryl, heterocycle, unsubstituted or substituted with:

- a) C₁-4 alkoxy,
- b) halogen, or
- c) aryl or heterocycle;

R^{6a} is selected from:

C₁-4 alkyl or C₃-6 cycloalkyl, unsubstituted or substituted with:

- a) C₁-4 alkoxy,
- b) halogen, or
- c) aryl or heterocycle;

R⁸ is independently selected from:

- a) hydrogen,
- b) C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ perfluoroalkyl, F, Cl, R¹⁰O-, R¹⁰C(O)NR¹⁰-, CN, NO₂, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and

- 29 -

- c) C₁-C₆ alkyl substituted by C₁-C₆ perfluoroalkyl, R¹⁰O-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

5 R⁹ is selected from:

- a) hydrogen,
b) C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ perfluoroalkyl, F, Cl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, CN, NO₂, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and
10 c) C₁-C₆ alkyl unsubstituted or substituted by C₁-C₆ perfluoroalkyl, F, Cl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, CN, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

15

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

20

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR¹⁰-, O, -N(R¹⁰)-, or S(O)_m;

V is selected from:

25

- a) hydrogen,
b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,
c) aryl,
30 d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
e) C₂-C₂₀ alkenyl, and

- 30 -

provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

G is H₂ or O;

5

W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or isoquinolinyl;

10 X is -CH₂- or -C(=O)-;

Z is mono- or bicyclic aryl, mono- or bicyclic heteroaryl, mono- or bicyclic arylmethyl, mono- or bicyclic heteroarylmethyl, mono- or bicyclic arylsulfonyl, mono- or bicyclic heteroarylsulfonyl, unsubstituted or substituted with one or two of the following:

15

1) C₁₋₄ alkyl, unsubstituted or substituted with:

20

- a) C₁₋₄ alkoxy,
- b) NR⁶R⁷,
- c) C₃₋₆ cycloalkyl,
- d) aryl or heterocycle,
- e) HO,
- f) -S(O)_mR⁶, or
- g) -C(O)NR⁶R⁷,

25

2) aryl or heterocycle,

3) halogen,

4) OR⁶,

5) NR⁶R⁷,

6) CN,

30

7) NO₂,

8) CF₃;

9) -S(O)_mR⁶,

10) -C(O)NR⁶R⁷, or

11) C₃₋₆ cycloalkyl;

- 31 -

- m is 0, 1 or 2;
 n is 0, 1, 2, 3 or 4;
 p is 0, 1, 2, 3 or 4;
 5 r is 0 to 5, provided that r is 0 when V is hydrogen;
 s is 0 or 1;
 t is 0 or 1; and
 u is 4 or 5;

- 10 provided that when G is H₂ and W is imidazolyl, then the substituent (R⁸)_r-V-A¹(CR^{1a}₂)_nA²(CR^{1a}₂)_n - is not H and

provided that when X is -C(=O)-, or -S(=O)_m-, then t is 1 and the substituent (R⁸)_r-V-A¹(CR^{1a}₂)_nA²(CR^{1a}₂)_n - is not H;

15

or the pharmaceutically acceptable salts thereof.

The preferred compounds of this invention are as follows:

- 20 2(S)-Butyl-1-(2,3-diaminoprop-1-yl)-4-(1-naphthoyl)-piperazine

1-(3-Amino-2-(2-naphthylmethylamino)prop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine

25

2(S)-Butyl-1-{5-[1-(2-naphthylmethyl)]-4,5-dihydroimidazol}methyl-4-(1-naphthoyl)piperazine

1-[5-(1-Benzylimidazol)methyl]-2(S)-butyl-4-(1-naphthoyl)piperazine

30

1-{5-[1-(4-Nitrobenzyl)imidazolyl]methyl}-2(S)-butyl-4-(1-naphthoyl)piperazine

1-(3-Acetamidomethylthio-2(R)-aminoprop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine

35

- 32 -

2(S)-Butyl-1-[2-(1-imidazolyl)ethyl]sulfonyl-4-(1-naphthoyl)piperazine

2(R)-Butyl-1-imidazolyl-4-methyl-4-(1-naphthoyl)piperazine

5

2(S)-Butyl-4-(1-naphthoyl)-1-(3-pyridylmethyl)piperazine

1-2(S)-butyl-(2(R)-(4-nitrobenzyl)amino-3-hydroxypropyl)-4-(1-naphthoyl)piperazine

10

1-(2(R)-Amino-3-hydroxyheptadecyl)-2(S)-butyl-4-(1-naphthoyl)piperazine

2(S)-Benzyl-1-imidazolyl-4-methyl-4-(1-naphthoyl)piperazine

15

1-(2(R)-Amino-3-(3-benzylthio)propyl)-2(S)-butyl-4-(1-naphthoyl)-piperazine

1-(2(R)-Amino-3-[3-(4-nitrobenzylthio)propyl]))-2(S)-butyl-4-(1-naphthoyl)piperazine

20

2(S)-Butyl-1-[(4-imidazolyl)ethyl]-4-(1-naphthoyl)piperazine

2(S)-Butyl-1-[(4-imidazolyl)methyl]-4-(1-naphthoyl)piperazine

25

2(S)-Butyl-1-[(1-naphth-2-ylmethyl)-1H-imidazol-5-yl)acetyl]-4-(1-naphthoyl)piperazine

2(S)-Butyl-1-[(1-naphth-2-ylmethyl)-1H-imidazol-5-yl)ethyl]-4-(1-naphthoyl)piperazine

30

1-(2(R)-Amino-3-hydroxypropyl)-2(S)-butyl-4-(1-naphthoyl)piperazine

35 1-(2(R)-Amino-4-hydroxybutyl)-2(S)-butyl-4-(1-naphthoyl)piperazine

- 33 -

- 1-(2-Amino-3-(2-benzyloxyphenyl)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine
- 5 1-(2-Amino-3-(2-hydroxyphenyl)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine
- 1-[3-(4-imidazolyl)propyl]-2(S)-butyl-4-(1-naphthoyl)piperazine
- 10 2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(1-naphthylmethyl)imidazol-5-ylmethyl]-piperazine
- 2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(2-naphthylmethyl)imidazol-5-ylmethyl]-piperazine
- 15 2(S)-*n*-Butyl-1-[1-(4-cyanobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine
- 2(S)-*n*-Butyl-1-[1-(4-methoxybenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine
- 20 2(S)-*n*-Butyl-1-[1-(3-methyl-2-butenyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine
- 25 2(S)-*n*-Butyl-1-[1-(4-fluorobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine
- 2(S)-*n*-Butyl-1-[1-(4-chlorobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine
- 30 1-[1-(4-Bromobenzyl)imidazol-5-ylmethyl]-2(S)-*n*-butyl-4-(1-naphthoyl)piperazine
- 1-[1-(4-Bromobenzyl)imidazol-5-ylmethyl]-2(S)-*n*-butyl-4-(1-naphthoyl)piperazine
- 35

- 34 -

2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(4-trifluoromethylbenzyl)imidazol-5-ylmethyl]-piperazine

5 2(S)-*n*-Butyl-1-[1-(4-methylbenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)-piperazine

2(S)-*n*-Butyl-1-[1-(3-methylbenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)-piperazine

10 1-[1-(4-Phenylbenzyl)imidazol-5-ylmethyl]-2(S)-*n*-butyl-4-(1-naphthoyl)-piperazine

15 2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(2-phenylethyl)imidazol-5-ylmethyl]-piperazine

2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(4-trifluoromethoxy)-imidazol-5-ylmethyl]piperazine

20 1-[[1-(4-cyanobenzyl)-1H-imidazol-5-yl]acetyl]-2(S)-*n*-butyl-4-(1-naphthoyl)piperazine

5(S)-*n*-Butyl-1-(2,3-dimethylphenyl)-4-(4-imidazolylmethyl)-piperazin-2-one

25 5(S)-*n*-Butyl-4-[1-(4-cyanobenzyl)imidazol-5-ylmethyl]-1-(2,3-dimethylphenyl)piperazin-2-one

30 4-[1-(4-Cyanobenzyl)imidazol-5-ylmethyl]-1-(2,3-dimethylphenyl)-5(S)-(2-methoxyethyl)piperazin-2-one

(S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(methanesulfonyl)ethyl]-2-piperazinone

35 (S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)ethyl]-2-piperazinone

- 35 -

(S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)methyl]-2-piperazinone

5 (S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[N-ethyl-2-acetamido]-2-piperazinone

(±)-5-(2-Butynyl)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone

10 1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone

15 5(S)-Butyl-4-[1-(4-cyanobenzyl-2-methyl)-5-imidazolylmethyl]-1-(2,3-dimethylphenyl)-piperazin-2-one

4-[1-(2-(4-Cyanophenyl)-2-propyl)-5-imidazolylmethyl]-1-(3-chlorophenyl)-5(S)-(2-methylsulfonylethyl)piperazin-2-one

20 5(S)-*n*-Butyl-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-1-(2-methylphenyl)piperazin-2-one

4-[1-(4-Cyanobenzyl)-5-imidazolylmethyl]-5(S)-(2-fluoroethyl)-1-(3-chlorophenyl)piperazin-2-one

25 4-[3-(4-Cyanobenzyl)pyridin-4-yl]-1-(3-chlorophenyl)-5(S)-(2-methylsulfonylethyl)-piperazin-2-one

4-[5-(4-Cyanobenzyl)-1-imidazolylethyl]-1-(3-chlorophenyl)piperazin-2-one.


30

or the pharmaceutically acceptable salts thereof.


Specific examples of the compounds of the invention are:

35 1-{5-[1-(4-Nitrobenzyl)imidazolyl]methyl}-2(S)-butyl-4-(1-naphthoyl)piperazine

- 36 -




1-[5-(1-Benzylimidazol)methyl]-2(S)-butyl-4-(1-naphthoyl)piperazine




5

1-(2(R)-Amino-3-(3-benzylthio)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine




10 1-(2(R)-Amino-3-[3-(4-nitrobenzylthio)propyl])-2(S)-butyl-4-(1-naphthoyl)piperazine

- 37 -




2(S)-*n*-Butyl-1-[1-(4-cyanobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine

5




2(S)-*n*-Butyl-1-[1-(4-cyanobenzyl)imidazol-5-ylmethyl]-4-(2,3-dimethylphenyl)piperazin-5-one

10




- 38 -

2(S)-*n*-Butyl-1-[1-(4-chlorobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine



5


1-[[1-(4-Cyanobenzyl)-1H-imidazol-5-yl]acetyl]-2(S)-*n*-butyl-4-(1-naphthoyl)piperazine



10


1-[1-(4-Cyanobenzyl)imidazol-5-ylmethyl]-4-(2,3-dimethylphenyl)-2(S)-(2-methoxyethyl)piperazin-5-one

- 39 -




5(S)-*n*-Butyl-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-1-(2-methylphenyl)piperazin-2-one

5




10 (S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(methanesulfonyl)ethyl]-2-piperazinone




- 40 -

(S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)ethyl]-2-piperazinone




5

(S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)methyl]-2-piperazinone



10

1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolyl-methyl]-2-piperazinone



15

- 41 -

or the pharmaceutically acceptable salts thereof.

The compounds of the present invention may have asymmetric centers and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers, including optical isomers, being included in the present invention. When any variable (e.g. aryl, heterocycle, R¹, R² etc.) occurs more than one time in any constituent, its definition on each occurrence is independent at every other occurrence. Also, combinations of substituents/or variables are permissible only if such combinations result in stable compounds.

As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms; "alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge. "Halogen" or "halo" as used herein means fluoro, chloro, bromo and iodo.

As used herein, "aryl" is intended to mean any stable monocyclic or bicyclic carbon ring of up to 7 members in each ring, wherein at least one ring is aromatic. Examples of such aryl elements include phenyl, naphthyl, tetrahydronaphthyl, indanyl, biphenyl, phenanthryl, anthryl or acenaphthyl.

The term heterocycle or heterocyclic, as used herein, represents a stable 5- to 7-membered monocyclic or stable 8- to 11-membered bicyclic heterocyclic ring which is either saturated or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from the group consisting of N, O, and S, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heterocyclic elements include, but are not limited to, azepinyl, benzimidazolyl, benzisoxazolyl, benzofurazanyl, benzopyranyl, benzothiopyranyl, benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, cinnoliny, dihydrobenzofuryl, dihydrobenzothieryl, dihydrobenzothiopyranyl,

- 42 -

5 dihydrobenzothiopyranyl sulfone, furyl, imidazolidinyl, imidazoliny, imidazolyl, indolinyl, indolyl, isochromanyl, isoindolinyl, isoquinolinyl, isothiazolidinyl, isothiazolyl, isothiazolidinyl, morpholinyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, 2-oxopiperazinyl,
10 2-oxopiperdiny, 2-oxopyrrolidinyl, piperidyl, piperazinyl, pyridyl, pyrazinyl, pyrazolidinyl, pyrazolyl, pyridazinyl, pyrimidinyl, pyrrolidinyl, pyrrolyl, quinazolinyl, quinolinyl, quinoxaliny, tetrahydrofuryl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiazolyl, thiazolinyl,
15 thienofuryl, thienothienyl, and thienyl.

As used herein, "heteroaryl" is intended to mean any stable monocyclic or bicyclic carbon ring of up to 7 members in each ring, wherein at least one ring is aromatic and wherein from one to four carbon atoms are replaced by heteroatoms selected from the group
15 consisting of N, O, and S. Examples of such heterocyclic elements include, but are not limited to, benzimidazolyl, benzisoxazolyl, benzofurazanyl, benzopyranyl, benzothiopyranyl, benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, cinnolinyl, dihydrobenzofuryl, dihydrobenzothienyl, dihydrobenzothiopyranyl,
20 dihydrobenzothiopyranyl sulfone, furyl, imidazolyl, indolinyl, indolyl, isochromanyl, isoindolinyl, isoquinolinyl, isothiazolyl, naphthyridinyl, oxadiazolyl, pyridyl, pyrazinyl, pyrazolyl, pyridazinyl, pyrimidinyl, pyrrolyl, quinazolinyl, quinolinyl, quinoxaliny, tetrahydroisoquinolinyl, tetrahydroquinolinyl, thiazolyl, thienofuryl,
25 thienothienyl, and thienyl.

As used herein in the definition of R² and R³, the term "the substituted group" intended to mean a substituted C₁₋₈ alkyl, substituted C₂₋₈ alkenyl, substituted C₂₋₈ alkynyl, substituted aryl or substituted heterocycle from which the substituent(s) R² and R³ are selected.

30 As used herein in the definition of R⁶, R⁷ and R^{7a}, the substituted C₁₋₈ alkyl, substituted C₃₋₆ alkenyl, substituted aroyl, substituted aryl, substituted heteroaroyl, substituted arylsulfonyl, substituted heteroarylsulfonyl and substituted heterocycle include

- 43 -


moieties containing from 1 to 3 substituents in addition to the point of attachment to the rest of the compound.

As used herein, the term "substituted aryl" is intended to include the aryl group which is substituted with 1 or 2 substituents
 5 selected from the group which includes but is not limited to F, Cl, Br, CF₃, NH₂, N(C₁-C₆ alkyl)₂, NO₂, CN, (C₁-C₆ alkyl)O-, -OH, (C₁-C₆ alkyl)S(O)_m-, (C₁-C₆ alkyl)C(O)NH-, H₂N-C(NH)-, (C₁-C₆ alkyl)C(O)-, (C₁-C₆ alkyl)OC(O)-, N₃, (C₁-C₆ alkyl)OC(O)NH- and C₁-C₂₀ alkyl.

10 When R² and R³ are combined to form - (CH₂)_u -, cyclic moieties are formed. Examples of such cyclic moieties include, but are not limited to:



15 In addition, such cyclic moieties may optionally include a heteroatom(s). Examples of such heteroatom-containing cyclic moieties include, but are not limited to:




20 Lines drawn into the ring systems from substituents (such as from R², R³, R⁴ etc.) indicate that the indicated bond may be attached to any of the substitutable ring carbon atoms.

- 44 -

Preferably, R^{1a} and R^{1b} are independently selected from:
 hydrogen, $-N(R^{10})_2$, $R^{10}C(O)NR^{10}$ - or unsubstituted or substituted
 C_1 - C_6 alkyl wherein the substituent on the substituted C_1 - C_6 alkyl is
 selected from unsubstituted or substituted phenyl, $-N(R^{10})_2$, $R^{10}O$ - and
 5 $R^{10}C(O)NR^{10}$.




Preferably, R^2 is selected from: H,








and an unsubstituted or substituted group, the
 group selected from C_1 -8 alkyl, C_2 -8 alkenyl and C_2 -8 alkynyl;

wherein the substituted group is substituted with one or more of:

- 10 1) aryl or heterocycle, unsubstituted or substituted with:
 a) C_1 -4 alkyl,
 b) $(CH_2)_pOR^6$,
 c) $(CH_2)_pNR^6R^7$,
 d) halogen,
 15 2) C_3 -6 cycloalkyl,
 3) OR^6 ,
 4) SR^{6a} , $S(O)R^{6a}$, SO_2R^{6a} ,



- 45 -



5 Preferably, R^3 is selected from: hydrogen and C1-C6 alkyl.
Preferably, R^4 and R^5 are hydrogen.

Preferably, R^6 , R^7 and R^{7a} is selected from: hydrogen, unsubstituted or substituted C1-C6 alkyl, unsubstituted or substituted aryl and unsubstituted or substituted cycloalkyl.

10 Preferably, R^{6a} is unsubstituted or substituted C1-C6 alkyl, unsubstituted or substituted aryl and unsubstituted or substituted cycloalkyl.

Preferably, R^9 is hydrogen or methyl. Most preferably, R^a is hydrogen.

- 46 -

Preferably, R^{10} is selected from H, C_1 - C_6 alkyl and benzyl.

Preferably, A^1 and A^2 are independently selected from: a bond, $-C(O)NR^{10}$ -, $-NR^{10}C(O)$ -, O, $-N(R^{10})$ -, $-S(O)_2N(R^{10})$ - and-
 5 $N(R^{10})S(O)_2$ -.

Preferably, V is selected from hydrogen, heterocycle and aryl. More preferably, V is phenyl.

Preferably, Y is selected from unsubstituted or substituted phenyl, unsubstituted or substituted naphthyl, unsubstituted or
 10 substituted pyridyl, unsubstituted or substituted furanyl and unsubstituted or substituted thienyl. More preferably, Y is unsubstituted or substituted phenyl.

Preferably, Z is selected from unsubstituted or substituted phenyl, unsubstituted or substituted naphthyl, unsubstituted or
 15 substituted pyridyl, unsubstituted or substituted furanyl and unsubstituted or substituted thienyl. More preferably, Z is unsubstituted or substituted phenyl.

Preferably, W is selected from imidazolyl, imidazolyl, oxazolyl, pyrazolyl, pyrrolidinyl, thiazolyl and pyridyl. More
 20 preferably, W is selected from imidazolyl and pyridyl.

Preferably, n and r are independently 0, 1, or 2.

Preferably p is 1, 2 or 3.

Preferably s is 0.

Preferably t is 1.

25 Preferably, in compounds of the formula B, when G is H_2 and W is imidazolyl, then the substituent $(R^8)_r$ - V - $A^1(CR^{1a}_2)_nA^2(CR^{1a}_2)_n$ - is not H.

Preferably, in compounds of the formula B, when X is $-C(=O)$ -, or $-S(=O)_m$ -, then t is 1 and the substituent $(R^8)_r$ - V -
 30 $A^1(CR^{1a}_2)_nA^2(CR^{1a}_2)_n$ - is not H;

It is intended that the definition of any substituent or variable (e.g., R^{1a} , R^9 , n, etc.) at a particular location in a molecule be independent of its definitions elsewhere in that molecule. Thus,

- 47 -

-N(R¹⁰)₂ represents -NHH, -NHCH₃, -NHC₂H₅, etc. It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials.

The pharmaceutically acceptable salts of the compounds of this invention include the conventional non-toxic salts of the compounds of this invention as formed, e.g., from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxy-benzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic and the like.

The pharmaceutically acceptable salts of the compounds of this invention can be synthesized from the compounds of this invention which contain a basic moiety by conventional chemical methods. Generally, the salts are prepared either by ion exchange chromatography or by reacting the free base with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various combinations of solvents.

Reactions used to generate the compounds of this invention are prepared by employing reactions as shown in the Schemes 1-22, in addition to other standard manipulations such as ester hydrolysis, cleavage of protecting groups, etc., as may be known in the literature or exemplified in the experimental procedures. Substituents R, R^a and R^b, as shown in the Schemes, represent the substituents R², R³, R⁴, and R⁵; however their point of attachment to the ring is illustrative only and is not meant to be limiting.

These reactions may be employed in a linear sequence to provide the compounds of the invention or they may be used to

- 48 -

synthesize fragments which are subsequently joined by the alkylation reactions described in the Schemes.

Synopsis of Schemes 1-22:

5 The requisite intermediates are in some cases commercially available, or can be prepared according to literature procedures, for the most part. In Scheme 1, for example, the synthesis of 2-alkyl substituted piperazines is outlined, and is essentially that described by J. S. Kiely and S. R. Priebe in Organic Preparations and Proceedings Int.,
10 1990, 22, 761-768. Boc-protected amino acids I, available commercially or by procedures known to those skilled in the art, can be coupled to N-benzyl amino acid esters using a variety of dehydrating agents such as DCC (dicyclohexycarbodiimide) or EDC-HCl (1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride) in a solvent such
15 as methylene chloride, chloroform, dichloroethane, or in dimethylformamide. The product II is then deprotected with acid, for example hydrogen chloride in chloroform or ethyl acetate, or trifluoroacetic acid in methylene chloride, and cyclized under weakly basic conditions to give the diketopiperazine III. Reduction of III with
20 lithium aluminum hydride in refluxing ether gives the piperazine IV, which is protected as the Boc derivative V. The N-benzyl group can be cleaved under standard conditions of hydrogenation, e.g., 10% palladium on carbon at 60 psi hydrogen on a Parr apparatus for 24-48 h. The product VI can be treated with an acid chloride, or a
25 carboxylic acid under standard dehydrating conditions to furnish the carboxamides VII; a final acid deprotection as previously described gives the intermediate VIII (Scheme 2). The intermediate VIII can be reductively alkylated with a variety of aldehydes, such as IX. The aldehydes can be prepared by standard procedures, such as that
30 described by O. P. Goel, U. Krolls, M. Stier and S. Kesten in Organic Syntheses, 1988, 67, 69-75, from the appropriate amino acid (Scheme 3). The reductive alkylation can be accomplished at pH 5-7 with a variety of reducing agents, such as sodium triacetoxyborohydride or sodium cyanoborohydride in a solvent such as dichloroethane, methanol

- 49 -

or dimethylformamide. The product X can be deprotected to give the final compounds XI with trifluoroacetic acid in methylene chloride. The final product XI is isolated in the salt form, for example, as a trifluoroacetate, hydrochloride or acetate salt, among others. The product diamine XI can further be selectively protected to obtain XII, which can subsequently be reductively alkylated with a second aldehyde to obtain XIII. Removal of the protecting group, and conversion to cyclized products such as the dihydroimidazole XV can be accomplished by literature procedures.

Alternatively, the protected piperazine intermediate VII can be reductively alkylated with other aldehydes such as 1-trityl-4-imidazolyl-carboxaldehyde or 1-trityl-4-imidazolylacetaldehyde, to give products such as XVI (Scheme 4). The trityl protecting group can be removed from XVI to give XVII, or alternatively, XVI can first be treated with an alkyl halide then subsequently deprotected to give the alkylated imidazole XVIII. Alternatively, the intermediate VIII can be acylated or sulfonylated by standard techniques. The imidazole acetic acid XIX can be converted to the acetate XXI by standard procedures, and XXI can be first reacted with an alkyl halide, then treated with refluxing methanol to provide the regiospecifically alkylated imidazole acetic acid ester XXII. Hydrolysis and reaction with piperazine VIII in the presence of condensing reagents such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC) leads to acylated products such as XXIV.

If the piperazine VIII is reductively alkylated with an aldehyde which also has a protected hydroxyl group, such as XXV in Scheme 6, the protecting groups can be subsequently removed to unmask the hydroxyl group (Schemes 6, 7). The alcohol can be oxidized under standard conditions to *e.g.* an aldehyde, which can then be reacted with a variety of organometallic reagents such as Grignard reagents, to obtain secondary alcohols such as XXIX. In addition, the fully deprotected amino alcohol XXX can be reductively alkylated (under conditions described previously) with a variety of aldehydes to obtain secondary amines, such as XXXI (Scheme 7), or tertiary amines.

- 50 -

The Boc protected amino alcohol XXVII can also be utilized to synthesize 2-aziridinylmethylpiperazines such as XXXII (Scheme 8). Treating XXVII with 1,1'-sulfonyldiimidazole and sodium hydride in a solvent such as dimethylformamide led to the formation of aziridine XXXII. The aziridine reacted in the presence of a nucleophile, such as a thiol, in the presence of base to yield the ring-opened product XXXIII.

In addition, the piperazine VIII can be reacted with aldehydes derived from amino acids such as O-alkylated tyrosines, according to standard procedures, to obtain compounds such as XXXIX. When R' is an aryl group, XXXIX can first be hydrogenated to unmask the phenol, and the amine group deprotected with acid to produce XL. Alternatively, the amine protecting group in XXXIX can be removed, and O-alkylated phenolic amines such as XLI produced.

Depending on the identity of the amino acid I, various side chains can be incorporated into the piperazine. For example when I is the Boc-protected β -benzyl ester of aspartic acid, the intermediate diketopiperazine XLII where $n=1$ and $R=\text{benzyl}$ is obtained, as shown in Scheme 10. Subsequent lithium aluminum hydride reduction reduces the ester to the alcohol XLIII, which can then be reacted with a variety of alkylating agents such as an alkyl iodide, under basic conditions, for example, sodium hydride in dimethylformamide or tetrahydrofuran. The resulting ether XLIV can then be carried on to final products as described in Schemes 3-9.

N-Aryl piperazines can be prepared as described in Scheme 11. An aryl amine XLV is reacted with *bis*-chloroethyl amine hydrochloride (XLVI) in refluxing *n*-butanol to furnish compounds XLVII. The resulting piperazines XLVII can then be carried on to final products as described in Schemes 3-9.

Piperazin-5-ones can be prepared as shown in Scheme 12. Reductive amination of Boc-protected amino aldehydes XLIX (prepared from I as described previously) gives rise to compound L. This is then reacted with bromoacetyl bromide under Schotten-Baumann conditions; ring closure is effected with a base such as sodium hydride

- 51 -

in a polar aprotic solvent such as dimethylformamide to give LI. The carbamate protecting group is removed under acidic conditions such as trifluoroacetic acid in methylene chloride, or hydrogen chloride gas in methanol or ethyl acetate, and the resulting piperazine can then be

5 carried on to final products as described in Schemes 3-9.

The isomeric piperazin-3-ones can be prepared as described in Scheme 13. The imine formed from arylcarboxamides LII and 2-aminoglycinal diethyl acetal (LIII) can be reduced under a variety of conditions, including sodium triacetoxyborohydride in dichloroethane,

10 to give the amine LIV. Amino acids I can be coupled to amines LIV under standard conditions, and the resulting amide LV when treated with aqueous acid in tetrahydrofuran can cyclize to the unsaturated LVI. Catalytic hydrogenation under standard conditions gives the requisite intermediate LVII, which is elaborated to final products as

15 described in Schemes 3-9.

Access to alternatively substituted piperazines is described in Scheme 14. Following deprotection with trifluoroacetic acid, the N-benzyl piperazine V can be acylated with an aryl carboxylic acid. The resulting N-benzyl aryl carboxamide LIX can be hydrogenated in the

20 presence of a catalyst to give the piperazine carboxamide LX which can then be carried on to final products as described in Schemes 3-9.

Reaction Scheme 15 provides an illustrative example the synthesis of compounds of the instant invention wherein the substituents R^2 and R^3 are combined to form $-(CH_2)_n-$. For example, 1-aminocyclohexane-1-carboxylic acid LXI can be converted to the spiropiperazine LXVI essentially according to the procedures outlined in Schemes 1 and 2. The piperazine intermediate LXIX can be deprotected as before, and carried on to final products as described in Schemes 3-9. It is understood that reagents utilized to provide the

30 substituent Y which is 2-(naphthyl) and the imidazolylalkyl substituent may be readily replaced by other reagents well known in the art and readily available to provide other N-substituents on the piperazine.

The aldehyde XLIX from Scheme 12 can also be reductively alkylated with an aniline as shown in Scheme 16. The

- 52 -


product LXXI can be converted to a piperazinone by acylation with chloroacetyl chloride to give LXXII, followed by base-induced cyclization to LXXIII. Deprotection, followed by reductive alkylation with a protected imidazole carboxaldehyde leads to LXXV, which can be alkylation with an arylmethylhalide to give the imidazolium salt LXXVI. Final removal of protecting groups by either solvolysis with a lower alkyl alcohol, such as methanol, or treatment with triethylsilane in methylene chloride in the presence of trifluoroacetic acid gives the final product LXXVII.

10 Scheme 17 illustrates the use of an optionally substituted homoserine lactone LXXIX to prepare a Boc-protected piperazinone LXXXII. Intermediate LXXXII may be deprotected and reductively alkylated or acylated as illustrated in the previous Schemes. Alternatively, the hydroxyl moiety of intermediate LXXXII may be mesylated and displaced by a suitable nucleophile, such as the sodium salt of ethane thiol, to provide an intermediate LXXXIII. Intermediate LXXXII may also be oxidized to provide the carboxylic acid on intermediate LXXXIV, which can be utilized form an ester or amide moiety.


20 Amino acids of the general formula LXXXVI which have a sidechain not found in natural amino acids may be prepared by the reactions illustrated in Scheme 18 starting with the readily prepared imine LXXXV.

25 Schemes 19-22 illustrate syntheses of suitably substituted aldehydes useful in the syntheses of the instant compounds wherein the variable W is present as a pyridyl moiety. Similar synthetic strategies for preparing alkanols that incorporate other heterocyclic moieties for variable W are also well known in the art.


- 53 -

SCHEME 1


- 54 -

SCHEME 2

- 55 -


SCHEME 3

- 56 -


SCHEME 3 (continued)

- 57 -


SCHEME 4




- 58 -

SCHEME 5


- 59 -

SCHEME 5 (continued)


- 60 -

SCHEME 6




- 61 -

SCHEME 6 (CONTINUED)




XXVIII




- 62 -

SCHEME 7


- 63 -

SCHEME 8


- 64 -

SCHEME 9


- 65 -

SCHEME 9 (continued)





- 66 -

SCHEME 10


- 67 -

SCHEME 11


- 68 -

SCHEME 12


- 69 -

SCHEME 12 (CONT'D)


- 70 -

SCHEME 13


- 71 -

SCHEME 14




- 72 -

SCHEME 15




- 73 -

SCHEME 15 (continued)


- 74 -

SCHEME 16


- 75 -

SCHEME 16 (continued)


- 76 -

SCHEME 17


- 77 -

SCHEME 17 (continued)


- 78 -

SCHEME 18


- 79 -

REACTION SCHEME 19


- 80 -

REACTION SCHEME 20

- 81 -

REACTION SCHEME 21

- 82 -

REACTION SCHEME 22

- 83 -

The instant compounds are useful as pharmaceutical agents for mammals, especially for humans. These compounds may be administered to patients for use in the treatment of cancer. Examples of the type of cancer which may be treated with the compounds of this invention include, but are not limited to, colorectal carcinoma, exocrine pancreatic carcinoma, myeloid leukemias and neurological tumors. Such tumors may arise by mutations in the *ras* genes themselves, mutations in the proteins that can regulate Ras activity (i.e., neurofibromin (NF-1), neu, scr, abl, lck, fyn) or by other mechanisms.

The compounds of the instant invention inhibit farnesyl-protein transferase and the farnesylation of the oncogene protein Ras. The instant compounds may also inhibit tumor angiogenesis, thereby affecting the growth of tumors (J. Rak et al. *Cancer Research*, 55:4575-4580 (1995)). Such anti-angiogenesis properties of the instant compounds may also be useful in the treatment of certain forms of blindness related to retinal vascularization.

The compounds of this invention are also useful for inhibiting other proliferative diseases, both benign and malignant, wherein Ras proteins are aberrantly activated as a result of oncogenic mutation in other genes (i.e., the Ras gene itself is not activated by mutation to an oncogenic form) with said inhibition being accomplished by the administration of an effective amount of the compounds of the invention to a mammal in need of such treatment. For example, a component of NF-1 is a benign proliferative disorder.

The instant compounds may also be useful in the treatment of certain viral infections, in particular in the treatment of hepatitis delta and related viruses (J.S. Glenn et al. *Science*, 256:1331-1333 (1992)).

The compounds of the instant invention are also useful in the prevention of restenosis after percutaneous transluminal coronary angioplasty by inhibiting neointimal formation (C. Indolfi et al. *Nature medicine*, 1:541-545(1995)).

The instant compounds may also be useful in the treatment and prevention of polycystic kidney disease (D.L. Schaffner et al.

- 84 -

American Journal of Pathology, 142:1051-1060 (1993) and B. Cowley, Jr. et al. *FASEB Journal*, 2:A3160 (1988)).

The instant compounds may also be useful for the treatment of fungal infections.

5 The compounds of this invention may be administered to mammals, preferably humans, either alone or, preferably, in combination with pharmaceutically acceptable carriers or diluents, optionally with known adjuvants, such as alum, in a pharmaceutical composition, according to standard pharmaceutical practice. The
10 compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

 For oral use of a chemotherapeutic compound according to this invention, the selected compound may be administered, for
15 example, in the form of tablets or capsules, or as an aqueous solution or suspension. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch, and lubricating agents, such as magnesium stearate, are commonly added. For oral
administration in capsule form, useful diluents include lactose and dried
20 corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavoring agents may be added. For intramuscular, intraperitoneal, subcutaneous and intravenous use,
sterile solutions of the active ingredient are usually prepared, and the
25 pH of the solutions should be suitably adjusted and buffered. For intravenous use, the total concentration of solutes should be controlled in order to render the preparation isotonic.

 The compounds of the instant invention may also be co-administered with other well known therapeutic agents that are selected
30 for their particular usefulness against the condition that is being treated. For example, the instant compounds may be useful in combination with known anti-cancer and cytotoxic agents. Similarly, the instant compounds may be useful in combination with agents that are effective in the treatment and prevention of NF-1, retinosis, polycystic kidney

- 85 -

disease, infections of hepatitis delta and related viruses and fungal infections.

If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range
5 described below and the other pharmaceutically active agent(s) within its approved dosage range. Compounds of the instant invention may alternatively be used sequentially with known pharmaceutically acceptable agent(s) when a combination formulation is inappropriate.

The present invention also encompasses a pharmaceutical
10 composition useful in the treatment of cancer, comprising the administration of a therapeutically effective amount of the compounds of this invention, with or without pharmaceutically acceptable carriers or diluents. Suitable compositions of this invention include aqueous solutions comprising compounds of this invention and pharmacolo-
15 gically acceptable carriers, e.g., saline, at a pH level, e.g., 7.4. The solutions may be introduced into a patient's blood-stream by local bolus injection.

When a compound according to this invention is administered into a human subject, the daily dosage will normally be
20 determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms.

In one exemplary application, a suitable amount of compound is administered to a mammal undergoing treatment for
25 cancer. Administration occurs in an amount between about 0.1 mg/kg of body weight to about 60 mg/kg of body weight per day, preferably of between 0.5 mg/kg of body weight to about 40 mg/kg of body weight per day.

The compounds of the instant invention are also useful
30 as a component in an assay to rapidly determine the presence and quantity of farnesyl-protein transferase (FPTase) in a composition. Thus the composition to be tested may be divided and the two portions contacted with mixtures which comprise a known substrate of FPTase (for example a tetrapeptide having a cysteine at the amine

- 86 -

terminus) and farnesyl pyrophosphate and, in one of the mixtures, a compound of the instant invention. After the assay mixtures are incubated for an sufficient period of time, well known in the art, to allow the FPTase to farnesylate the substrate, the chemical content of the assay mixtures may be determined by well known immunological, radiochemical or chromatographic techniques. Because the compounds of the instant invention are selective inhibitors of FPTase, absence or quantitative reduction of the amount of substrate in the assay mixture without the compound of the instant invention relative to the presence of the unchanged substrate in the assay containing the instant compound is indicative of the presence of FPTase in the composition to be tested.

It would be readily apparent to one of ordinary skill in the art that such an assay as described above would be useful in identifying tissue samples which contain farnesyl-protein transferase and quantitating the enzyme. Thus, potent inhibitor compounds of the instant invention may be used in an active site titration assay to determine the quantity of enzyme in the sample. A series of samples composed of aliquots of a tissue extract containing an unknown amount of farnesyl-protein transferase, an excess amount of a known substrate of FPTase (for example a tetrapeptide having a cysteine at the amine terminus) and farnesyl pyrophosphate are incubated for an appropriate period of time in the presence of varying concentrations of a compound of the instant invention. The concentration of a sufficiently potent inhibitor (i.e., one that has a K_i substantially smaller than the concentration of enzyme in the assay vessel) required to inhibit the enzymatic activity of the sample by 50% is approximately equal to half of the concentration of the enzyme in that particular sample.

30

EXAMPLES

Examples provided are intended to assist in a further understanding of the invention. Particular materials employed, species

- 87 -

and conditions are intended to be further illustrative of the invention and not limitative of the reasonable scope thereof. Purification by HPLC was accomplished with a 40 X 100 mm Waters PrepPak® reverse phase HPLC column (Delta-Pak™ C18 15 µm, 100 Å).

- 5 Gradient elution employed 0.1% trifluoroacetic acid in water (Solvent A) and 0.1% trifluoroacetic acid in acetonitrile (Solvent B). Chloride salts were obtained by passing an aqueous solution of the trifluoroacetic acid salt through a Biorad AG® 3X4 ion exchange resin column (100-200 mesh, C1-form). Purification by HPLC was utilized for each of the
- 10 Examples 1-23, 27, 48 and 49 as set forth below.

EXAMPLE 1

- 15 2(S)-Butyl-1-(2,3-diaminoprop-1-yl)-4-(1-naphthoyl)piperazine trihydrochloride

Step A: 1-Benzyl-3(S)-*n*-butylpiperazine-2,5-dione

- The title compound was prepared according to the procedure described by John S. Kiely and Stephen R. Priebe in Organic
- 20 Preparations and Procedures Int., 22 (6), 761-768 (1990). Thus dicyclohexylcarbodiimide (9.33 g, 45.2 mmol) in methylene chloride (0.5 M) was added to methylene chloride (250 mL). This solution was cooled to 0°C under nitrogen and Boc-L-norleucine (10.5 g, 45.2 mmol) was added. The resulting slurry was stirred for 5 min, and then
- 25 ethyl N-benzylglycinate (8.72 g, 45.2 mmol) was added. The reaction was stirred for 2 h at 0°C, then at 20°C overnight. The precipitate was removed by filtration, and hydrogen chloride gas bubbled through the methylene chloride solution for 2-4 h, until the reaction was shown to be complete by tlc. The solvent was removed in vacuo, and the residue
- 30 partitioned between ethyl acetate (150 mL) and saturated sodium bicarbonate solution (42 mL). The organic phase was washed with saturated sodium chloride, dried over magnesium sulfate, filtered and evaporated. The crude diketopiperazine was triturated with hexane to give the title compound as a white powder. ¹HNMR (300 MHz, CDCl₃)

- 88 -

δ 7.24-7.40 (5H, m), 6.22 (1H, br s), 4.07 (1H, dt, $J=3, 6$ Hz), 3.87 (1H, d, $J=17$ Hz), 3.80 (1H, d, $J=17$ Hz), 1.88 (2H, m), 1.35 (4H, m), 0.91 (3H, t, $J=7$ Hz).

5 **Step B:** 4-Benzyl-1-*tert*-butoxycarbonyl-2(S)-*n*-butylpiperazine

The product from Step A (4.95 g, 0.019 mol) was dissolved in THF (200 mL) and cooled under nitrogen to 0°C with mechanical stirring. Lithium aluminum hydride (2.60 g, 0.0685 mol) was added slowly. The reaction was refluxed for 18 h, cooled to 0°C, and
10 quenched by the sequential slow addition of 5 mL H₂O, 5 mL 10% sodium hydroxide solution and 5 mL H₂O. The reaction was stirred for 30 min and filtered. The solvent was removed in vacuo, the crude product taken up in methylene chloride and dried over magnesium sulfate. The drying agent was removed by filtration, and the filtrate
15 treated with di-*tert*-butyl dicarbonate ((4.35 g, 0.020 mol). After 2 h at 20°C, saturated sodium bicarbonate was added. The layers were separated, and the organic phase washed with saturated sodium chloride solution, then dried over magnesium sulfate. Filtration and evaporation
20 gave the crude product which was purified by column chromatography on silica gel, eluting with 5% ethyl acetate in hexane. The title compound was obtained as a foam. ¹HNMR (300 MHz, DMSO-d₆) δ 7.25 (5H, m), 3.90 (1H, br s.), 3.73 (1H, d, $J=13$ Hz), 3.51 (1H, d, $J=13$ Hz), 3.34 (1H, d, $J=13$ Hz), 2.93 (1H, m), 2.75 (1H, d, $J=11$ Hz), 2.62 (1H, d, $J=11$ Hz), 1.90 (2H, m), 1.60 (2H, m), 1.38 (9H, s), 1.26 (2H,
25 m), 1.04 (2H, m), 0.84 (3H, t, $J=7$ Hz).

Step C: 1-*tert*-Butoxycarbonyl-2(S)-*n*-butylpiperazine

The product from Step B (3.75 g, 11.3 mmol) was dissolved in methanol (75 mL) in a Parr bottle, and the vessel purged
30 with argon. To this was added 10 % palladium on carbon (0.80 g) and the reaction hydrogenated under 60 psi hydrogen for 24 h. The catalyst was removed by filtration through Celite, and the filtrate evaporated in vacuo to give the title compound as an oil. ¹HNMR (300 MHz, CDCl₃)

- 89 -

δ 4.08 (1H, br s), 3.90 (1H, d, $J=12$ Hz), 2.5-3.8 (6H, m), 1.80 (1H, m), 1.60 (1H, m), 1.46 (9H, s), 1.30 (4H, m), 0.90 (3H, t, $J=7$ Hz).

Step D: 1-tert -Butoxycarbonyl-2(S)-n -butyl-4-(1-naphthoyl)-piperazine

5
1-tert -butoxycarbonyl-2(S)-n -butylpiperazine (0.325 g, 1.34 mmol), 1-hydroxybenzotriazole (HOBT) (0.203 g, 1.34 mmol) and 1-ethyl-3-(3-dimethylamino-propyl)carbodiimide hydrochloride (EDC · HCl) (0.254 g, 1.34 mmol) were added to dry, degassed
10 dimethylformamide (7 mL). The pH of the reaction was adjusted to 7 with triethylamine, and the reaction stirred for 2 h. The dimethylformamide (DMF) was distilled in vacuo, and the residue partitioned between ethyl acetate and water. The organic phase was washed with 2% aqueous potassium hydrogen sulfate, saturated sodium
15 bicarbonate solution, saturated sodium chloride solution, and dried over magnesium sulfate. The title compound was obtained as a thick oil.

Step E: 2(S)-n -Butyl-4-(1-naphthoyl)piperazine hydrochloride

The product from Step D was dissolved in ethyl acetate,
20 cooled to -40°C under nitrogen, and the solution saturated with HCl(g). The solution was warmed to 0°C for 30 min, and then purged with nitrogen. The solvent was removed in vacuo. The product was evaporated from ethyl acetate three times. The title compound was obtained as a white solid.

25
Step F: 2,3-(bis-tert-Butoxycarbonylamino)propanoic acid

Diaminopropanoic acid monohydrochloride (2.86 g, 0.0204 mol) was suspended in 1:1 water-dioxane (100 mL) containing triethylamine (5.97 mL, 0.0204 mol). BOC-ON (11.0 g, 0.0448 mol)
30 was added along with additional triethylamine to adjust the pH to 9.5. The reaction was stirred under nitrogen overnight at 20°C . The clear solution was diluted with water and extracted with diethyl ether (5 x 100 mL). The aqueous solution was adjusted to pH 1 with cold 5% aqueous hydrochloric acid and extracted with ethyl acetate. The organic

- 90 -

layer was washed with water, saturated sodium chloride, then dried over magnesium sulfate. The resulting foam (5.46 g) was crystallized from ethyl acetate to give the title compound as a white solid.

5 Step G: N-Methoxy-N-methyl-2,3-(bis-tert-butoxycarbonyl-amino)propionamide

2,3-bis-(tert-Butoxycarbonylamino)propanoic acid (1.80 g, 5.92 mmol) in dimethylformamide (25 mL) was stirred overnight with N-methoxy-N-methylamine hydrochloride (0.635 g, 6.51 mmol), EDC ·
10 HCl (1.24 g, 6.51 mmol), N-hydroxybenzotriazole (0.80 g, 5.92 mmol) and triethylamine (0.825 mL, 5.92 mmol). The dimethylformamide was removed in vacuo and the residue partitioned between ethyl acetate and water. The organic phase was washed with 10% hydrochloric acid, saturated sodium bicarbonate solution, saturated brine, and dried over
15 magnesium sulfate. The crude product was chromatographed on silica gel with 30% ethyl acetate in hexane. The title compound was obtained as a foam. ¹HNMR (CDCl₃, 300 MHz) δ 5.51 (1H, br d), 4.87 (1H, br s), 4.72 (1H, br s), 3.77 (3H, s), 3.50 (1H, m), 3.40 (1H, dt, J=12, 6 Hz), 3.20 (3H, s), 1.44 (9H, s), 1.42 (9H, s).

20

Step H: 2,3-(bis-tert-Butoxycarbonylamino)propanal

Lithium aluminum hydride (0.384 g, 10.14 mmol) was suspended in diethyl ether (20 mL) and cooled to -45°C under nitrogen. N-Methoxy-N-methyl-2,3-(bis-tert-butoxycarbonylamino)-
25 propionamide (2.07 g, 5.96 mmol) in 1:1 ether-tetrahydrofuran (60 mL) was added at such a rate so as to keep the reaction temperature less than -35°C. The reaction was allowed to warm to 5°C, then cooled to -45°C and quenched with a solution of potassium hydrogen sulfate (3.08 g, 22.6 mmol) in water (20 mL). The reaction was stirred at 20°C 1h,
30 then filtered through celite. The organic phase was washed with 10% citric acid and saturated brine, then dried over magnesium sulfate. The title compound was obtained as a foam. NMR (CDCl₃, 300 MHz) δ 9.61 (1H, s), 5.60 (1H, br s), 4.88 (1H, br s), 4.24 (1H, br d, J=6 Hz), 3.68 (1H, m), 3.50 (1H, m), 1.40 (9H, s), 1.39 (9H, s).

- 91 -

Step I: 1-[(2,3-*bis-tert*-Butoxycarbonylamino)prop-1-yl]-2(S)-butyl-4-(1-naphthoyl)piperazine

A solution of 3(S)-butyl-1-(1-naphthoyl)piperazine (1.83 g, 6.20 mmol) (free base of the product from Step E), acetic acid (0.17 mL, 2.9 mmol) in dichloroethane (20 mL) was adjusted to pH 5.5 with triethylamine. Sodium triacetoxyborohydride (1.87 g, 8.79 mmol) and crushed molecular sieves (1 g) were added, and the reaction cooled to 0°C under nitrogen. A solution of 2,3-(*bis-tert*-butoxycarbonyl-amino)propanal (1.69 g, 5.86 mmol) in dichloroethane (10 mL) was added dropwise, and the reaction stirred at 20°C overnight. The reaction was cooled to 0°C, quenched with saturated sodium bicarbonate and stirred for 1h. The layers were separated and the organic layer washed with saturated brine, then dried over magnesium sulfate. The crude product was chromatographed on silica gel with 40-50% ethyl acetate in hexane, and the title compound was isolated as a foam (Rf 0.30, 50% ethyl acetate in hexane).

Step J: 1-(2,3-Diaminoprop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine trihydrochloride

Trifluoroacetic acid (25 mL) was added to a solution of 1-[(2,3-*bis-tert*-butoxycarbonylamino)prop-1-yl]-2(S)-butyl-4-(1-naphthoyl)piperazine (2.13 g, 3.75 mmol) in dichloromethane (75 mL). After 25 min at 20°C, the solvent was evaporated and the residue partitioned between chlorform and 20% aqueous sodium hydroxide. The organic layer was washed with saturated brine and dried over magnesium sulfate. The free base of the title compound was obtained as a yellow gum (1.72 g). A portion of this material (52 mg) was purified by preparative HPLC using a 100 mm Waters PrepPak[®] reverse phase column (DeltaPak[™] C18, 50 mM, 100 Å), and pure product isolated by gradient elution using 100% 0.1% trifluoroacetic acid in water (Solvent A) to 50% Solvent A and 50% 0.1% trifluoroacetic acid in acetonitrile (Solvent B). The pure fractions were combined and the solvent evaporated, and the crude product dissolved in water and passed

- 92 -

through an ion exchange column (Biorad AG[®] 3X4 ion exchange resin, chloride form). After lyophilization, the title compound was obtained as a white solid. Anal. Calc. for C₂₂H₃₂N₄O · 4.45 HCl C, 49.78; H, 6.92; N, 10.56. Found: C, 49.75; H, 6.72; N, 10.36.

5

EXAMPLE 2

1-(3-Amino-2-(2-naphthylmethylamino)prop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine trihydrochloride

10

Step A: 1-[(2-Amino-3-*tert*-butoxycarbonylaminoprop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine

Di-*tert*-butyl dicarbonate (0.282 g, 1.29 mmol) was added to a solution of 4-(2,3-diaminoprop-1-yl)-2(S)-butyl-1-(1-naphthoyl)piperazine (0.476 g, 1.29 mmol) in dichloromethane (10 mL). The reaction was stirred a total of 2h at 20°C, then quenched and extracted with water. The organic phase was washed with saturated brine and dried over magnesium sulfate. The crude product was chromatographed on silica gel with 5% methanol in chloroform followed by 5% (9:1 methanol-ammonium hydroxide) in chloroform. The title compound was isolated as the major product. FAB ms (m+1) 469.

15

20

Step B: 1-(3-*tert*-Butoxycarbonylamino-2-(2-naphthylmethylamino)prop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine

25

The title compound was prepared according to the procedure described in Example 1, Step D except using 1-[(2-amino-3-*tert*-butoxycarbonylaminoprop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine (0.287 g, 0.613 mmol), naphthalene-2-carboxaldehyde (0.95 g, 0.613 mmol), sodium triacetoxymethylborohydride (0.194 g, 0.919 mmol), in dichloroethane (15 mL) at pH 6. The crude product was chromatographed on silica gel with 5% methanol in

30

- 93 -

chloroform (R_f 0.30), and the title compound isolated as a foam. FAB ms ($m+1$) 609.

Step C: 1-(3-Amino-2-(2-naphthylmethylamino)prop-1-yl)-2(S)-
butyl-4-(1-naphthoyl)piperazine trihydrochloride

5 A solution of 1-(3-*tert*-butoxycarbonylamino-2-(2-naphthylmethylamino)prop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine (0.313 g, 0.514 mmol) in methylene chloride (10 mL) was deprotected with trifluoroacetic acid (5 mL) and converted to the free base (255
10 mg) according to the procedure described in Example 1, Step E. Purification of 40 mg by preparative HPLC used gradient elution with solvents A and B (from Example 1; 95% to 5% solvent A). Ion
exchange and lyophilization as described furnished the title compound as
a white solid. FAB ms ($m+1$) 509. Anal. Calc. for $C_{33}H_{40}N_4O \cdot 0.05$
15 H_2O 4.45 HCl: C, 59.00; H, 6.68; N, 8.34. Found: C, 59.00; H, 6.51; N, 8.44.

EXAMPLE 3

20 2(S)-Butyl-1-{5-[1-(2-naphthylmethyl)]-4,5-dihydroimidazol}methyl-4-(1-naphthoyl)piperazine ditrifluoroacetate

1-(3-Amino-2-(2-naphthylmethylamino)prop-1-yl)-2(S)-
butyl-4-(1-naphthoyl)piperazine (0.105 g, 0.207 mmol), *tert*-
butylisocyanate (1.5 mL) and silver cyanide (0.023 g, 0.207 mmol)
25 were heated in a sealed tube under nitrogen at 90°C overnight. The volatiles were removed in vacuo, and the residue chromatographed on silica gel with 5-10% methanol in chloroform to give the free base (73 mg). This was purified by preparative HPLC as described in Example 1
by gradient elution with solvents A and B (from Example 1; 95% to 5%
30 solvent A). Lyophilization furnished the title compound as a white solid. FAB ms ($m+1$) 519. Anal. Calc. for $C_{34}H_{38}N_4O \cdot 0.85 H_2O$ 3.75 TFA: C, 51.84; H, 4.55; N, 5.83. Found: C, 51.83; H, 4.56; N, 6.32.

- 94 -

EXAMPLE 41-[5-(1-Benzylimidazol)methyl]-2(S)-butyl-4-(1-naphthoyl)piperazine dihydrochloride

5 Benzyl bromide (0.012 mL, 0.103 mmol) was added to a solution of 2(S)-butyl-1-[5-(3-triphenylmethylimidazol)]methyl-4-(1-naphthoyl)piperazine (63 mg, 0.103 mmol) in acetonitrile (0.5 mL) at 20°C under nitrogen. The reaction was stirred overnight, concentrated in vacuo, and taken up in dichlormethane (2 mL) containing
10 triethylsilane (0.100 mL). Trifluoroacetic acid was added and the reaction stirred for 1h at 20°C. The solvent was evaporated and the residue purified by HPLC (95% to 5% solvent A). Pure fractions were combined and converted to the HCl salt as described in Example 1, Step E. After lyophilization, the title compound was isolated as a white
15 solid. FAB ms (m+1) 467. Anal. Calc. for C₃₀H₃₄N₄O · 0.05 H₂O · 3.70 HCl: C, 59.81; H, 6.32; N, 9.30. Found: C, 59.78; H, 6.33; N, 9.30.

EXAMPLE 5

20 1-{5-[1-(4-nitrobenzyl)]imidazolylmethyl}-2(S)-butyl-4-(1-naphthoyl)piperazine ditrifluoroacetate

 The title compound was prepared according to the procedure described in Example 4, except using p-nitrobenzylbromide
25 (0.043 g, 0.199 mmol) and 2(S)-butyl-1-[5-(3-triphenylmethylimidazol)]methyl-4-(1-naphthoyl)piperazine (123 mg, 0.199 mmol) in acetonitrile (2 mL). The crude product was treated with triethylsilane (0.127 mL, 0.80 mmol) and trifluoroacetic acid (2 mL) in dichloroethane (4 mL). Preparative HPLC (95-5% solvent A)
30 provided the title compound as a white solid. FAB ms (m+1) 512. Anal. Calc. for C₃₀H₃₃N₅O₃ · 2 CF₃CO₂H · 0.03 H₂O: C, 41.42; H, 3.98; N, 10.18. Found: C, 41.43; H, 3.96; N, 10.51.

- 95 -

EXAMPLE 6

1-(3-Acetamidomethylthio-2(R)-aminoprop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine ditrifluoroacetate

- 5 1-(3-Acetamidomethylthio-2(R)-aminoprop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine dihydrochloride and N-hydroxymethylacetamide (0.010 g, 0.105 mmol) (prepared as described in M. Bodansky, A. Bodansky, "The Practice of Peptide Synthesis", Springer-Verlag, 1984, p.82) were dissolved in trifluoroacetic acid for
10 0.5h. The solvent was removed in vacuo and the residue purified by preparative HPLC (85-60% solvent A). After lyophilization, the title compound was isolated as a solid. FAB ms (m+1) 457. Anal. Calc. for $C_{25}H_{36}N_4O_2 \cdot 2 CF_3CO_2H \cdot 1.9 H_2O$. C, 44.38 H, 5.12; N, 6.64. Found: C, 44.35; H, 5.11; N, 6.97.

15

EXAMPLE 7

2(S)-Butyl-1-[2-(1-imidazolyl)ethyl]sulfonyl-4-(1-naphthoyl)piperazine ditrifluoroacetate

20

Step A: 2(S)-Butyl-4-(1-naphthoyl)-1-vinylsulfonylpiperazine

- Chloroethylsulfonylchloride (0.038 mL, 0.314 mmol) was added to a solution of 3(S)-butyl-1-(1-naphthoyl)piperazine (0.095 g, 0.285 mmol) and diisopropylethylamine (0.119 mL, 0.685 mmol) in
25 dichloromethane (3 mL). The reaction was stirred overnight under nitrogen, quenched with saturated sodium bicarbonate and extracted into ethyl acetate. After drying with magnesium sulfate, the title compound was isolated.

- 30 Step B: 2(S)-Butyl-1-[2-(1-imidazolyl)ethyl]sulfonyl-4-(1-naphthoyl)piperazine ditrifluoroacetate
-

Imidazole (0.043 g, 0.627 mmol) was added to sodium hydride (60% dispersion in oil, 0.024 g, 0.598 mmol) suspended in dimethylformamide (2 mL). The reaction was cooled to 0°C under

- 96 -

nitrogen, and 2(S)-butyl-4-(1-naphthoyl)-1-vinylsulfonylpiperazine (0.011 g, 0.29 mmol) in dimethylformamide (5 mL) was added. The reaction was stirred at 20°C overnight. The dimethylformamide was removed in vacuo and the residue dissolved in ethyl acetate. This was
5 extracted with saturated sodium bicarbonate solution, saturated brine and dried over magnesium sulfate. The crude product was first purified by silica gel chromatography using 8% methanol in chloroform, then by preparative HPLC (80 to 40% solvent A). The trifluoroacetate salt was dissolved in water and partitioned between saturated sodium bicarbonate
10 and ethyl acetate. The organic phase was washed with saturated brine and dried over magnesium sulfate. The title compound was obtained as a foam. FAB ms (m+1) 455. Anal. Calc. for C₂₅H₃₆N₄O₂ S · 0.8 H₂O. C, 61.46; H, 6.79; N, 11.95. Found: C, 61.44; H, 6.97; N, 10.72.

15

EXAMPLE 8

2(R)-Butyl-1-imidazolyl-4-methyl-4-(1-naphthoyl)piperazine

Step A: 2(R)-Butyl-4-(1-naphthoyl)-1-[4-(1-
20 triphenylmethylimidazolyl)methyl-piperazine
3(R)-Butyl-1-(1-naphthoyl)piperazine (0.202 g, 0.607 mmol) (prepared as described for the (S) enantiomer in Example 1) was reacted with 1-triphenylmethylimidazole-4-carboxaldehyde (0.226 g, 0.667 mmol), sodium triacetoxyborohydride (0.321 g, 1.52 mmol), in
25 dichloroethane (7 mL) in the presence of crushed molecular sieves. The pH was adjusted to 5-6 with triethylamine/acetic acid. The reaction was stirred overnight, then quenched with saturated sodium bicarbonate. The organic layer was washed with saturated brine, and dried over magnesium sulfate. The crude product was purified by chromatography
30 on silica gel with 30% ethyl acetate in hexane followed by 5% methanol in chloroform, to obtain the title compound.

Step B: 2(R)-Butyl-1-imidazolyl-4-methyl-4-(1-
naphthoyl)piperazine ditrifluoroacetate

- 97 -

Triethylsilane (1.0 mL, 11.80 mmol) was added to a solution of 2(R)-butyl-4-(1-naphthoyl)-1-[4-(1-triphenylmethyl-imidazolyl)methyl-piperazine (0.381 g, 0.616 mmol) in dichloromethane, followed by trifluoroacetic acid (8 mL). After 1h, the solvents were evaporated and the residue partitioned between water and hexane. The aqueous phase was injected directly onto a preparative HPLC column (100-40% solvent A) and the title compound isolated after lyophilization. FAB ms (m+1) 377. Anal. Calc. for $C_{25}H_{36}N_4O_2 \cdot 2.35 CF_3CO_2H \cdot 0.32 H_2O$. C, 48.93; H, 4.52; N, 7.98. Found: C, 48.93; H, 4.55; N, 8.26.

EXAMPLE 9

2(S)-Butyl-4-(1-naphthoyl)-1-(3-pyridylmethyl)piperazine dihydrochloride

3(S)-Butyl-1-(1-naphthoyl)piperazine hydrochloride (0.200 g, 0.601 mmol) was reacted with pyridine-3-carboxaldehyde (0.062 mL, 0.661 mmol), sodium triacetoxyborohydride (0.321 g, 1.52 mmol), in dichloroethane (7 mL) at pH 5-6 in the presence of crushed molecular sieves as described in Example 8, Step A. The crude product was purified by silica gel chromatography with 30% acetone in hexane, followed by preparative HPLC (80-75% solvent A). After ion exchange, the title compound was isolated. FAB ms (m+1) 388. Anal. Calc. for $C_{25}H_{29}N_3O \cdot 2.3 HCl \cdot 0.95 H_2O$. C, 61.49; H, 6.85; N, 8.60. Found: C, 61.49; H, 7.01; N, 8.76.

EXAMPLE 10

1-2(S)-Butyl-(2(R)-(4-nitrobenzyl)amino-3-hydroxypropyl)-4-(1-naphthoyl)piperazine dihydrochloride

1-(2(R)-amino-3-hydroxypropyl)-2(S)-butyl-4-(1-naphthoyl)piperazine (120 mg, 0.326 mmol) was converted to the title compound according to the procedure described in Example 8, Step A using 4-nitrobenzaldehyde (0.493 g, 0.327 mmol), sodium

- 98 -

triacetoxyborohydride (0.173 g, 0.817 mmol), in dichloroethane. The crude product was purified by preparative HPLC (100-75% solvent A), and after ion exchange to the HCl salt and lyophilization, the title compound was obtained. FAB ms (m+1) 505. Anal. Calc. for C₂₅H₂₉N₃O · 3.6 HCl · 0.10 H₂O. C, 54.69; H, 6.30; N, 8.80. Found: C, 54.66; H, 5.85; N, 8.31.

EXAMPLE 11

10 1-(2(R)-Amino-3-hydroxyheptadecyl)-2(S)-butyl-4-(1-naphthoyl)piperazine ditrifluoroacetate

Step A: 1-(2(R)-*t*-Butyoxycarbonylamino-2-formylethyl)-2(S)-butyl-4-(1-naphthoyl)piperazine

15 A solution of oxalyl chloride (1.36 mL, 14.9 mmol) in dichloromethane (35 mL) was cooled to -65°C under nitrogen, and dimethylsulfoxide (2.30 mL, 32.4 mmol) in methylene chloride (7 mL) added, and the reaction stirred 2 min. A solution of 1-(2(R)-amino-3-hydroxypropyl)-2(S)-butyl-4-(1-naphthoyl)piperazine (3.19 g, 6.79 mmol) was added to this solution at -10°C and the reaction stirred at this temperature for 15 min. The reaction was cooled to -55°C and triethylamine (4.76 mL, 34 mmol) added. The reaction was stirred for 5 min then warmed to room temperature. Additional methylene chloride is added and the reaction is extracted with water. The organic phase was washed with 2% potassium hydrogen sulfate, water, dilute sodium bicarbonate solution, and saturated brine. After drying with magnesium sulfate, the title compound was obtained.

25 Step B: 1-(2(R)-Amino-3-hydroxyheptadecyl)-2(S)-butyl-4-(1-naphthoyl)piperazine ditrifluoroacetate

30 A solution of 0.25 g, 0.53 mmol) 1-(2(R)-*t*-butoxycarbonylamino-2-formylethyl)-2(S)-butyl-4-(1-naphthoyl)piperazine in dry tetrahydrofuran (5 mL) was cooled to 0°C under nitrogen in a flame-dried three necked flask. A solution of

- 99 -

heptadecylmagnesium chloride (1.01 mL of a 1M solution in ether, 1.01 mmol) was added via syringe, and the reaction allowed to warm to room temperature. The reaction was quenched with saturated sodium bicarbonate solution, and then extracted with ethyl acetate. After
5 drying over magnesium sulfate, the crude product was chromatographed on silica gel with 25% ethyl acetate in hexane followed by 5% methanol in chloroform. The purified product was dissolved in methylene chloride (7 mL) and treated with trifluoroacetic acid (3.5 mL). After 45 min, the solvents were removed in vacuo and
10 the residue purified by preparative HPLC (95-40% solvent A). Two isomers were separated. After lyophilization, the title compound was isolated as diastereomer A (retention time 8.405 min, gradient 100-50% solvent A over 15 min), FAB ms (m+1) 566, Anal. Calc. for $C_{36}H_{59}N_3O_2 \cdot 2.35 CF_3CO_2H \cdot 0.35 H_2O$. C, 58.19; H, 7.44; N,
15 5.00. Found: C, 58.21; H, 7.46; N, 5.36. After lyophilization, the title compound was also isolated as diastereomer B (retention time 9.269 min, gradient 100-50% solvent A over 15 min), FAB ms (m+1) 566, Anal. Calc. for $C_{36}H_{59}N_3O_2 \cdot 2.35 CF_3CO_2H \cdot 0.05 H_2O$. C, 58.56; H, 7.42; N, 5.03. Found: C, 58.53; H, 7.41; N, 5.17.

20

EXAMPLE 12

2(S)-Benzyl-1-imidazolyl-4-methyl-4-(1-naphthoyl)piperazine

25 Step A: 1.(3S)-Dibenzylpiperazine-2,5-dione

The title compound was prepared according to the procedure described in Example 1, Step A, except using Boc-L-phenylalanine (12.8 g, 48.2 mmol), ethyl N-benzylglycinate (9.32 g, 48.2 mmol) and dicyclohexylcarbodiimide (96.5 mL 0.5 M in
30 dichloromethane, 48.2 mmol). The crude diketopiperazine was triturated with hexane to give the title compound as a white powder. 1H NMR (300 MHz, CD_3OD) δ 7.0-7.4 (10H, m), 4.61 (1H, d, J=16 Hz), 4.37 (1H, t, J=5 Hz), 4.24 (1H, d, J=16 Hz), 3.42 (1H, d, J=18 Hz),

- 100 -

3.28 (1H, dd, J=4, 16 Hz), 2.96 (1H, dd, J=6, 16 Hz), 2.55 (1H, d, J=18 Hz).

Step B: 1-tert -Butoxycarbonyl-2(S)-(S)-4-dibenzylpiperazine

5 The title compound was prepared according to the procedure described in Example 1, Step B, except using 1,3(S)-dibenzylpiperazine-2,5-dione (5.01 g, 17.1 mmol) and lithium aluminum hydride (2.33 g, 61.4 mmol), followed by di-tert -butyl dicarbonate (4.02 g, 18.4 mmol). The crude product was purified by
10 column chromatography on silica gel, eluting with 7.5% ethyl acetate in hexane. The title compound was obtained as a white solid. ¹HNMR (300 MHz, CD₃OD) δ 7.2-7.4 (5H, m), 7.0-7.2 (5H, m), 4.15 (1H, m), 3.90 (1H, d, J=15 Hz), 3.60 (1H, d, J=15 Hz), 3.15 (1H, m), 2.95 (3H, m), 2.7 (1H, d, J=13 Hz), 2.02 (1H, dt, J=6, 13 Hz), 1.95 (1H, br d),
15 1.35 (9H, s).

Step C: 2(S)-Benzyl-1-tert -butoxycarbonylpiperazine

The title compound was prepared according to the procedure described in Example 1, Step C, except using 1-tert -
20 butoxycarbonyl-2(S)-4-dibenzylpiperazine (4.78 g, 11.3 mmol) and 10% palladium on carbon (1.04 g). The title compound was obtained as an oil. ¹HNMR (300 MHz, CD₃OD) δ 7.25 (5H, m), 4.35 (1H, m), 4.00 (1H, d, J=12 Hz), 2.7-3.3 (7H, m), 1.25 (9H, s).

25 **Step D:** 2(S)-Benzyl-1-tert -butoxycarbonyl-4-(1-naphthoyl)piperazine

The title compound was prepared according to the procedure described for Example 1, Step A except using 2(S)-benzyl-1-
tert -butoxycarbonylpiperazine (0.292 g, 1.06 mmol), 2,3-dimethyl-
30 benzoic acid (0.159 g, 1.06 mmol), HOBT (0.157 g, 1.02 mmol), EDC·HCl (0.213 g, 1.11 mmol) and triethylamine to adjust the pH to 7. The title compound was obtained as a thick oil. ¹HNMR (DMSO-d₆, 300 MHz) δ 7.15 (2H, m), 6.06 (1H, m), 4.42 (1H, m), 3.6-4.2 (2H, m),

- 101 -

2.7-3.24 (4H, m), 2.24 (3H, s), 2.03-2.20 (3H, 4s), 1.10-1.6 (15H, m), 0.72-1.00 (3H, m).

Step E: 2(S)-Benzyl-4-(1-naphthoyl)-1-[4-(1-triphenylmethyylimidazolyl)methyl-piperazine

5
3(S)-Benzyl-1-(1-naphthoyl)piperazine (0.173 g, 0.472 mmol) was reacted with 1-triphenylmethyylimidazole-4-carboxaldehyde (0.160 g, 0.472 mmol), sodium triacetoxymethylborohydride (0.300 g, 1.42 mmol), in dichloroethane (7 mL) in the presence of crushed molecular
10 sieves as described in Example 8, Step A. The title compound was obtained as an oil.

Step F: 2(S)-Benzyl-1-imidazolyl-4-methyl-4-(1-naphthoyl)piperazine ditrifluoroacetate

15 Triethylsilane (0.300 mL, 1.89 mmol) was added to a solution of 2(S)-benzyl-4-(1-naphthoyl)-1-[4-(1-triphenylmethyylimidazolyl)methyl-piperazine (0.310 g, 0.472 mmol) in dichloromethane (5 mL), followed by trifluoroacetic acid (5 mL). After 1h, the solvents were evaporated and the residue partitioned
20 between water and hexane. The aqueous phase was injected directly onto a preparative HPLC column (85-45% solvent A) and the title compound isolated after lyophilization. FAB ms (m+1) 411. Anal. Calc. for C₂₆H₂₆N₄O₂ · 2.75 CF₃CO₂H · 0.05 H₂O. C, 52.11; H, 4.14; N, 7.72. Found: C, 52.10; H, 4.03; N, 8.16.

25

EXAMPLE 13

1-(2(R)-Amino-3-(3-benzylthio)propyl)-2(S)-butyl-4-(1-naphthoyl)-piperazine ditrifluoroacetate

30

Step A: 1-[(1-Aziridinyl)methyl]-2(S)-butyl-4-(1-naphthoyl)-piperazine

A solution of 1-(2(R)-butoxycarbonylamino-3-hydroxypropyl)-2(S)-butyl-4-(1-naphthoyl)piperazine (1.67 g, 3.56

- 102 -

mmol) in dimethylformamide (10 mL) was cooled to 0°C under nitrogen. Sodium hydride (0.427 g, 10.6 mmol, 60% dispersion in oil) was added, followed by 1,1'-sulfonyldiimidazole (0.704 g, 3.56 mol). The reaction was warmed to 20°C for 1h, cooled to 0°C and quenched with water. Dimethylformamide was distilled in vacuo, and the residue partitioned between ethyl acetate and water. The organic phase was washed with saturated brine and dried over magnesium sulfate. The crude product was chromatographed on silica gel with 70% ethyl acetate in hexane, followed by 5% methanol in chloroform. The title compound was obtained as the major product, FAB ms (m+1) 352. A lesser amount of 1-[(1-butoxycarbonylaziridinyl)methyl]-2(S)-butyl-4-(1-naphthoyl)piperazine was also isolated.

Step B: 1-(2(R)-Amino-3-(3-benzylthio)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine ditrifluoroacetate
1-[(1-Aziridinyl)methyl]-2(S)-butyl-4-(1-naphthoyl)piperazine (0.050 g, 0.142 mmol) was refluxed for 18h with benzyl mercaptan (0.100 mL, 0.852 mmol) and triethylamine (0.200 mL) in methanol (4 mL). The crude product was first chromatographed on silica gel with 3% methanol in chloroform, and then purified by preparative HPLC (85% to 10% solvent A). The title compound was obtained after lyophilization. FAB ms (m+1) 476. Anal. Calc. for C₂₉H₃₇N₃OS · 2.6 CF₃CO₂H · 0.3 H₂O. C, 52.83; H, 5.21; N, 5.40. Found: C, 52.78; H, 5.17; N, 5.66.

EXAMPLE 14

1-(2(R)-Amino-3-[3-(4-nitrobenzylthio)propyl]))-2(S)-butyl-4-(1-naphthoyl)piperazine ditrifluoroacetate
A solution of 1-[(1-butoxycarbonylaziridinyl)methyl]-2(S)-butyl-4-(1-naphthoyl)piperazine (0.050 g, 0.111 mmol) in methanol (4 mL) was relaxed with p-nitrobenzylthioacetate (0.070 g, 0.333 mmol) and triethylamine (0.200 mL) for 2h. The crude product was chromatographed on silica gel with 3% methanol in chloroform. The

- 103 -

purified product was treated with 33% trifluoroacetic acid in methylene chloride for 20 min. The solvents were evaporated and the product purified by preparative HPLC (85%-10% solvent A). The title compound was obtained after lyophilization. FAB ms (m+1) 529.

- 5 Anal. Calc. for $C_{29}H_{36}N_4O_3 \cdot 2 CF_3CO_2H \cdot 0.08 H_2O$. C, 51.94; H, 5.23; N, 7.34. Found. C, 51.87; H, 5.06; N, 7.47.

EXAMPLE 15

- 10 2(S)-Butyl-1-[(4-imidazolyl)ethyl]-4-(1-naphthoyl)piperazine dihydrochloride

Step A: N-Methyl-N-methoxy-2-(1-triphenylmethyl-1H-imidazol-4-yl)acetamide

- 15 To a solution of 4-imidazoleacetic acid (1.04g, 6.40 mmol) and triphenylmethyl bromide (2.48 g, 7.68 mmol) in dimethylformamide (40 ml) was added triethylamine (4.46 ml, 32 mmol) and the suspension allowed to stir for 18 hours at room temperature. After this time, the mixture was treated with 3-hydroxy-20 1,2,3-benzotriazin-4(3)-one (HOBT) (1.31g, 8 mmol), N, O-dimethylhydroxyamine hydrochloride (1.56g, 16 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (1.53 g, 8 mmol) and stirred for 24 hours at room temperature. After this time, sat. aq. sodium bicarbonate (50 ml) and water (50 ml) were 25 added and the mixture was extracted with ethyl acetate (2 x 100 ml). The combined extracts were washed with sat. aq. $NaHCO_3$ (100 ml) and then brine (50 ml) and the solvent evaporated in vacuo. The residue was suspended in ether (20 ml) and the white solid filtered to give the title compound as a white solid.
- 30 $^1NMR(CD_3OD, 300 MHz)$ δ 7.37(10H, m), 7.16(6H, m), 6.84(1H, s), 3.73(2H, s), 3.68(3H, s) and 3.18(3H, s)ppm.

- 104 -

Step B: 2-(1-triphenylmethyl-1H-imidazol-4-yl)acetaldehyde

To a solution of the product from Step A (300 mg, 0.73 mmol) in freshly distilled THF (15 ml) cooled to -40°C over dry ice/acetone was added lithium aluminum hydride (33.2 mg, 0.874 mmol). The resulting suspension was allowed to warm to +5°C and then maintained at 0°C for 30 min. After this time, the reaction mixture was recooled to -40°C and quenched sequentially with water (33 µl), 1.0N NaOH (33 µl) and water (100 µl). The resulting suspension was stirred for 30 min, filtered and the solvent removed in vacuo. The residue was dissolved in methylene chloride (5 ml) and washed with 10% aq. citric acid (5 ml) and then water (5ml). The organic layer was dried(MgSO₄) and the solvent evaporated in vacuo to give the title compound.

¹HNMR (CDCl₃, 300 MHz) δ 9.79(1H, t, J=3Hz), 7.44(1H, s), 7.4-7.1(15H, m), 6.76(1H, s) and 3.63(2H, d, J=3Hz)ppm.

Step C: 2(S)-Butyl-1-[(4-imidazolyl)ethyl]-4-(1-naphthoyl)piperazine ditrifluoroacetate

To a solution of the product from Step B (89 mg, 0.263 mmol) and 3(S)-butyl-1-naphthoylpiperazine hydrochloride (62.6 mg, 0.188 mmol) in 1,2-dichloroethane (4 ml) was added 3A molecular sieves (400 mg) and sodium triacetoxyborohydride (200 mg, 0.94 mmol). This mixture was stirred at room temperature for 4 days. After this time, filtered the mixture through sintered glass. The filtrate was diluted with methylene chloride (50 ml) and washed with water (25 ml). The solvent was evaporated in vacuo and the residue was purified by flash chromatography eluting with 2-5% methanol/methylene chloride to provide the trityl protected title compound as an oil. This oil was dissolved in methylene chloride (2 ml) and trifluoroacetic acid (1 ml) and treated with triethylsilane (2 drops) to give a colorless solution. This solution was stirred at room temperature for 3 hours and then the solvent was evaporated in vacuo. The resulting residue was dissolved in water (20 ml) and washed with hexanes (20 ml). The aqueous layer was lyophilized to give the title compound.

- 105 -

¹HNMR (CD₃OD, 300 MHz) δ 8.83(1H, s), 8.08-7.75(3H, m), 7.67-7.49(4H, m), 7.45(1H, s), 4.5-3.8(2H, m), 3.65-2.95(9H, m), 2.0-1.3(4H, m) and 2.2-0.2(5H, m)ppm.

Anal. calc'd for C₂₄H₃₀N₄O 3.45 TFA 0.75H₂O: C, 46.54; H, 4.42; N, 7.03. Found: C, 46.54; H, 4.41; N, 7.35.

FAB HRMS exact mass calc'd for C₂₃H₃₁N₄O 391.249787 (MH⁺), found 391.249028

EXAMPLE 16

2(S)-Butyl-1-[(4-imidazolyl)methyl]-4-(1-naphthoyl)piperazine ditrifluoroacetate

To a solution of the product from Step C (102 mg, 0.30 mmol) and 3(S)-butyl-1-naphthoylpiperazine hydrochloride (62.6 mg, 0.188 mmol) in 1,2-dichloroethane (5 ml) was added sodium triacetoxyborohydride (200 mg, 0.94 mmol) and triethylamine to pH = 5.5 and stirred at room temp. for 18 hours. After this time, the mixture was filtered. The filtrate was concentrated in vacuo and dissolved in ethyl acetate (25 ml) and washed with sat. aq. NaHCO₃ (10 ml) and then brine (10 ml). The solvent was evaporated in vacuo and the residue was purified by flash chromatography eluting with 3-8% methanol/methylene chloride to provide the trityl protected title compound. This oil was dissolved in methylene chloride (4 ml) and trifluoroacetic acid (2 ml) and treated with triethylsilane until the yellow color disappeared to give a colorless solution. Stirring continued at room temp. for 10 min. and the solvent evaporated in vacuo. The resulting white solid was partially dissolved in water (7 ml) and filtered. The filtrate was purified by Prep HPLC using a Nova Prep 5000 Semi Preparative HPLC system and a Waters PrepPak cartridge (47 X 300mm, C18, 15 μm, 100Å) eluting with 5 - 95% acetonitrile/water (0.1% TFA) at 100 ml/min (Chromatography A) to give the title compound after lyophilization

¹HNMR (DMSO-d₆, 400 MHz, 150°C) δ 8.53(1H, s), 7.94(2H, m), 7.79(1H, m), 7.53(3H, m), 7.41(1H, m), 7.40(1H, s), 4.12(1H, d,

- 106 -

J=4.8Hz), 3.95(1H, d, J=4.8Hz), 3.70(1H, s), 3.63(1H, s), 3.48(1H, s), 3.40(1H, s), 3.01(1H, s), 2.82(1H, s), 2.74(1H, s), 1.70(1H, m), 1.49(1H, m), 1.18(2H, s), 1.08(2H, s) and 0.77(3H, t, J=5.5Hz)ppm.
Anal. calc'd for C₂₃H₂₉N₄O 2 TFA 0.70H₂O: C, 52.55; H, 5.13; N, 9.08. Found: C, 52.54; H, 5.11; N, 9.35.

5 FAB MS mass calc'd for C₂₃H₃₀N₄O 377 (MH⁺), found 377.

EXAMPLE 17

10 2(S)-Butyl-1-[(1-naphth-2-ylmethyl)-1H-imidazol-5-yl]acetyl]-4-(1-naphthoyl)piperazine dihydrochloride

Step A: Preparation of 1H-Imidazole-4- acetic acid methyl ester hydrochloride

15 To a solution of 1H-imidazole-4-acetic acid hydrochloride (4.00g, 24.6 mmol) in methanol (100 ml) was bubbled hydrogen chloride gas until saturated. This solution was allowed to stand for 18h at room temperature and the solvent evaporated in vacuo to give the title compound as a white solid.

20 ¹H NMR(CDCl₃, 400 MHz) δ 8.85(1H, s), 7.45(1H, s), 3.89(2H, s) and 3.75(3H, s) ppm.

Step B: Preparation of 1-(Triphenylmethyl)-1H-imidazol-4-ylacetic acid methyl ester

25 To a suspension of the product from Step A (7.48g, 42.4 mmol) in methylene chloride (200ml) was added triethylamine (17.7 ml, 127 mmol) and triphenylmethyl bromide(16.4g, 50.8 mmol) and stirred for 72h. After this time, the reaction mixture was washed with sat. aq. NaHCO₃ (100 ml) and water (100 ml). The organic layer was
30 evaporated in vacuo and the residue was purified by flash chromatography(30-100% ethyl acetate in hexanes gradient elution) to provide the title compound as a white solid. ¹H NMR(CDCl₃, 400 MHz) δ 7.35(1H, s), 7.31(9H, m), 7.22(6H, m), 6.76(1H, s), 3.68(3H, s) and 3.60(2H, s) ppm.

- 107 -

Step C: Preparation of 2-[1-(Naphth-2-ylmethyl)-1H-imidazol-5-yl]acetic acid methyl ester

To a solution of the product of Step B (4.36g, 11.4 mmol) in acetonitrile (70 ml) was added 2-(bromomethyl)naphthalene and heated to 55°C for 4h. After this time, the reaction was cooled to room temperature and the resulting white precipitate was collected by filtration. The filtrate was concentrated to 30 ml and heated to 55°C for 18h. After this time, the reaction was cooled to room temperature and filtered the resulting white precipitate collected by filtration. The filtrate was concentrated to 10 ml volume and heated to 55°C for 1h. After this time, the reaction was again cooled to room temperature and diluted with ethyl acetate (25 ml). The resulting precipitate was collected by filtration and combined with the previous 2 precipitates in methanol (100 ml) and heated to reflux for 30m. After this time, the solvent was removed in vacuo and the resulting residue was partitioned between methylene chloride(200 ml) and sodium bicarbonate (100 ml). The organic layer was evaporated in vacuo to dryness and the residue was purified by flash chromatography (0-6% methanol/methylene chloride gradient) to provide the title compound as an off white solid : ¹HNMR (CDCl₃, 400 MHz) δ 7.82(2H, m), 7.75(1H, m), 7.70(1H, s), 7.49(3H, m), 7.20(1H, d, J=8.4Hz), 7.06(1H, s), 5.32(2H, s), 3.57(3H, s) and 3.49(2H, s) ppm.

Step D: Preparation of 2-[1-(Naphth-2-ylmethyl)-1H-imidazol-5-yl]acetic acid hydrochloride

2-[1-(Naphth-2-ylmethyl)-1H-imidazol-5-yl]acetic acid methyl ester (0.92g, 3.28mmol) was dissolved in 2.5N hydrochloric acid (50ml) and heated to 55°C for 3h. After this time, the solution was concentrated to dryness in vacuo to give the title compound as a white solid. ¹HNMR (CD₃OD, 400 MHz) δ 8.92(1H, s), 7.94(1H, d, J=8.6Hz), 7.88(2H, m), 7.83(1H, s), 7.54(3H, m), 7.43(1H, d, J=14Hz), 5.60(2H, s) and 3.82(2H, s) ppm.

- 108 -

Step E: Preparation of 2(S)-Butyl-1-[(1-(naphth-2-ylmethyl)-1H-imidazol-5-yl)acetyl]-4-(1-naphthoyl)piperazine dihydrochloride

5 To a solution of the product from Step D (100 mg, 0.330mmol), 3(S)-butyl-1-naphthoylpiperazine hydrochloride (100 mg, 0.300 mmol) and 3-hydroxy-1,2,3-benzotriazin-4(3H)-one (HOBT) (54 mg, 0.33mmol) in dimethylformamide (2 ml) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) 10 (63mg, 0.33 mmol) and triethylamine (161ul, 1.16mmol) and the resulting suspension stirred for 18 hours. After this time, sat. aq. NaHCO₃ (7 ml) was added and the resulting precipitate filtered. The solid was redissolved in methylene chloride and washed with water (25 ml) and brine (20 ml). The solvent was evaporated in vacuo and the 15 residue purified by Preparative HPLC (Chromatography A) to give after lyophilization the title compound.

¹HNMR (CDCl₃, 400 MHz) δ 8.98(1H, m), 8.03-7.27(15H, m), 5.69-5.50(2H, m), 4.79-4.34(2H, m), 4.15-2.50(7H, m), 1.8-0.2(9H, m)ppm. Anal. calc'd for C₃₅H₃₅N₄O₂ HCl 0.65H₂O: C, 67.30; H, 6.16; N, 20 8.97. Found: C, 67.25; H, 6.16; N, 9.15. FAB HRMS exact mass calc'd for C₃₅H₃₆N₄O₂ 545.291652 (MH⁺); found 545.292050.

EXAMPLE 18

25

2(S)-Butyl-1-[(1-naphth-2-ylmethyl)-1H-imidazol-5-yl)ethyl]-4-(1-naphthoyl)piperazine ditrifluoroacetate

Step A: Preparation of N-Methyl-N-methoxy-2-[1-(naphth-2-ylmethyl)-1H-imidazol-5-yl]acetamide

30

To a solution of 2-[1-(naphth-2-ylmethyl)-1H-imidazol-5-yl]acetic acid hydrochloride (0.819 mg, 2.70 mmol) in dimethylformamide (15 ml) was added sequentially N, O-dimethylhydroxylamine hydrochloride (293 mg, 3.0 mmol), 3-

- 109 -

hydroxy-1,2,3-benzotriazin-4(3H)-one (HOObt) (489 mg, 3.0 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (575 mg, 3.0 mmol) and triethylamine (1.67 ml, 12.0 mmol). This mixture was stirred at room temperature for 18h. After this time, sat. aq. NaHCO₃ (30ml) and water (30ml) were added and the mixture was extracted with methylene chloride (2 x 50ml). The combined organic extracts were washed with brine (50 ml) and the solvent evaporated in vacuo. The residue was purified by flash chromatography eluting with 2-4% methanol/methylene chloride gradient to provide the title compound as an oil.

¹HNMR (CDCl₃, 400 MHz) δ 7.80(2H, m), 7.74(1H, m), 7.56(1H, s), 7.47(3H, m), 7.22(1H, d, J=8.6Hz), 6.97(1H, s), 5.37(2H, s), 3.58(2H, s), 3.51(3H, s) and 3.12(3H, s)ppm.

15 **Step B:** Preparation of 2-[1-(Naphth-2-ylmethyl)(1H-imidazol-5-yl)]acetaldehyde

To a suspension of lithium aluminum hydride (40.8mg, 1.07mmol) in tetrahydrofuran (5ml) cooled to -45°C over a dry ice/acetone bath was added a solution of the product from Step A (243 mg, 0.895 mmol) in tetrahydrofuran (5 ml) via canula at such a rate to maintain temperature the temperature at <-35°C. After the addition was complete, the solution was allowed to warm to +5°C and then recooled to -35°C. To this solution was added a solution of potassium bisulfate (272 mg) in water (1 ml). This mixture was stirred for 30 min at room temperature and filtered through celite. The celite pad was washed with ethyl acetate (25 ml). The combined filtrates were washed with sat. sodium bicarbonate (10 ml) and then water (10ml). The organic layer was dried over magnesium sulfate, filtered and evaporated to give the title compound as a clear oil.

¹HNMR (CDCl₃, 400 MHz) δ 9.50(1H, t, 2Hz), 7.85-7.70(3H, m), 7.64(1H, s), 7.53-7.40(3H, m), 7.16(1H, d, J=12Hz), 7.06(1H, s), 5.20(2H, s) and 3.53(2H, m)ppm.

- 110 -

Step C: Preparation of 2(S)-Butyl-1-[(1-naphth-2-ylmethyl)-1H-imidazol-5-yl]ethyl]-4-(1-naphthoyl)piperazine
ditrifluoroacetate

To a solution of the product from Step B (58.4 mg, 0.232 mmol) and 3(S)-butyl-1-naphthoylpiperazine hydrochloride (96.5 mg, 0.279 mmol) in 1,2-dichloroethane (10ml) and dimethylformamide (5ml) was added 3Å molecular sieves (250mg) and sodium triacetoxyborohydride (236.5 mg, 1.12 mmol). This mixture was stirred at room temperature for 18 h. After this time, the mixture was filtered. The filtrate was diluted with methylene chloride (100ml) and washed with sat sodium bicarbonate (50ml). The organic layer was dried (MgSO₄), filtered and evaporated in vacuo. The residue was purified first by flash chromatography eluting with 2-5% methanol / methylene chloride and then by Prep HPLC (Chromatography A) to provide the title compound.

¹H NMR (CD₃OD, 400 MHz) δ 9.04(1H, s), 8.17-7.30(15H, m), 5.65(2H, s), 4.6-2.2(11H, m) and 1.6-0.2(9H, m) ppm.
FAB HRMS exact mass calc'd for C₃₅H₃₉N₄O 531.312387 (MH⁺), found 531.313011.

EXAMPLE 19

1-(2(R)-Amino-3-hydroxypropyl)-2(S)-butyl-4-(1-naphthoyl)piperazine
bis trifluoroacetate salt

Step A: N-Boc-O-Benzylserine-(N'-Methoxy) methyl amide

N-Boc-O-Benzylserine (Bachem; 5.0g, 16.9 mmol) and HOBT (2.29g, 16.9 mmol) were dissolved in dry DMF (100 mL) under argon. To this solution was added N,O-dimethylhydroxyl amine hydrochloride (1.98 g, 20.3 mmol) and then, at 0°C, EDC hydrochloride (3.56 g, 18.6 mmol). 4-Methylmorpholine was added to bring the pH to ~7 (4.5 mL) and the mixture was stirred at room temperature for 3h. The solution was diluted with EtoAc and poured into 0.5N HCl. After extraction with EtoAc (twice), the organic layers

- 111 -

were washed with water then brine, dried (MgSO₄) and evaporated in vacuo to give a pale yellow oil. Column chromatograph (silica gel; hexane/EtoAc 2:1) gave the title compound as an oil R_f (silica; hexane/EtoAc 1:1) = 0.45.

5

Step B: N-Boc-O-Benzyl-serine aldehyde

The amide from Step A (5.7 g, 16.9 mmol) in dry ether (20 mL) was added dropwise to a suspension of LAH (0.705 g, 18.5 mmol) in 80 mL of ether at -50°C. After the addition was complete, the
10 solution was stirred at 0°C for 45 min. then cooled back to -50°C and a solution of KHSO₄ (4 g in 11 mL H₂O) was added slowly. This mixture was then stirred at room temperature for 1 h, filtered through celite, washed successively with 10% citric acid solution, saturated NaHCO₃ solution and brine, dried (MgSO₄) and evaporated to give an
15 oil which was used as such in the next step. R_f (silica; hexane/EtoAc 2:1) = 0.58.

Step C: 1-(2(R)-N-Boc-Amino-3-benzyloxy propyl)-2(S)-butyl-4-(1-naphthoyl) piperazine

A solution of the piperazine hydrochloride from Example 1, Step E (1.7 g, 574 mmol) in CH₂Cl₂ (25 mL) was adjusted to pH 6 using Et₃N then freshly ground and activated 4 Å sieves were added followed by sodium triacetoxyborohydride (4.85 g, 22.9 mmol). The aldehyde from Step B (2.08g, 747 mmol) dissolved in 20 mL CH₂Cl₂
20 was added dropwise at 0°C over 20 min. then the mixture was stirred at room temperature for 16 h. After this time, the mixture was filtered through celite, diluted with EtoAc and washed successively with H₂O, KHSO₄ solution, NaHCO₃ solution and then brine. The dried (MgSO₄) solution was evaporated in vacuo to give a pale yellow oil
25 which was subjected to column chromatography (silica gel; hexane/EtOAc 1:1) to give the title compound as a foam. R_f (silica; hexane/EtoAc 2:1) = 0.15.
30

- 112 -

Step D: 1-(2(R)-N-Boc-Amino-3-hydroxy propyl)-2-(S)-butyl-4-(1-naphthoyl) piperazine

The benzyl ether from Step C (700 mg, 1.25 mmol) was dissolved in 20 mL MeOH with 150 μ L of acetic acid and then 20% Pd(OH)₂ on carbon (500 mg) was added and the mixture hydrogenated at 50 psi for 16 hr. After filtration through celite, the solvent was removed and the residue was chromatographed on silica gel (EtoAc/hexane 1:1 then 5% MeOH/EtoAc) to give the title compound as an oil.

Step E: 1-(2(R)-Amino-3-hydroxy propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine bis trifluoroacetate salt

A solution of the N-Boc amine from Step D (140 mg) in 10 mL EtoAc was treated with HCl (gas) until saturated. After 5 min, the solution was purged with argon then the solvent was removed to give a solid which was purified by preparative HPLC (C-18 column; H₂O/CH₃CN with 0.1% TGA; gradient). The aqueous solution was frozen and lyophilized to give the title compound as a hygroscopic powder. FAB mass spectrum $m/z = 370$ (M+1).

Analysis calculated for C₂₂H₃₂N₃O₂•2.35 TFA
C, 50.31; H, 5.27; N, 6.59
Found: C, 50.28; H, 5.49; N, 6.70

EXAMPLE 20

1-(2(R)-Amino-4-hydroxybutyl)-2(S)-butyl-4-(1-naphthoyl)piperazine bishydrochloride salt

Following the procedure of Example 19, Steps A to E but using N-Boc-O-benzyl homoserine (Bachem) as starting material, the title compound was obtained as the bis hydrochloride salt. FAB Mass spectrum, $m/z = 384$ (M+1).

Analysis calculated for C₂₃H₃₃N₃O₂•1.5 HCl
C, 54.04; H, 7.20; N, 8.22
Found: C, 53.95; H, 7.23; N, 8.50

- 113 -

EXAMPLE 21

1-(2-Amino-3-(2-benzyloxyphenyl)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine bistrifluoroacetate salt

5 Step A: D,L-N-Boc-ortho-tyrosine methyl ester

The title compound was prepared as a crystalline solid from D,L-orthotyrosine (Sigma) in two steps ((Boc)₂O/K₂CO₃ in THF/H₂O followed by diazomethane in EtOAc).

10 Step B: 3-(2-Hydroxyphenyl)-2-(N-Boc-Amino)propanol

To a solution of the ester from Step A (1.34 g, 4.54 mmol) in THF (20 mL) at 0°C was added LAH (400 mg, 10.5 mmol) in portions. After 4 hs at room temperature, 0.4 mL H₂O was added dropwise followed by 0.4 mL 1N NaOH and then 1.2 mL H₂O. The slurry was stirred for 1 h, filtered through celite rinsing with THF and the solvent was removed chromatography of the residue (silica gel; hexane/EtOAc 1:1) gave the title compound as a solid. R_f (silica; hexane/EtOAc 1:1) = 0.45.

20 Step C: 3-(2-Benzyloxyphenyl)-2-(N-Boc-amino)propanol

A mixture of the alcohol from Step B (280 mg, 1.05 mmol) benzyl bromide (150 µL, 1.26 mmol) and Cs₂CO₃ (513 mg, 1.57 mmol) in DMF (10 mL) was stirred at room temperature under argon for 16 h. The mixture was poured into H₂O, extracted twice with EtOAc, washed with water then brine, dried (MgSO₄) and evaporated to give an oil. Purification by chromatography (silica gel; hexane/EtOAc 2:1) gave the title compound as an oil. R_f (silica; hexane/EtOAc 2:1) = 0.36.

30 Step D: 3-(2-Benzyloxyphenyl)-2-(N-Boc-amino)propanol

To a solution of the alcohol from Step C (280 mg, 0.78 mmol) in 3 mL DMSO, 3 mL CH₂Cl₂ and 0.55 mL Et₃N at room temperature under argon was added pyridine SO₃ complex (500 mg,

- 114 -

3.14 mmol) and the mixture stirred for 1h. The solution was poured into saturated NaHCO₃ solution, extracted twice with EtOAc, washed with water then brine, dried (MgSO₄) and concentrated in vacuo. The title compound was thus obtained as an oil and was used as such in the following step. R_f (silica; hexane/EtOAc 4:1) = 0.41.

Step E: 1-(2-N-Boc-Amino-3-(2-benzyloxyphenyl)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine

The aldehyde from Step D and the piperazine hydrochloride from Example 1, Step E were coupled via reductive alkylation using the procedure described for Example 19, Step C. Purification by column chromatography (silica gel; hexane/EtOAc 3:2) gave the title compound as a 1:1 mixture of diastereomers. R_f (silica; hexane/EtOAc 1:1) = 0.51 and 0.45.

Step F: 1-(2-Amino-3-(2-benzyloxyphenyl)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine bis trifluoroacetate salt

A solution of the N-Boc amine from Step E (70 mg) in EtOAc (15 mL) was treated with HCl (g) until saturated. After 15 min., the solution was purged with argon and then the solvent removed to yield a solid. Purification by preparative HPLC (C-18 column, H₂O/CH₃CN with 0.1% TFA; gradient) afforded (after lyophilization) the title compound as a powder. FAB mass spectrum, m/z = 536 (M+1).

Analysis calculated for C₃₅H₄₁N₃O₂•2.2 TFA•0.35 H₂O
C, 59.68; H, 5.58; N, 5.30
Found: C, 59.70; H, 5.60; N, 5.56

- 115 -

EXAMPLE 22

1-(2-Amino-3-(2-hydroxyphenyl)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine bis trifluoroacetate salt, diastereomer A

5

Step A: 1-(2-N-Boc-Amino-3-(2-hydroxyphenyl)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine

A mixture of 1-(2-N-Boc-Amino-3-(2-benzyloxyphenyl)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine (from Example 19, Step E; 290 mg, 0.5 mmol), acetic acid (60 μ L, 1.0 mmol) and 20% Pd(OH)₂ on carbon (100 mg) in MeOH (20 mL) was hydrogenated at 67 psi for 5 h. The solution was filtered through celite and the solvent evaporated to give an oil which showed 2 well resolved spots on silica tlc. Column chromatography (silica gel; hexane:EtoAc 1:1) gave:

- 15 a) diastereomer A of the title compound RF (silica; hexane/EtoAc 1:1) = 0.49.
 b) diastereomer B of the title compound RF (silica; hexane/EtoAc 1:1) = 0.35.

20 Step B: 1-(2-Amino-3-(2-hydroxyphenyl)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine bis trifluoroacetate salt, diastereomer A

The N-Boc amine, diastereomer A from Step A, was deprotected using HCl (gas) in EtoAc. Removal of the solvent followed by preparative HPLC (C-18 column; H₂O/CH₃CN with 0.1% TFA; gradient) gave (after lyophilization) the title compound as a powder.

FAB mass spectrum, m/z = 446 (M+1).

Analysis calculated for C₂₈H₃₅N₃O₂•2 TFA

C, 57.05; H, 5.54; N, 6.24

30 Found: C, 57.08; H, 5.64; N, 6.32

EXAMPLE 23

1-[3-(4-imidazolyl)propyl]-2(S)-butyl-4-(1-naphthoyl)piperazine bis trifluoroacetate salt

35

- 116 -

Step A: Ethyl 3-(4-imidazolyl)propionate

The title compound was prepared from urocanic acid (Aldrich) in 2 steps using standard chemical procedures (esterification using HCl in EtOH followed by hydrogenation with 10% Pd-C in EtOH).
5 ^1H NMR (CDCl_3): δ 1.23 (3H, t), 2.65 (2H, t), 2.94 (2H, t), 4.15 (2H, q), 6.81 (1H, s), 7.56 (1H, s).

10 Step B: 3-(4-imidazolyl) propanol

To a solution of the ester from Step A (120 g, 71.4 mmol) in dry THF (230 mL) at 0°C was added LAH (2.99 g, 78.6 mmol) portionwise over 30 min. The mixture was stirred for 2 h at room temperature then cooled back to 0°C and H_2O (4.2 mL) was added
15 (care!) dropwise. This was followed by the dropwise addition of 10.6 mL of 1N NaOH then the resulting slurry was stirred at room temperature for 1 h. After filtration through celite washing with EtOAc, the solvent was removed to give the title compound as a clear oil (8.16 g).
20 ^1H NMR (CD_3OD): δ 1.84 (2H, pentet), 2.68 (2H, t), 3.58 (2H, t), 6.88 (1H, s), 7.80 (1H, s).

Step C: 3-(4-N-BOC-imidazolyl)propanol

The imidazole from Step B (166 mg, 1.32 mmol), (Boc) $_2\text{O}$ (302 mg, 1.38 mmol) and K_2CO_3 (190 mg, 1.38 mmol) were stirred in THF (10 mL) for 2 h. After filtration, the solvent was removed in vacuo to give the desired compound as an oil. R_f (silica; 5% MeOH/ CHCl_3) = 0.17. This was used as such in the next step.
25

30 Step D: 1-[3-(4-imidazolyl)propyl]-2(S)-butyl-4-(1-naphthoyl)-piperazine bis trifluoroacetate salt

The alcohol from Step C (~1.32 mmol) was dissolved in 4 mL DMSO, 4 mL CH_2Cl_2 and 0.92 mL Et_3N and then pyridine SO_3 complex (600 mg, 5.28 mmol) was added in portions. After 3 h, the

- 117 -

mixture was poured into EtOAc, extracted with saturated NaHCO₃ solution then brine, dried and evaporated to afford the corresponding aldehyde which was used without further purification.

- This aldehyde was coupled with the piperazine from Example 1, Step E by reductive alkylation following the procedure described for Example 19, Step C and the product was obtained by column chromatography (silica gel; 5% MeOH/CHCl₃). This product was dissolved in 10 mL EtOAc saturated with HCl (gas). The solvent was removed and the residue purified by preparative HPLC (C-18; H₂O/CH₃CN with 0.1% TFA; gradient). Lyophilization of the frozen aqueous solution gave the title compound as a hygroscopic powder. FAB mass spectrum, $m/z = 405$ (M+1).
Analysis calculated for C₂₅H₃₂N₄O•2.35 TFA 0.4 H₂O
 C, 52.48; H, 5.21; N, 8.24
15 Found: C, 52.45; H, 5.22; N, 8.27

EXAMPLE 24

- 20 2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(1-naphthylmethyl)imidazol-5-ylmethyl]-piperazine ditrifluoroacetic acid salt

- The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 1-bromomethylnaphthalene (0.046 g, 0.21 mmol) according to the procedure described in Example 4. The title
25 compound was obtained after purification by reverse phase preparative HPLC (gradient elution with 30%-75% acetonitrile/0.1% TFA; 70%-25% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 517. Anal. Calc. for C₃₄H₃₆N₄O · 0.05 H₂O · 2.0 TFA: C, 60.55; H, 5.22; N, 7.43. Found: C, 60.48; H, 5.12; N, 7.42.

30

- 118 -

EXAMPLE 25

2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(2-naphthylmethyl)imidazol-5-ylmethyl]-piperazine ditrifluoroacetic acid salt

- 5 The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 2-bromomethylnaphthalene (0.046 g, 0.21 mmol) according to the procedure described in Example 4. The title compound was obtained after purification by reverse phase preparative
- 10 HPLC (gradient elution with 30%-75% acetonitrile/0.1% TFA; 70%-25% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 517. Anal. Calc. for C₃₄H₃₆N₄O · 1.7 H₂O · 2.0 TFA: C, 58.87; H, 5.38; N, 7.23. Found: C, 58.90; H, 4.93; N, 7.13.

EXAMPLE 26

15

2(S)-*n*-Butyl-1-[1-(4-cyanobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine ditrifluoroacetic acid salt

- 20 The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 4-cyanobenzylbromide (0.041 g, 0.21 mmol) according to the procedure described in Example 4. The title compound was obtained after purification by reverse phase preparative
- 25 HPLC (gradient elution with 25%-65% acetonitrile/0.1% TFA; 75%-35% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 492. Anal. Calc. for C₃₁H₃₃N₅O · 0.35 H₂O · 2.0 TFA: C, 57.91; H, 4.96; N, 9.65. Found: C, 57.93; H, 4.91; N, 9.55.

EXAMPLE 27

30

2(S)-*n*-Butyl-1-[1-(4-methoxybenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine ditrifluoroacetic acid salt

- The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124

- 119 -

g, 0.200 mmol) and 4-methoxybenzylchloride (0.041 mL, 0.21 mmol) according to the procedure described in Example 4, with the addition of potassium iodide (100 mg) to the reaction mixture. The title compound was obtained after purification by reverse phase preparative HPLC (gradient elution with 25%-65% acetonitrile/0.1% TFA; 75%-35% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 497. Anal. Calc. for $C_{31}H_{36}N_4O_2 \cdot 1.7 H_2O \cdot 2.0 TFA$: C, 53.40; H, 5.30; N, 7.12. Found: C, 53.37; H, 4.78; N, 7.00.

10

EXAMPLE 28

2(S)-*n*-Butyl-1-[1-(3-methyl-2-butenyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine ditrifluoroacetic acid salt

The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 4-bromo-2-methyl-2-butene (0.024 mL, 0.21 mmol) according to the procedure described in Example 4. The title compound was obtained after purification by reverse phase preparative HPLC (gradient elution with 5%-95% acetonitrile/0.1% TFA; 95%-5% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 445. Anal. Calc. for $C_{28}H_{36}N_4O \cdot 1.8 H_2O \cdot 2.0 TFA$: C, 54.51; H, 5.95; N, 7.95. Found: C, 54.54; H, 5.39; N, 7.73.

25

EXAMPLE 29

2(S)-*n*-Butyl-1-[1-(4-fluorobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine ditrifluoroacetic acid salt

The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 4-fluorobenzylbromide (0.026 mL, 0.21 mmol) according to the procedure described in Example 4. The title compound was obtained after purification by reverse phase preparative HPLC (gradient elution with 25%-65% acetonitrile/0.1% TFA; 75%-35% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms

- 120 -

(m+1) 485. Anal. Calc. for $C_{30}H_{33}FN_4O \cdot 3.0 H_2O \cdot 2.0 TFA$: C, 53.26; H, 5.39; N, 7.31. Found: C, 53.21; H, 4.56; N, 7.08.

EXAMPLE 30

5 2(S)-n-Butyl-1-[1-(4-chlorobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine ditrifluoroacetic acid salt

The title compound was prepared from 2(S)-n-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 4-chlorobenzylchloride (0.034 mg, 0.21 mmol) according to the procedure described in Example 4, with the addition of sodium iodide (100 mg) to the reaction mixture. The title compound was obtained after purification by reverse phase preparative HPLC (gradient elution with 25%-65% acetonitrile/0.1% TFA; 75%-35% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 15 501. Anal. Calc. for $C_{30}H_{33}ClN_4O \cdot 4.8 H_2O \cdot 2.0 TFA$: C, 50.07; H, 5.51; N, 6.87. Found: C, 50.10; H, 4.25; N, 6.48.

EXAMPLE 31

20 1-[1-(4-Bromobenzyl)imidazol-5-ylmethyl]-2(S)-n-butyl-4-(1-naphthoyl)piperazine ditrifluoroacetic acid salt

The title compound was prepared from 2(S)-n-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 4-bromobenzylbromide (0.053 mg, 0.21 mmol) according to the procedure described in Example 4, with the addition of sodium iodide (100 mg) to the reaction mixture. The title compound was obtained after purification by reverse phase preparative HPLC (gradient elution with 30%-65% acetonitrile/0.1% TFA; 70%-35% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 25 545. Anal. Calc. for $C_{30}H_{33}BrN_4O \cdot 1.7 H_2O \cdot 2.0 TFA$: C, 50.78; H, 4.81; N, 6.97. Found: C, 50.81; H, 4.39; N, 6.88.

- 121 -

EXAMPLE 32

2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(4-trifluoromethylbenzyl)imidazol-5-ylmethyl]-piperazine ditrifluoroacetic acid salt

5 The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 4-trifluoromethylbenzylbromide (0.053 mg, 0.21 mmol) according to the procedure described in Example 4. The title compound was obtained after purification by reverse phase preparative
10 HPLC (gradient elution with 30%-65% acetonitrile/0.1% TFA; 70%-35% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 535. Anal. Calc. for C₃₀H₃₃BrN₄O · 2.0 TFA: C, 55.12; H, 4.63; N, 7.35. Found: C, 57.46; H, 4.98; N, 7.84.

15 EXAMPLE 33

2(S)-*n*-Butyl-1-[1-(4-methylbenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)-piperazine ditrifluoroacetic acid salt

The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 4-methylbenzylbromide (0.029 mL, 0.21 mmol) according to the procedure described in Example 4, with the addition of sodium iodide (100 mg) to the reaction mixture. The title compound was obtained after purification by reverse phase preparative HPLC (gradient elution with 30%-65% acetonitrile/0.1% TFA; 70%-35% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms ($m+1$) 481. Anal. Calc. for $C_{31}H_{36}N_4O \cdot 2.6 H_2O \cdot 2.0 TFA$: C, 55.64; H, 5.76; N, 7.42. Found: C, 55.61; H, 5.09; N, 7.43.

- 122 -

EXAMPLE 34

2(S)-*n*-Butyl-1-[1-(3-methylbenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)-piperazine ditrifluoroacetic acid salt

- 5 The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 3-methylbenzylbromide (0.029 mL, 0.21 mmol) according to the procedure described in Example 4. The title compound was obtained after purification by reverse phase preparative
10 HPLC (gradient elution with 25%-65% acetonitrile/0.1% TFA; 75%-35% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 481. Anal. Calc. for C₃₁H₃₆N₄O · 2.0 TFA: C, 67.26; H, 6.92; N, 10.12. Found: C, 69.60; H, 6.98; N, 10.51.

EXAMPLE 35

15

1-[1-(4-Phenylbenzyl)imidazol-5-ylmethyl]-2(S)-*n*-butyl-4-(1-naphthoyl)-piperazine ditrifluoroacetic acid salt

- The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124 g, 0.200 mmol) and 4-phenylbenzylbromide (0.029 mL, 0.21 mmol) according to the procedure described in Example 4. The title compound was obtained after purification by reverse phase preparative
20 HPLC (gradient elution with 30%-65% acetonitrile/0.1% TFA; 70%-35% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms
25 (m+1) 543. Anal. Calc. for C₃₆H₃₈N₄O · 4.95 H₂O · 2.0 TFA: C, 55.87; H, 5.85; N, 6.52. Found: C, 55.55; H, 4.58; N, 6.23.

EXAMPLE 36

30

2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(2-phenylethyl)imidazol-5-ylmethyl]-piperazine ditrifluoroacetic acid salt

- The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124

- 123 -

g, 0.200 mmol) and 2-phenylethylbromide (0.029 mL, 0.21 mmol) according to the procedure described in Example 4, except with the addition of sodium iodide (120 mg) and refluxing for 12 h. The title compound was obtained after purification by reverse phase preparative
5 HPLC (gradient elution with 30%-65% acetonitrile/0.1% TFA; 70%-35% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 481. Anal. Calc. for $C_{36}H_{38}N_4O \cdot 4.20 H_2O \cdot 2.0 TFA$: C, 53.60; H, 5.96; N, 7.14. Found: C, 53.54; H, 4.86; N, 6.86.

10

EXAMPLE 37

2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(4-trifluoromethoxy)imidazol-5-ylmethyl]piperazine ditrifluoroacetic acid salt

The title compound was prepared from 2(S)-*n*-butyl-1-[5-(3-triphenylmethylimidazolyl)methyl]-4-(1-naphthoyl)piperazine (0.124
15 g, 0.200 mmol) and 4-trifluoromethoxybenzylbromide (0.032 mL, 0.21 mmol) according to the procedure described in Example 4, except with the addition of sodium iodide (120 mg) and refluxing for 12 h. The title compound was obtained after purification by reverse phase
20 preparative HPLC (gradient elution with 35%-70% acetonitrile/0.1% TFA; 65%-30% 0.1% aqueous TFA over 50 min.) and lyophilization. FAB ms (m+1) 551. Anal. Calc. for $C_{31}H_{33}F_3N_4O_2 \cdot 4.00 H_2O \cdot 2.0 TFA$: C, 49.42; H, 5.09; N, 6.59. Found: C, 48.95; H, 4.06; N, 6.26.

25

EXAMPLE 38

Preparation of 1-[[1-(4-cyanobenzyl)-1H-imidazol-5-yl]acetyl]-2(S)-*n*-butyl-4-(1-naphthoyl)piperazine trifluoroacetate

30 Step A: Preparation of 1H-Imidazole-4- acetic acid methyl ester hydrochloride.

A solution of 1H-imidazole-4-acetic acid hydrochloride (4.00g, 24.6 mmol) in methanol (100 ml) was saturated with gaseous hydrogen chloride. The resulting solution was allowed to stand at room

- 124 -

temperature (RT) for 18hr. The solvent was evaporated in vacuo to afford the title compound as a white solid.

^1H NMR(CDCl_3 , 400 MHz) δ 8.85(1H, s), 7.45(1H, s), 3.89(2H, s) and 3.75(3H, s) ppm.

5

Step B: Preparation of 1-(Triphenylmethyl)-1H-imidazol-4-ylacetic acid methyl ester.

To a solution of the product from Step A (24.85g, 0.141mol) in dimethyl formamide (DMF) (115ml) was added triethylamine (57.2 ml, 0.412mol) and triphenylmethyl bromide(55.3g, 0.171mol) and the suspension was stirred for 24hr. After this time, the reaction mixture was diluted with ethyl acetate (EtOAc) (1 L) and water (350 ml). The organic phase was washed with sat. aq. NaHCO_3 (350 ml), dried (Na_2SO_4) and evaporated in vacuo. The residue was purified by flash chromatography (SiO_2 , 0-100% ethyl acetate in hexanes; gradient elution) to provide the title compound as a white solid.

10

^1H NMR (CDCl_3 , 400 MHz) δ 7.35(1H, s), 7.31(9H, m), 7.22(6H, m), 6.76(1H, s), 3.68(3H, s) and 3.60(2H, s) ppm.

15

Step C: Preparation of [1-(4-cyanobenzyl)-1H-imidazol-5-yl]acetic acid methyl ester.

To a solution of the product from Step B (8.00g, 20.9mmol) in acetonitrile (70 ml) was added bromo-p-toluenitrile (4.10g, 20.92 mmol) and heated at 55°C for 3 hr. After this time, the reaction was cooled to room temperature and the resulting imidazolium salt (white precipitate) was collected by filtration. The filtrate was heated at 55°C for 18hr. The reaction mixture was cooled to room temperature and evaporated in vacuo. To the residue was added EtOAc (70 ml) and the resulting white precipitate collected by filtration. The precipitated imidazolium salts were combined, suspended in methanol (100 ml) and heated to reflux for 30min. After this time, the solvent was removed in vacuo, the resulting residue was suspended in EtOAc (75ml) and the solid isolated by filtration and washed (EtOAc). The

20

25

30

- 125 -

solid was treated with sat aq NaHCO₃ (300ml) and CH₂Cl₂ (300ml) and stirred at room temperature for 2 hr. The organic layer was separated, dried (MgSO₄) and evaporated in vacuo to afford the title compound as a white solid :

- 5 ¹H NMR(CDCl₃, 400 MHz) δ 7.65(1H, d, J=8Hz), 7.53(1H, s), 7.15(1H, d, J=8Hz), 7.04(1H, s), 5.24(2H, s), 3.62(3H, s) and 3.45(2H, s) ppm.

Step D: Preparation of [1-(4-cyanobenzyl)-1H-imidazol-5-yl]acetic acid

- 10 A solution of [1-(4-cyanobenzyl)-1H-imidazol-5-yl]acetic acid methyl ester (4.44g, 17.4mmol) in THF (100ml) and 1 M lithium hydroxide (17.4 ml, 17.4 mmol) was stirred at RT for 18 hr. 1 M HCl (17.4 ml) was added and the THF was removed by evaporation in vacuo. The aqueous solution was lyophilised to afford the title
15 compound containing lithium chloride as a white solid.
¹H NMR(CD₃OD, 400 MHz) δ 8.22(1H, s), 7.74(1H, d, J=8.4Hz), 7.36(1H, d, J=8.4Hz), 7.15(1H, s), 5.43(2H, s) and 3.49(2H, s) ppm.

Step E: Preparation of 1-[[1-(4-cyanobenzyl)-1H-imidazol-5-yl]acetyl]-2(S)-n-butyl-4-(1-naphthoyl)piperazine trifluoroacetate

- 20 To a solution of the acid from step D (100 mg, 0.35 mmol), the amine hydrochloride salt from example step (117 mg, 0.35mmol), HOOBT (58 mg, 0.35 mmol), and triethylamine (0.123ml, 0.88 mmol) in DMF (2ml) was added EDC (75 mg, 0.38mmol). The
25 reaction was stirred at room temperature for 16hrs, diluted with EtOAc and the organic layer washed with sat. aq NaHCO₃ , brine , dried (Na₂SO₄). and evaporated in vacuo. The residue was purified by preparative HPLC (C-18; 95:5 to 5:95 water:CH₃CN containing 0.1% trifluoroacetic acid; gradient elution). Lyophilisation of the collected
30 fractions afforded the title compound as a white solid.
¹H NMR(CD₃OD, 400 MHz) δ 9.00-8.90(1H, m), 8.05-7.94(2H,m), 7.94-7.40(10H,m), 5.60-5.40(2H,m), 5.00-2.80(9H,m), 1.90-1.15(4H,m) and 1.05-0.40(5H, m)ppm.

- 126 -

Anal. calc'd for $C_{32}H_{33}N_5O_2 \cdot 1.40 \text{ TFA} \cdot 0.55H_2O$: C, 60.65; H, 5.19; N, 10.16. Found: C, 60.66; H, 5.17; N, 10.06.

EXAMPLE 39

5

5(S)-*n*-Butyl-1-(2,3-dimethylphenyl)-4-(4-imidazolylmethyl)piperazin-2-one ditrifluoroacetic acid salt

10

Step A: N-Methoxy-N-methyl 2(S)-(tert-butoxycarbonylamino)-hexanamide

15

2(S)-(tert-Butoxycarbonylamino)hexanoic acid (24.6 g, 0.106 mol), N,O-dimethylhydroxylamine hydrochloride (15.5 g, 0.15 mol), EDC hydrochloride (22.3 g, 0.117 mol) and HOBT (14.3 g, 0.106 mol) were stirred in dry, degassed DMF (300 mL) at 20°C under nitrogen. N-Methylmorpholine was added to obtain pH 7. The reaction was stirred overnight, the DMF distilled under high vacuum, and the residue partitioned between ethyl acetate and 2% potassium hydrogen sulfate. The organic phase was washed with saturated sodium bicarbonate, water, and saturated brine, and dried with magnesium sulfate. The solvent was removed in vacuo to give the title compound.

20

Step B: 2(S)-(tert-Butoxycarbonylamino)hexanal

25

A mechanically stirred suspension of lithium aluminum hydride (5.00 g, 0.131 mol) in ether (250 mL) was cooled to -45°C under nitrogen. A solution of the product from Step A (28.3 g, 0.103 mol) in ether (125 mL) was added, maintaining the temperature below -35°C. When the addition was complete, the reaction was warmed to 5°C, then re-cooled to -45°C. A solution of potassium hydrogen sulfate (27.3 g, 0.200 mol) in water was slowly added, maintaining the temperature below -5°C. After quenching, the reaction was stirred at room temperature for 1h. The mixture was filtered through Celite, the ether evaporated, and the remainder partitioned between ethyl acetate and 2% potassium hydrogen sulfate. After washing with saturated

30

- 127 -

brine, drying over magnesium sulfate and solvent removal, the title compound was obtained.

Step C: N-(2,3-Dimethylphenyl)-2(S)-(tert-butoxycarbonylamino)-hexanamine

5 2,3-Dimethylaniline (8.32 mL, 68.3 mmol) was dissolved in dichloroethane under nitrogen. Acetic acid was added to obtain pH 5, and sodium triacetoxyborohydride (17.2 g, 80.8 mmol) and crushed molecular sieves (4 g) were added. A solution of the product from Step
10 B (13.3 g, 62.1 mmol) in dichloroethane (80 mL) was added slowly dropwise at 20°C. The reaction was stirred overnight, then quenched with saturated sodium bicarbonate solution. The aqueous layer was removed, the organic phase washed with saturated brine and dried over magnesium sulfate. Crystallization from hexane gave the title
15 compound.

Step D: 4-tert-Butoxycarbonyl-5(S)-n-butyl-1-(2,3-dimethylphenyl)piperazin-2-one

20 A solution of the product from Step C (8.50 g, 26.5 mmol) in ethyl acetate (250 mL) was vigorously stirred at 0°C with saturated sodium bicarbonate (150 mL). Chloroacetyl chloride (2.33 mL, 29.1 mmol) was added, and the reaction stirred at) 0°C for 1h. The layers were separated, and the ethyl acetate phase was washed with saturated brine, and dried over magnesium sulfate. The crude product was
25 dissolved in DMF (300 mL) and cooled to 0°C under nitrogen. Sodium hydride (1.79 g, 60% dispersion in oil, 44.9 mmol) was added portionwise to maintain moderate hydrogen evolution. After 30 min, an additional amount of sodium hydride was added (0.8 g). The reaction was stirred another 30 min, then quenched with saturated
30 ammonium chloride. The DMF was distilled *in vacuo*, and the residue partitioned between ethyl acetate and water. The organic phase was washed with water, saturated brine, and dried over magnesium sulfate. The crude product was chromatographed on silica gel with 20-30% ethyl acetate in hexane to obtain the title compound.

- 128 -

Step E: 5(S)-*n*-Butyl-1-(2,3-dimethylphenyl)-4-[4-(1-triphenylmethylimidazolyl)methyl]piperazin-2-one

A solution of the product from Step D (0.570 g, 1.58 mmol) in ethyl acetate (50 mL) was cooled to -15°C under nitrogen. HCl gas was bubbled through for 15 min, and the reaction solution warmed to 0°C for 2h. The solvent was removed *in vacuo*, and the resulting solid was dissolved in dichloroethane (20 mL). Sodium triacetoxyborohydride (0.502 g, 2.37 mmol) and 1-triphenylmethyl-4-imidazolyl carboxaldehyde (0.534 g, 1.58 mmol) was added. The reaction was stirred overnight at 20°C then poured into saturated sodium bicarbonate solution. The organic phase was washed with saturated brine and dried over magnesium sulfate. Silica gel chromatography using 4% methanol in dichloromethane as eluant yielded the title compound.

Step F: 5(S)-*n*-Butyl-1-(2,3-dimethylphenyl)-4-(4-imidazolylmethyl)piperazin-2-one ditrifluoroacetic acid salt

To a solution of the compound from Step E (0.233 g, 0.40 mmol) in dichloromethane (6 mL) was added triethylsilane (0.254 mL) and trifluoroacetic acid (2 mL) were added, and the reaction stirred at 20°C for 2h. The volatiles were removed *in vacuo*, and the residue partitioned between hexane and water-methanol. The aqueous phase was injected onto a preparative HPLC column and purified with a mixed gradient of 15%-60% acetonitrile/0.1% TFA; 85%-40% 0.1% aqueous TFA over 50 min. The title compound was isolated after lyophilization. FAB ms (*m*+1) 341. Anal. Calc. for C₂₀H₂₈N₄O · 2.0 TFA: C, 50.80; H, 5.15; N, 9.87. Found: C, 51.31; H, 5.41; N, 10.11.

- 129 -

EXAMPLE 40

5(S)-*n*-Butyl-4-[1-(4-cyanobenzyl)imidazol-5-ylmethyl]-1-(2,3-dimethylphenyl)piperazin-2-one ditrifluoroacetic acid salt

5 4-Cyanobenzylbromide (0.043 g, 0.22 mmol) was added at 20°C to a solution of 5(S)-*n*-butyl-1-(2,3-dimethylphenyl)-4-[4-(1-triphenylmethylimidazolyl)methyl]piperazin-2-one (0.120 g, 0.21 mmol) from Example 39, Step E, in acetonitrile (10 mL). After 48 h, the solvent was removed *in vacuo*, and the crude product dissolved in 10 dichloromethane (6 mL). Triethylsilane (0.13 mL) and trifluoroacetic acid (2 mL) were added, and the reaction stirred at 20°C for 2h. The volatiles were removed *in vacuo*, and the residue partitioned between hexane and water-methanol. The aqueous phase was injected onto a reverse phase preparative HPLC column and purified with a mixed 15 gradient of 30%-60% acetonitrile/0.1% TFA; 70%-40% 0.1% aqueous TFA over 50 min. The title compound was isolated after lyophilization from water-acetonitrile solution. FAB ms (m+1) 456. Anal. Calc. for C₂₈H₃₃N₅O · 0.7 H₂O · 2.0 TFA: C, 55.28; H, 5.13; N, 10.07. Found: C, 55.27; H, 5.20; N, 10.41.

EXAMPLE 41

4-[1-(4-Cyanobenzyl)imidazol-5-ylmethyl]-1-(2,3-dimethylphenyl)-5(S)-(2-methoxyethyl)piperazin-2-one ditrifluoroacetic acid salt

25 Step A: N-Methoxy-N-methyl 4-benzyloxy-2(S)-(tert-butoxycarbonylamino)butanamide
 4-Benzyloxy-2(S)-(tert-butoxycarbonylamino)butanoic acid (1.00 g, 3.23 mmol) was converted to the title compound following the 30 procedure described in Example 39, Step A, using EDC · HCl (0.680 g, 3.55 mmol). HOBt (0.436 g, 3.23 mmol) and N,O-dimethylhydroxylamine hydrochloride (0.473 g, 4.85 mmol) in DMF (50 mL) at pH 7. After workup, the title compound was obtained as a clear gum.

- 130 -

Step B: 4-(1-Benzyloxyethyl)-2(S)-(tert-butoxycarbonylamino)butanal

5 The title compound was obtained by lithium aluminum hydride reduction of the product of Step A using the procedure described in Example 39, Step B.

Step C: N-(2,3-Dimethylphenyl)-4-(2-benzyloxyethyl)-2-(S)-(tert-butoxycarbonylamino)butanamine

10 The title compound was prepared from the product of Step C according to the procedure described in Example 39, Step B, using 2,3-dimethylaniline (0.505 mL, 4.14 mmol), sodium triacetoxymethylborohydride (1.20 g, 5.65 mmol) and crushed molecular sieves (1 g) at pH 5 in dichloroethane (20 mL). The title compound was
15 obtained after purification on silica gel, eluting with 15% ethyl acetate in hexane.

Step D: 5(S)-(2-Benzyloxyethyl)-4-tert-butoxycarbonyl-1-(2,3-dimethylphenyl)piperazin-2-one

20 The title compound was prepared from the product of Step C according to the procedure described in Example 39, Step D, using chloroacetyl chloride (0.21 mL, 2.57 mmol) in 60 mL 1:1 ethyl acetate:saturated sodium bicarbonate, followed by reaction of the crude product with sodium hydride (0.373 g, 60% dispersion in oil, 9.32
25 mmol) in DMF (30 mL). After workup, the crude product was chromatographed on silica gel with 30% ethyl acetate in hexane to obtain the title compound.

Step E: 4-tert-Butoxycarbonyl-1-(2,3-dimethylphenyl)-5(S)-(2-hydroxyethyl)piperazin-2-one

30 The product from Step D was dissolved in methanol (40 mL) and 10% Pd/C was added (0.160 g). The reaction was shaken under 60 psi hydrogen overnight. The catalyst was removed by filtration, and the solvent evaporated to give the title compound.

- 131 -

Step F: 4-*tert*-Butoxycarbonyl-1-(2,3-dimethylphenyl)-5(S)-(2-methoxyethyl)piperazin-2-one

The product from Step E (0.241 g, 0.688 mmol) was dissolved in DMF (10 mL) containing methyl iodide (0.21 mL, 3.44 mmol) and the stirred solution cooled to 0°C under nitrogen. Sodium hydride (0.070 g, 60% dispersion in oil, 1.72 mmol) was added and the reaction stirred for 1h. The reaction was quenched with water, and the DMF removed under vacuum. The residue was partitioned between ethyl acetate and water, and the organic phase washed with saturated brine and dried over magnesium sulfate. The crude product was chromatographed on silica gel with 40% ethyl acetate in hexane to give the title compound.

Step G: 1-(2,3-Dimethylphenyl)-5(S)-(2-methoxyethyl)-4-[4-(1-triphenylmethylimidazolyl)methyl]piperazin-2-one

The product from Step F (0.113 g, 0.312 mmol) was converted to the title compound according to the procedure described in Example 39, Step E, except using 30% trifluoroacetic acid in dichloromethane (10 mL) for 1 h for the initial deprotection. The volatiles were removed *in vacuo*, and the residue dissolved in dichloroethane. Triethylamine was added to obtain pH 5. Sodium triacetoxymethylborohydride (0.100 g, 0.468 mmol) and 1-triphenylmethyl-4-imidazolylcarboxaldehyde (0.1164 g, 0.343 mmol) was added. The reaction was stirred overnight at 20°C then poured into saturated sodium bicarbonate solution. The organic phase was washed with saturated brine and dried over magnesium sulfate. Silica gel chromatography using 5% methanol in chloroform as eluant yielded the title compound.

30

- 132 -

Step H: 4-[1-(4-Cyanobenzyl)imidazol-5-ylmethyl]-1-(2,3-dimethylphenyl)-5(S)-(2-methoxyethyl)piperazin-2-one
ditrifluoroacetic acid salt

The product from Step G (0.182 g, 0.312 mmol) was converted to the title compound according to the procedure described in Example 40, using 4-cyanobenzylbromide (0.061 g, 0.312 mmol) in acetonitrile (10 mL), followed by reaction of the crude imidazolium salt with triethylsilane (0.13 mL) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL). Purification was accomplished by reverse phase preparative HPLC with a mixed gradient of 0%-70% acetonitrile/0.1% TFA; 100%-30% 0.1% aqueous TFA over 60 min. The title compound was isolated after lyophilization from water. FAB ms (m+1) 458. Anal. Calc. for $C_{27}H_{31}N_5O_2 \cdot 0.35 H_2O \cdot 2.0 TFA$: C, 53.81; H, 4.91; N, 10.21. Found: C, 53.83; H, 4.95; N, 10.29.

EXAMPLE 42

Preparation of (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(methanesulfonyl)ethyl]-2-piperazinone
dihydrochloride

Step A: Preparation of 1-triphenylmethyl-4-(hydroxymethyl)-imidazole

To a solution of 4-(hydroxymethyl)imidazole hydrochloride (35.0 g, 260 mmol) in 250 mL of dry DMF at room temperature was added triethylamine (90.6 mL, 650 mmol). A white solid precipitated from the solution. Chlorotriphenylmethane (76.1 g, 273 mmol) in 500 mL of DMF was added dropwise. The reaction mixture was stirred for 20 hours, poured over ice, filtered, and washed with ice water. The resulting product was slurried with cold dioxane, filtered, and dried *in vacuo* to provide the titled product as a white solid which was sufficiently pure for use in the next step.

- 133 -

Step B: Preparation of 1-triphenylmethyl-4-(acetoxymethyl)-
imidazole

Alcohol from Step A (260 mmol, prepared above) was suspended in 500 mL of pyridine. Acetic anhydride (74 mL, 780 mmol) was added dropwise, and the reaction was stirred for 48 hours during which it became homogeneous. The solution was poured into 2 L of EtOAc, washed with water (3 x 1 L), 5% aq. HCl soln. (2 x 1 L), sat. aq. NaHCO₃, and brine; then dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the crude product. The acetate was isolated as a white powder which was sufficiently pure for use in the next reaction.

Step C: Preparation of 1-(4-cyanobenzyl)-5-(acetoxymethyl)-
imidazole hydrobromide

A solution of the product from Step B (85.8 g, 225 mmol) and α -bromo-*p*-tolunitrile (50.1 g, 232 mmol) in 500 mL of EtOAc was stirred at 60 °C for 20 hours, during which a pale yellow precipitate formed. The reaction was cooled to room temperature and filtered to provide the solid imidazolium bromide salt. The filtrate was concentrated *in vacuo* to a volume 200 mL, reheated at 60 °C for two hours, cooled to room temperature, and filtered again. The filtrate was concentrated *in vacuo* to a volume 100 mL, reheated at 60 °C for another two hours, cooled to room temperature, and concentrated *in vacuo* to provide a pale yellow solid. All of the solid material was combined, dissolved in 500 mL of methanol, and warmed to 60 °C. After two hours, the solution was reconcentrated *in vacuo* to provide a white solid which was triturated with hexane to remove soluble materials. Removal of residual solvents *in vacuo* provided the titled product hydrobromide as a white solid which was used in the next step without further purification.

- 134 -

Step D: Preparation of 1-(4-cyanobenzyl)-5-(hydroxymethyl)-
imidazole

To a solution of the acetate from Step C (50.4 g, 150 mmol) in 1.5 L of 3:1 THF/water at 0 °C was added lithium hydroxide monohydrate (18.9 g, 450 mmol). After one hour, the reaction was concentrated *in vacuo*, diluted with EtOAc (3 L), and washed with water, sat. aq. NaHCO₃ and brine. The solution was then dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the crude product as a pale yellow fluffy solid which was sufficiently pure for use in the next step without further purification.

Step E: Preparation of 1-(4-cyanobenzyl)-5-
imidazolecarboxaldehyde

To a solution of the alcohol from Step D (21.5 g, 101 mmol) in 500 mL of DMSO at room temperature was added triethylamine (56 mL, 402 mmol), then SO₃-pyridine complex (40.5 g, 254 mmol). After 45 minutes, the reaction was poured into 2.5 L of EtOAc, washed with water (4 x 1 L) and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the aldehyde as a white powder which was sufficiently pure for use in a subsequent step (Step L) without further purification.

Step F: Preparation of (S)-2-(*tert*-butoxycarbonylamino)-*N*-
methoxy-*N*-methyl-4-(methylthio)butanamide

L-*N*-Boc-methionine (30.0 g, 0.120 mol), *N,O*-dimethylhydroxylamine hydrochloride (14.1 g, 0.144 mol), EDC hydrochloride (27.7 g, 0.144 mol) and HOBT (19.5 g, 0.144 mol) were stirred in dry DMF (300 mL) at 20°C under nitrogen. More *N,O*-dimethylhydroxylamine hydrochloride (2.3 g, 23 mmol) was added to obtain pH 7-8. The reaction was stirred overnight, the DMF distilled to half the original volume under high vacuum, and the residue partitioned between ethyl acetate and sat. NaHCO₃ soln. The organic phase was washed with saturated sodium bicarbonate, water, 10% citric acid, and

- 135 -

brine, and dried with sodium sulfate. The solvent was removed *in vacuo* to give the title compound.

Step G: Preparation of (S)-2-(*tert*-butoxycarbonylamino)-4-(methylthio)butanal

5 A suspension of lithium aluminum hydride (5.02 g, 0.132 mol) in ether (500 mL) was stirred at room temperature for one hour. The solution was cooled to -50 °C under nitrogen, and a solution of the product from Step F (39.8 g, ca. 0.120 mol) in ether (200 mL) was
10 added over 30 min, maintaining the temperature below -40 °C. When the addition was complete, the reaction was warmed to 5°C, then recooled to -45°C. Analysis by tlc revealed incomplete reaction. The solution was rewarmed to 5 °C, stirred for 30 minutes, then cooled to -50 °C. A solution of potassium hydrogen sulfate (72 g, 0.529 mol) in
15 200 mL water was slowly added, maintaining the temperature below -20 °C. The mixture was warmed to 5 °C, filtered through Celite, and concentrated *in vacuo* to provide the title aldehyde.

Step H: Preparation of (S)-2-(*tert*-butoxycarbonylamino)-N-(3-chlorophenyl)-4-(methylthio)butanamine

20 To a solution of 3-chloroaniline (10.3 mL, 97.4 mmol), the product from Step G (23.9 g, 97.4 mmol), and acetic acid (27.8 mL, 487 mmol) in dichloroethane (250 mL) under nitrogen was added sodium triacetoxyborohydride (41.3 g, 195 mmol). The reaction was
25 stirred overnight, then quenched with saturated sodium bicarbonate solution. The solution was diluted with CHCl₃, and the organic phase was washed with water, 10% citric acid and brine. The solution was dried over sodium sulfate and concentrated *in vacuo* to provide the crude product (34.8 g) which was chromatographed on silica gel with
30 20% ethyl acetate in hexane to obtain the title compound.

- 136 -

Step I: Preparation of (S)-4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-5-[2-(methylthio)ethyl]piperazin-2-one

A solution of the product from Step H (22.0 g, 63.8 mmol) in ethyl acetate (150 mL) was vigorously stirred at 0°C with saturated sodium bicarbonate (150 mL). Chloroacetyl chloride (5.6 mL, 70.2 mmol) was added dropwise, and the reaction stirred at 0°C for 2h. The layers were separated, and the ethyl acetate phase was washed with 10% citric acid and saturated brine, and dried over sodium sulfate. After concentration *in vacuo*, the resulting crude product (27.6 g) was dissolved in DMF (300 mL) and cooled to 0°C under argon. Cesium carbonate (63.9 g, 196 mmol) was added, and the reaction was stirred for two days, allowing it to warm to room temperature. Another portion of cesium carbonate (10 g, 30 mmol) was added, and the reaction was stirred for 16 hours. The DMF was distilled *in vacuo*, and the residue partitioned between ethyl acetate and water. The organic phase was washed with saturated brine, and dried over sodium sulfate. The crude product was chromatographed on silica gel with 20-25% ethyl acetate in hexane to obtain the title compound.

Step J: Preparation of (S)-4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-5-[2-(methanesulfonyl)ethyl]piperazin-2-one

A solution of the product from Step I (14.2 g, 37 mmol) in methanol (300 mL) was cooled to 0 °C, and a solution of magnesium monoperoxyphthalate (54.9 g, 111 mmol) in 210 mL MeOH was added over 20 minutes. The ice bath was removed, and the solution was allowed to warm to room temperature. After 45 minutes, the reaction was concentrated *in vacuo* to half the original volume, then quenched by the addition of 2N Na₂S₂O₃ soln. The solution was poured into EtOAc and sat NaHCO₃ solution, and the organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the crude sulfone. This material was chromatographed on silica gel with 60-100% ethyl acetate in hexane to obtain the titled compound.

- 137 -

Step K: Preparation of (S)-1-(3-chlorophenyl)-5-[2-(methanesulfonyl)ethyl]piperazin-2-one

Through a solution of Boc-protected piperazinone from Step J (1.39 g, 3.33 mmol) in 30 mL of EtOAc at 0 °C was bubbled anhydrous HCl gas. The saturated solution was stirred for 35 minutes, then concentrated *in vacuo* to provide the hydrochloride salt as a white powder. This material was suspended in EtOAc and treated with dilute aqueous NaHCO₃ solution. The aqueous phase was extracted with EtOAc, and the combined organic mixture was washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The resulting amine was reconcentrated from toluene to provide the titled material suitable for use in the next step.

Step L: Preparation of (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(methanesulfonyl)ethyl]-2-piperazinone dihydrochloride

To a solution of the amine from Step K (898 mg, 2.83 mmol) and imidazole carboxaldehyde from Step E (897 mg, 4.25 mmol) in 15 mL of 1,2-dichloroethane was added sodium triacetoxyborohydride (1.21 g, 5.7 mmol). The reaction was stirred for 23 hours, then quenched at 0 °C with sat. NaHCO₃ solution. The solution was poured into CHCl₃, and the aqueous layer was back-extracted with CHCl₃. The combined organics were washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The resulting product was purified by silica gel chromatography (95:5:0.5-90:10:0 EtOAc:MeOH:NH₄Cl), and the resultant product was taken up in EtOAc/methanol and treated with 2.1 equivalents of 1 M HCl/ether solution. After concentrated *in vacuo*, the product dihydrochloride was isolated as a white powder.

30

- 138 -

EXAMPLE 43

Preparation of (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)ethyl]-2-piperazinone
5 dihydrochloride

Step A: Preparation of (S)-N-(*tert*-butoxycarbonyl)homoserine
lactone

To a solution of (S)-homoserine lactone hydrochloride
10 (11.0 g, 79.9 mmol) and di-*tert*-butylpyrocarbonate (19.2 g, 88.0 mmol) in 160 mL of dichloromethane at 0 °C was added diisopropylethylamine (13.9 mL, 79.9 mmol) over 3 min. The solution was allowed to warm to room temperature. After 3 hours, another
15 portion of di-*tert*-butylpyrocarbonate (1.75 g, 8.0 mmol) and diisopropylethylamine (0.70 mL, 4.0 mmol) were added, and the mixture was stirred for an additional 2.5 hours. The solution was washed with 10% citric acid, sat. NaHCO₃, and brine, dried (Na₂SO₄),
20 filtered, and concentrated *in vacuo*. The resulting material was purified by silica gel chromatography (50% EtOAc/hexane) to provide pure titled compound.

Step B: Preparation of (S)-N-(*tert*-butoxycarbonyl)homoserine
lactol

To a solution of the lactone from Step A (7.0 g, 35 mmol)
25 in 175 mL of THF at -78 °C was added diisobutylaluminum hydride (72.0 mL, 1M in THF, 72 mmol) dropwise, while maintaining the reaction temperature below -72 °C. After 3 hours, another portion of diisobutylaluminum hydride (10.0 mL, 10 mmol) was added, followed by another after 1 hour (20.0 mL, 20 mmol). After an additional hour,
30 the reaction was quenched with EtOAc at -78 °C, followed by sat. Na-K-tartrate soln., then warmed to room temperature. The solution was poured into EtOAc, washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The resulting material was purified by silica gel chromatography (50% EtOAc/hexane) to give the titled lactol.

- 139 -

Step C: Preparation of (S)-3-(tert-butoxycarbonylamino)-N-(3-chlorophenyl)-4-hydroxy-1-butanamine

To a solution of lactol from Step B (4.49 g, 22.2 mmol) and 3-chloroaniline (2.58 mL, 24.4 mmol) in 50 mL of dichloromethane at room temperature was added acetic acid (1.27 mL, 22.2 mmol). After 10 min, sodium triacetoxyborohydride (6.59 g, 31.1 mmol) was added, and the solution was stirred for 1.5 hours. The reaction was quenched with sat. aq. NaHCO₃, diluted with CH₂Cl₂, and the layers were separated. The organic material was dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide a solid which was purified by silica gel chromatography (EtOAc/hexane) to give the titled amine.

Step D: Preparation of (S)-N-[2-(tert-butoxycarbonylamino)-4-hydroxybutyl]-2-chloro-N-(3-chlorophenyl)acetamide

The aniline derivative from Step C (5.29 g, 16.9 mmol) was dissolved in 60 mL of EtOAc and 60 mL of sat. NaHCO₃ soln., then cooled to 0 °C. With vigorous stirring, chloroacetyl chloride (1.48 mL, 18.5 mmol) was added dropwise. After 2 hours, the reaction was diluted with water and EtOAc, and the organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the titled chloroacetamide, which was used without further purification.

Step E: Preparation of (S)-4-(tert-butoxycarbonyl)-1-(3-chlorophenyl)-5-(2-hydroxyethyl)piperazin-2-one

To a solution of the chloroacetamide from Step D (6.32 g, 16.1 mmol) in 80 mL of DMF at 0 °C was added cesium carbonate (15.8 g, 48.3 mmol). The solution was stirred until tlc analysis indicated consumption of the starting material (ca. 5 hours). The solution was poured into EtOAc, washed with water and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the crude product. This material was purified by silica gel chromatography

- 140 -

(99:1:0-95:5:0.15 CHCl₃:MeOH:NH₄OH) to yield the product containing a minor amount of DMF impurity.

5 Step F: Preparation of (S)-4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-5-[2-(methanesulfonyloxy)ethyl]piperazin-2-one

10 To a solution of the alcohol from Step E (3.58 g, 10.1 mmol) in 50 mL of dichloromethane at 0 °C was added diisopropylethylamine (3.5 mL, 20.2 mmol), followed by methanesulfonyl chloride (0.936 mL, 12.1 mmol). The solution was stirred for 45 minutes, then quenched with 10% citric acid. The solution was washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the crude product which was used in the next step without further purification.

15 Step G: Preparation of (S)-4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-5-[2-(ethylthio)ethyl]piperazin-2-one

20 To a solution of the mesylate from Step F (3.6 g, 8.3 mmol) in 100 mL of DMF at 0 °C was added sodium ethanethiolate (1.4 g, 16.6 mmol). After 2 hours, the reaction was poured into EtOAc, washed with sat. NaHCO₃ and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the crude product which was used in the next step without further purification.

25 Step H: Preparation of (S)-4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-5-[2-(ethanesulfonyl)ethyl]piperazin-2-one

30 To solution of the product from Step G (3.12 g, 7.82 mmol) in methanol (50 mL) was added a solution of magnesium monoperoxyphthalate (11.6 g, 23.5 mmol) in 50 mL MeOH at room temperature. After 45 minutes, the reaction was quenched by the addition of 2N Na₂S₂O₃ soln. The solution was poured into EtOAc and sat NaHCO₃ solution, and the organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the

- 141 -

crude sulfone. This material was chromatographed on silica gel with 2% methanol in chloroform to obtain the titled compound.

Step I: Preparation of (S)-1-(3-chlorophenyl)-5-[2-(ethanesulfonyl)ethyl]piperazin-2-one

5 Through a solution of the Boc-protected piperazinone from Step H (1.75 g, 4.06 mmol) in 20 mL of EtOAc at 0 °C was bubbled anhydrous HCl gas. The saturated solution was stirred for 30 minutes, then concentrated *in vacuo* to provide the hydrochloride of title
10 compound as a white powder. This material was suspended in EtOAc and treated with dilute aqueous NaHCO₃ solution. The aqueous phase was extracted with EtOAc, and the combined organic mixture was washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The resulting titled amine was reconcentrated from toluene to provide
15 the title compound suitable for use in the next step.

Step J: Preparation of (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)ethyl]-2-piperazinone dihydrochloride

20 To a solution of the amine from Step I (480 mg, 1.45 mmol), imidazole carboxaldehyde from Step E of Example 42 (460 mg, 2.2 mmol), and acetic acid (0.415 mL, 7.25 mmol) in 10 mL of 1,2-dichloroethane was added sodium triacetoxyborohydride (615 mg, 2.9 mmol). The reaction was stirred for 18 hours, then quenched at 0 °C
25 with sat. NaHCO₃ solution. The solution was poured into CH₂Cl₂, and the organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The resulting product was purified by silica gel chromatography (2-5% MeOH:CHCl₃), to give the desired product and less polar boron complex. The latter compound was taken up in
30 dichloromethane (1 mL) and benzene (5 mL), treated with n-propylamine (1 mL) for 18 hours, and concentrated *in vacuo*. The residue was purified by silica gel chromatography (2-5% MeOH:CHCl₃), combined with the former batch, taken up in EtOAc/methanol, and treated with 2.1 equivalents of 1 M HCl/ether

- 142 -

solution. After concentrated *in vacuo*, the product dihydrochloride was isolated as a white powder.

EXAMPLE 44

5

Preparation of (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)methyl]-2-piperazinone dihydrochloride

- 10 Step A: Preparation of (S)-2-(*tert*-butoxycarbonylamino)-*N*-(3-chlorophenyl)-3-[(triphenylmethyl)thio]-1-propanamine
- To a solution of 3-chloroaniline (0.709 mL, 6.70 mmol) in 30 mL of dichloromethane at room temperature was added 1.2 g of crushed 4Å molecular sieves. Sodium triacetoxyborohydride (3.55 g, 16.7 mmol) was added, followed by dropwise addition of *N*-methylmorpholine to achieve a pH of 6.5. L-S-Trityl-*N*-Boc-cysteinal (3.15 g, 7.04 mmol) (prepared according to S.L. Graham et al. *J. Med. Chem.*, (1994) Vol. 37, 725-732) was added, and the solution was stirred for 48 hours. The reaction was quenched with sat. aq. NaHCO₃, diluted with EtOAc, and the layers were separated. The organic material was washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide an oil which was purified by silica gel chromatography (15% EtOAc/hexane) to give the title amine.

- 25 Step B: Preparation of (S)-*N*-[2-(*tert*-butoxycarbonylamino)-3-((triphenylmethyl)thio)propyl]-2-chloro-*N*-(3-chlorophenyl)acetamide
-

- The aniline derivative from Step A (2.77 g, 4.95 mmol) was dissolved in 73 mL of EtOAc and 73 mL of sat. NaHCO₃ soln., then cooled to 0 °C. With vigorous stirring, chloroacetyl chloride (0.533 mL, 6.69 mmol) was added dropwise. After 3 hours, the reaction was diluted with water and EtOAc, and the organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo*

- 143 -

to provide crude titled chloroacetamide which was used without further purification.

5 Step C: Preparation of (S)-4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-5-[S-(triphenylmethyl)thiomethyl]piperazin-2-one

10 To a solution of chloroacetamide from Step B (3.29 g crude, theoretically 4.95 mmol) in 53 mL of DMF at 0 °C was added cesium carbonate (4.84 g, 14.85 mmol). The solution was stirred for 48 hours, allowing it to warm to room temperature. The solution was poured into EtOAc, washed with water and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the crude product as an oil. This material was purified by silica gel chromatography (20% EtOAc/hexane) to yield the product as a white solid.

15 Step D: Preparation of (S)-4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-5-(thiomethyl)piperazin-2-one

20 A solution of piperazinone from Step C (625 mg, 1.04 mmol) in degassed EtOAc (38 mL) and EtOH (12 mL) was warmed to 30 °C. A solution of AgNO₃ (177 mg, 1.04 mmol) and pyridine (0.084 mL, 1.04 mmol) in 8 mL of EtOH was added, and the solution was heated to reflux. After 45 minutes, the reaction was concentrated *in vacuo*, then redissolved in 26 mL of degassed EtOAc. Through this solution was bubbled H₂S gas for 2.5 minutes, then activated charcoal was added after 4 minutes. The material was filtered through celite and rinsed with degassed EtOAc, concentrated *in vacuo*, then reconcentrated from degassed CH₂Cl₂ to provide the crude product which was used without further purification.

30 Step E: Preparation of (S)-4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-5-[(ethylthio)methyl]piperazin-2-one

A solution of the thiol from Step D (*ca.* 1.04 mmol) in 3 mL of THF was added *via* cannula to a suspension of NaH (51.4 mg, 60% disp. in mineral oil, 1.28 mmol) in 2 mL THF at 0 °C. After 10

- 144 -

minutes, iodoethane was added (0.079 mL, 0.988 mmol), and the solution was stirred for 1.5 hours. The reaction was poured into EtOAc, washed with sat. NaHCO₃ and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the crude product. This material was purified by silica gel chromatography (1% MeOH/CH₂Cl₂) to yield the titled product.

Step F: Preparation of (S)-4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-5-[(ethanesulfonyl)methyl]piperazin-2-one

To a solution of the sulfide from Step E (217 mg, 0.563 mmol) in 3 mL of MeOH at 0 °C was added a solution of magnesium monoperoxyphthalate (835 mg, 1.69 mmol) in 2 mL MeOH. The reaction was stirred overnight, allowing it to warm to room temperature. The solution was cooled to 0 °C, quenched by the addition of 4 mL 2N Na₂S₂O₃ soln., then concentrated *in vacuo*. The residue was partitioned between EtOAc and sat NaHCO₃ solution, and the organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the crude sulfone as a white waxy solid.

Step G: Preparation of (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)methyl]-2-piperazinone dihydrochloride

To a solution of the Boc-protected piperazinone from Step F (224 mg, 0.538 mmol) in 5 mL of dichloromethane at 0 °C was added 2.5 mL of trifluoroacetic acid (TFA). After 45 minutes, the reaction was concentrated *in vacuo*, then azeotroped with benzene to remove the excess TFA. The residue was taken up in 4 mL of 1,2-dichloroethane and cooled to 0 °C. To this solution was added 4 Å powdered molecular sieves (340 mg), followed by sodium triacetoxymethylborohydride (285 mg, 1.34 mmol) and several drops of triethylamine to achieve pH = 6. The imidazole carboxaldehyde from Step E of Example 42 (125 mg, 0.592 mmol) was added, and the reaction was stirred at 0 °C. After 2 days, the reaction was poured into EtOAc, washed with dilute aq. NaHCO₃,

- 145 -

- and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The crude product was taken up in methanol and injected onto a preparative HPLC column and purified with a mixed gradient of 15%-50% acetonitrile/0.1% TFA; 85%-50% 0.1% aqueous TFA over 60 minutes.
- 5 After concentration *in vacuo*, the resultant product was partitioned between dichloromethane and aq. NaHCO₃ soln., and the aqueous phase was extracted with CH₂Cl₂. The organic solution was washed with brine, dried (Na₂SO₄), filtered, and concentrated to dryness to provide the product free base, which was taken up in CH₂Cl₂ and treated with
- 10 2.1 equivalents of 1 M HCl/ether solution. After concentrated *in vacuo*, the product dihydrochloride was isolated as a white powder.

EXAMPLE 45

- 15 Preparation of (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[N-ethyl-2-acetamido]-2-piperazinone dihydrochloride
-

- 20 Step A: Preparation of (S)-4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-5-[2-(oxo)ethyl]piperazin-2-one
- To a solution of oxalyl chloride (0.608 mL, 6.97 mmol) in dichloromethane (40 mL) at -78 °C was added DMSO (0.990 mL, 13.9 mmol) over 2-3 minutes. The solution was stirred for 10 minutes, then a solution of the alcohol from Step E of Example 43 (2.06 g, 5.81
- 25 mmol) in 10 mL of dichloromethane was added over 5 minutes, keeping the reaction temperature below -70 °C. The reaction was stirred for 10 minutes, then triethylamine (2.43 mL) was added, and the reaction was stirred at -78 °C for 5 minutes, then allowed to warm to room temperature. After 45 minutes, the solution was poured into
- 30 dichloromethane, and washed with sat. NH₄Cl soln., 10% citric acid soln., water, and brine. The solution was dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the titled aldehyde.

- 146 -

Step B: Preparation of (S)-4-(*tert*-butoxycarbonyl)-5-
(carboxymethyl)-1-(3-chlorophenyl)piperazin-2-one

To a solution of the aldehyde from Step A (1.52 g, 4.31 mmol) in 2-methyl-2-propanol (50 mL) and 2-methyl-2-butene (10 mL) at room temperature was added a solution of sodium chlorite (585 mg, 5.17 mmol) and sodium dihydrogenphosphate (595 mg, 4.31 mmol) in 10 mL of water. The reaction turned yellow, then slowly faded to light pink over 45 minutes. The solution was poured into EtOAc, and washed with sat. sodium bisulfite soln. The aqueous layer was acidified to pH 3 with 2.75 M KHSO₄ soln., and extracted several times with EtOAc. The combined organic extracts were dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the titled carboxylic acid.

Step C: Preparation of (S)-4-(*tert*-butoxycarbonyl)-5-[*N*-ethyl-2-
acetamidol-1-(3-chlorophenyl)piperazin-2-one

The product from Step B (200 mg, 0.56 mmol), ethylamine hydrochloride (114 mg, 1.4 mmol), EDC hydrochloride (140 mg, 0.73 mmol) and HOBt hydrate (113 mg, 0.84 mmol) were stirred in dry DMF (3 mL) at 0°C under nitrogen. After one hour, the solution was warmed to room temperature, and stirred overnight. The DMF was removed *in vacuo*, and the residue was partitioned between ethyl acetate and water. The organic phase was washed with 10% citric acid, saturated sodium bicarbonate, water, and brine, and dried with sodium sulfate. The solvent was removed *in vacuo* to give the title compound.

Step D: Preparation of (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[*N*-ethyl-2-acetamido]-
2-piperazinone dihydrochloride

The titled product was prepared from the product of Step C (184 mg, 0.47 mmol) in analogy to Example 42, Steps K and L. The product was isolated as a white solid.

- 147 -

EXAMPLE 46

Preparation of (\pm) -5-(2-butynyl)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone dihydrochloride

5

Step A: Preparation of 1-(methanesulfonyl)-2-butyne

To a solution of 2-butyne (10.0 mL, 134 mmol) in 200 mL of dichloromethane at 0 °C was added methanesulfonyl chloride (23.4 g, 134 mmol), followed by dropwise addition of diisopropylethylamine (30 mL, 174 mmol). After 1.5 hours, the solution was poured into 0.5 N KHSO₄ soln, and the organic layer was washed with brine. The solution was dried over Na₂SO₄, filtered, and concentrated *in vacuo* to provide the titled product.

15 Step B: Preparation of (\pm) -ethyl 2-[(phenylmethyl)imino]-4-hexynoate

To a solution of glycine ethyl ester hydrochloride (10.11 g, 72.4 mmol) in 200 mL of dichloromethane was added benzaldehyde (7.36 mL, 72.4 mmol), triethylamine (20.0 mL, 143 mmol), and magnesium sulfate (6g). The solution was stirred at room temperature for 16 hours, filtered through a glass frit, and concentrated *in vacuo*. The residue was partitioned between ether and water, and the organic layer was washed with brine. The solution was dried over Na₂SO₄, filtered, and concentrated *in vacuo* to provide a pale yellow oil. A portion of this oil (9.90 g, 51.8 mmol) was dissolved in 200 mL of THF and cooled to -78 °C under nitrogen atmosphere. A solution of potassium *tert*-butoxide in THF (51.8 mL of 1 M, 51.8 mmol) was added dropwise to produce a bright red solution. After 20 minutes, a solution of the mesylate from Step A (8.05 g, 54.4 mmol) in 20 mL of THF was added dropwise via cannula, and the solution was allowed to warm to room temperature. After 2 hours, the reaction was poured into EtOAc and washed with sat. NaHCO₃ soln. and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the titled product.

- 148 -

Step C: Preparation of (\pm) -ethyl 2-[(*tert*-butoxycarbonyl)amino]-4-hexynoate

A solution of the product from Step B (ca. 51.8 mmol) was stirred at room temperature in 5% aqueous HCl solution (100 mL). After 12 hours, the solution was concentrated *in vacuo* to give an orange oil. This product was taken up in 50 mL of THF and sat. NaHCO₃ soln. was added (50 mL), followed by di-*tert*-butylpyrocarbonate (11.3 g, 51.8 mmol) at room temperature. After 6 hours, the reaction was poured into EtOAc and washed with sat. NaHCO₃ soln. and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the titled product.

Step D: Preparation of (\pm) -2-[(*tert*-butoxycarbonyl)amino]-4-hexynoic acid

To a solution of the product from Step C (ca. 51.8 mmol) in THF (100 mL) and water (20 mL) was added at 0 °C a solution of lithium hydroxide monohydrate (6.5 g, 155 mmol). The solution was stirred for 1 hour at 0 °C, then warmed to room temperature. After 48 hours, the solution was concentrated *in vacuo*. The aqueous mixture was extracted with EtOAc, acidified at 0 °C with 10% aq. HCl soln., then extracted with three portions of dichloromethane. The combined dichloromethane extracts were dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the titled product as an orange oil.

Step E: Preparation of (\pm) -2-(*tert*-butoxycarbonylamino)-*N*-methoxy-*N*-methyl-4-hexynamide

The product from Step D (10.58 g, 46.6 mmol), *N,O*-dimethylhydroxylamine hydrochloride (9.09 g, 93.2 mmol), HOBT hydrate (9.44 g, 69.9 mmol) and triethylamine (13.0 mL, 93.2 mmol) were stirred in dry DMF (150 mL) at 0 °C under nitrogen. EDC hydrochloride (11.5 g, 60.6 mmol) was added, and the reaction was stirred for 3 hours. The solution was partitioned between 2:1 ethyl acetate:hexane and water, washed with water, 10% aq. HCl, sat.

- 149 -

NaHCO₃ soln and brine, then dried with sodium sulfate. The solvent was removed *in vacuo* to give the title compound as an orange oil

Step F: Preparation of (±)-2-(tert-butoxycarbonylamino)-4-hexynal

5 A suspension of lithium aluminum hydride (1.56 g, 41.1 mmol) in ether (150 mL) was stirred at room temperature for 30 minutes. The solution was cooled to -55 °C under nitrogen, and a solution of the product from Step E (11.10 g, 41.1 mmol) in ether (150 mL) was added over 15 min, maintaining the temperature below -50 °C.
10 When the addition was complete, the reaction was warmed to 5 °C, then recooled to -40 °C. A solution of potassium hydrogen sulfate (21.8 g) in 25 mL water was slowly added, maintaining the temperature below -35 °C. The mixture was warmed to room temperature and stirred for one hour, filtered through Celite, and concentrated *in vacuo* to provide
15 the title aldehyde.

Step G: Preparation of (±)-2-(tert-butoxycarbonylamino)-N-(3-chlorophenyl)-4-butyamine

20 To a 0 °C solution of 3-chloroaniline (4.33 mL, 40.9 mmol), the product from Step F (ca. 41 mmol), and crushed 4 Å molecular sieves (10 g) in dichloroethane (100 mL) under nitrogen was added sodium triacetoxyborohydride (12.9 g, 61.5 mmol). The reaction was stirred for one hour, then warmed to room temperature. After 3 hours, the solution was poured into EtOAc and washed with water, sat.
25 NaHCO₃ soln. and brine. The solution was dried over sodium sulfate and concentrated *in vacuo* to provide the crude product.

Step H: Preparation of (±)-N-[2-(tert-butoxycarbonylamino)-4-hexynyl]-2-chloro-N-(3-chlorophenyl)acetamide

30 A solution of the product from Step G (1.68 g, 5.22 mmol) and triethylamine (1.20 mL, 8.61 mmol) in 15 mL of CH₂Cl₂ was cooled to 0 °C. Chloroacetyl chloride (0.457 mL, 5.74 mmol) was added dropwise, and the reaction was maintained at 0 °C with stirring. After 30 minutes, another portion of chloroacetyl chloride (0.20 mL)

- 150 -

and triethylamine (0.5 mL) was added dropwise. After 30 min, the reaction was poured into EtOAc and washed with 10% aq. HCl, sat. aq. NaHCO₃ soln., and brine. The solution was dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide a brown oil. This material was
5 purified by silica gel chromatography (20-35% EtOAc/hexane) to yield the titled product.

Step I: Preparation of (±)-4-(tert-butoxycarbonyl)-5-(2-butynyl)-1-(3-chlorophenyl)-2-piperazinone

10 To a solution of the chloroacetamide from Step H (1.68 g, 4.23 mmol) in 15 mL of dry DMF at 0 °C was added Cs₂CO₃ (3.08 g, 9.48 mmol). The solution was stirred for 30 minutes, then allowed to warm to room temperature. After 14 hours, the reaction was poured into 50% EtOAc/hexane, washed with water and brine, dried (Na₂SO₄),
15 filtered, and concentrated *in vacuo* to provide the crude product. This material was purified by silica gel chromatography (20-40% EtOAc/hexane) to yield the titled product.

20 Step J: Preparation of (±)-5-(2-butynyl)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone dihydrochloride

To a solution of the Boc-protected piperazinone from Step I (1.03 g, 2.85 mmol) in 7 mL of dichloromethane at 0 °C was added 4 mL of trifluoroacetic acid (TFA). After 4 hours, the reaction was
25 warmed to room temperature, and stirred an additional 6 hours. The solution was concentrated *in vacuo*, then azeotroped with benzene to remove the excess TFA. A portion of the residue (255 mg, 0.678 mmol) was taken up in 6 mL of 1,2-dichloroethane and cooled to 0 °C. To this solution was added 4Å powdered molecular sieves (600 mg),
30 followed by sodium triacetoxyborohydride (214 mg, 1.02 mmol). The imidazole carboxaldehyde from Step E of Example 42 (186 mg, 0.881 mmol) was added, and the reaction was stirred at 0 °C. After 24 hours, the reaction was poured into EtOAc, washed with dilute aq. NaHCO₃, and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo*. This

- 151 -

material was purified by silica gel chromatography (2-5% MeOH:CH₂Cl₂) to yield a white foam which was taken up in CH₂Cl₂ and treated with 2.1 equivalents of 1 M HCl/ether solution. After concentrated *in vacuo*, the product dihydrochloride was isolated as a white powder.

EXAMPLE 47

Preparation of 1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)imidazolyl-methyl]-2-piperazinone dihydrochloride

Step A: Preparation of N-(3-chlorophenyl)ethylenediamine hydrochloride

To a solution of 3-chloroaniline (30.0 mL, 284 mmol) in 500 mL of dichloromethane at 0 °C was added dropwise a solution of 4 N HCl in 1,4-dioxane (80 mL, 320 mmol HCl). The solution was warmed to room temperature, then concentrated to dryness *in vacuo* to provide a white powder. A mixture of this powder with 2-oxazolidinone (24.6 g, 282 mmol) was heated under nitrogen atmosphere at 160 °C for 10 hours, during which the solids melted, and gas evolution was observed. The reaction was allowed to cool, forming the crude diamine hydrochloride salt as a pale brown solid.

Step B: Preparation of N-(tert-butoxycarbonyl)-N'-(3-chlorophenyl)ethylenediamine

The amine hydrochloride from Step A (*ca.* 282 mmol, crude material prepared above) was taken up in 500 mL of THF and 500 mL of sat. aq. NaHCO₃ soln., cooled to 0 °C, and di-*tert*-butylpyrocarbonate (61.6 g, 282 mmol) was added. After 30 h, the reaction was poured into EtOAc, washed with water and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the titled carbamate as a brown oil which was used in the next step without further purification.

- 152 -

Step C: Preparation of *N*-[2-(*tert*-butoxycarbonyl)ethyl]-*N*-(3-chlorophenyl)-2-chloroacetamide

A solution of the product from Step B (77 g, *ca.* 282 mmol) and triethylamine (67 mL, 480 mmol) in 500 mL of CH₂Cl₂ was cooled to 0 °C. Chloroacetyl chloride (25.5 mL, 320 mmol) was added dropwise, and the reaction was maintained at 0 °C with stirring. After 3 h, another portion of chloroacetyl chloride (3.0 mL) was added dropwise. After 30 min, the reaction was poured into EtOAc (2 L) and washed with water, sat. aq. NH₄Cl soln, sat. aq. NaHCO₃ soln., and brine. The solution was dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the chloroacetamide as a brown oil which was used in the next step without further purification.

Step D: Preparation of 4-(*tert*-butoxycarbonyl)-1-(3-chlorophenyl)-2-piperazinone

To a solution of the chloroacetamide from Step C (*ca.* 282 mmol) in 700 mL of dry DMF was added K₂CO₃ (88 g, 0.64 mol). The solution was heated in an oil bath at 70-75 °C for 20 hours, cooled to room temperature, and concentrated *in vacuo* to remove *ca.* 500 mL of DMF. The remaining material was poured into 33% EtOAc/hexane, washed with water and brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the product as a brown oil. This material was purified by silica gel chromatography (25-50% EtOAc/hexane) to yield pure product, along with a sample of product (*ca.* 65% pure by HPLC) containing a less polar impurity.

Step E: Preparation of 1-(3-chlorophenyl)-2-piperazinone

Through a solution of Boc-protected piperazinone from Step D (17.19 g, 55.4 mmol) in 500 mL of EtOAc at -78 °C was bubbled anhydrous HCl gas. The saturated solution was warmed to 0 °C, and stirred for 12 hours. Nitrogen gas was bubbled through the reaction to remove excess HCl, and the mixture was warmed to room temperature. The solution was concentrated *in vacuo* to provide the hydrochloride as a white powder. This material was taken up in 300

- 153 -

mL of CH₂Cl₂ and treated with dilute aqueous NaHCO₃ solution. The aqueous phase was extracted with CH₂Cl₂ (8 x 300 mL) until tlc analysis indicated complete extraction. The combined organic mixture was dried (Na₂SO₄), filtered, and concentrated *in vacuo* to provide the
5 titled free amine as a pale brown oil.

Step F: Preparation of 1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone dihydrochloride

To a solution of the amine from Step E (55.4 mmol,
10 prepared above) in 200 mL of 1,2-dichloroethane at 0 °C was added 4Å powdered molecular sieves (10 g), followed by sodium triacetoxymethylborohydride (17.7 g, 83.3 mmol). The imidazole carboxaldehyde from Step E of Example 42 (11.9 g, 56.4 mmol) was added, and the reaction was stirred at 0 °C. After 26 hours, the reaction
15 was poured into EtOAc, washed with dilute aq. NaHCO₃, and the aqueous layer was back-extracted with EtOAc. The combined organics were washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The resulting product was taken up in 500 mL of 5:1 benzene:CH₂Cl₂, and propylamine (20 mL) was added. The mixture
20 was stirred for 12 hours, then concentrated *in vacuo* to afford a pale yellow foam. This material was purified by silica gel chromatography (2-7% MeOH/CH₂Cl₂), and the resultant white foam was taken up in CH₂Cl₂ and treated with 2.1 equivalents of 1 M HCl/ether solution. After concentrated *in vacuo*, the product dihydrochloride was isolated
25 as a white powder.

EXAMPLE 48

5(S)-Butyl-4-[1-(4-cyanobenzyl)-2-methyl-5-imidazolylmethyl]-1-(2,3-dimethylphenyl)-piperazin-2-one dihydrochloride
30

- 154 -

Step A: 4-Bromo-2-methylimidazole-5-carboxaldehyde

4-Bromo-5-hydroxymethyl-2-methylimidazole was prepared according to the procedure described by S. P. Watson, Synthetic Communications, 22, 2971-2977 (1992). A solution of 4-bromo-5-hydroxymethyl-2-methylimidazole (4.18 g, 21.9 mmol) was refluxed with manganese dioxide (16.1 g) in 1:1 methylene chloride:dioxane (200 mL) for 16 h. The cooled reaction was filtered through celite and concentrated to yield the title compound as a pale yellow solid. ¹H NMR (CDCl₃, 300 MHz) δ 9.57 (1H, s), 2.52 (3H, s).

10

Step B: 4-Bromo-1-(4-cyanobenzyl)-2-methylimidazole-5-carboxaldehyde

4-Cyanobenzylbromide (1.05 g, 5.39 mmol) was added to a solution of 4-bromo-2-methylimidazole-5-carboxaldehyde (1.02 g, 5.39 mmol) in dimethylacetamide (15 mL). The solution was cooled to -10°C and powdered potassium carbonate (0.745 g, 5.39 mmol) added. The reaction was stirred at -10°C for 2 h, and a further 4 h at 20°C. The reaction was diluted with water and extracted with ethyl acetate. The organic phase was washed with water, saturated brine, and dried over magnesium sulfate. Solvent evaporation yielded a white solid. ¹H NMR (CDCl₃, 400 MHz) δ 9.68 (1H, s), 7.64 (2H, d, J=7 Hz), 7.15 (2H, d, J=7Hz) 5.59 (2H, s), 2.40 (3H, s).

15

20

Step C: 1-(4-Cyanobenzyl)-2-methylimidazole-5-carboxaldehyde

A solution of 4-bromo-1-(4-cyanobenzyl)-2-methylimidazole-5-carboxaldehyde (1.33 g, 4.37 mmol) and imidazole (0.600 g, 8.74 mmol) in 1:1 ethyl acetate-alcohol (150 mL) was stirred with 10% palladium on carbon (0.020 g) under 1 atm hydrogen. After 2 h, the reaction was filtered through celite and concentrated to give the title compound as a white solid. ¹H NMR (DMSO-d₆, 400 MHz) δ 9.62 (1 H, s), 7.90 (1H, s), 7.81 (2H, d, J=8 Hz), 7.20 (2H, d, J=8 Hz), 5.64 (2H, s), 2.33 (3H, s).

25

30

- 155 -

Step D: 5(S)-Butyl-4-[1-(4-cyanobenzyl)-2-methyl-5-imidazolylmethyl]-1-(2,3-dimethylphenyl)piperazin-2-one dihydrochloride

Sodium triacetoxyborohydride (0.265 g, 1.25 mmol) was added to a solution of 1-(4-cyanobenzyl)-2-methylimidazole-5-carboxaldehyde (0.190 g, 0.843 mmol), 5(S)-butyl-1-(2,3-dimethylphenyl)piperazin-2-one hydrochloride (0.250 g, 0.843 mmol), N-methylmorpholine (0.093 mL, 0.843 mmol) in dichloroethane (10 mL), according to the procedure described in Example 39, Step E. The title compound was purified by preparative HPLC using a gradient of 70 to 45% Solvent A. The free base was isolated and converted to dihydrochloride salt. The title compound was obtained as a white solid. FAB ms: 470 (M+1). Anal. Calc for $C_{29}H_{35}N_5O \cdot 2 HCl \cdot 1.45 H_2O$, C, 61.25; H, 7.07; N, 12.32. Found: C, 61.56; H, 6.99; N, 11.32.

EXAMPLE 49

4-[1-(2-(4-Cyanophenyl)-2-propyl)-5-imidazolylmethyl]-1-(3-chlorophenyl)-5(S)-(2-methylsulfonyl)ethyl)piperazin-2-one dihydrochloride

Step A: 4-(tert-Butyldimethylsilyloxymethyl)-1-triphenylmethylimidazole

tert-Butyldimethylsilylchloride (2.83 g, 18.76 mmol) was added to a suspension of 4-hydroxymethyl-1-triphenylmethylimidazole (5.80 g, 17.05 mmol) in DMF (200 mL) containing imidazole (3.48 g, 51.1 mmol). After 15 min, a clear colorless solution was obtained which was stirred at room temperature. When reaction was complete, the DMF was removed in vacuo and the residue partitioned between ethyl acetate and water. The organic phase was washed with water, saturated brine, and dried over magnesium sulfate. The title compound was obtained as a clear gum.

- 156 -

Step B: 5-*tert*-Butyldimethylsilyloxymethyl-1-(4-cyanobenzyl)imidazole

A solution of 4-(*tert*-butyldimethylsilyloxymethyl)-1-triphenylmethylimidazole (4.66 g, 10.26 mmol) and 1-bromomethyl-4-cyanobenzene (2.01 g, 10.26 mmol) in acetonitrile (50 mL) was
5 refluxed 4 h. The reaction was cooled, acetonitrile removed in vacuo, and the residue dissolved in methanol (30 mL). This solution was refluxed for 2h, cooled and methanol evaporated. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate. The
10 crude product was chromatographed on silica with 3% methanol in chloroform. The title compound was obtained as a white solid.

Step C: 5-*tert*-Butoxydimethylsilyloxymethyl-1-[2-(4-cyanophenyl)-2-propyl]imidazole

A solution of 5-*tert*-butyldimethylsilyloxymethyl-1-(4-cyanobenzyl)imidazole (1.005 g, 3.07 mmol) in THF (25 mL) under
15 nitrogen was cooled to -78°C. A solution of lithium hexamethyldisilazide (4.61 mL, 1 M in THF) was added and the reaction stirred at -78°C for 1 h, then warmed to -60°C over 30 min. The
20 reaction was cooled to -78°C, methyl iodide added (0.287 mL, 4.61 mmol), and stirring continued at -78°C for an additional 2 h, before warming to 0°C over 2 h. After 30 min, the reaction was cooled to -78°C, and lithium hexamethyldisilazide (4.61 mL, 1 M in THF) added.
25 After 1 h, methyl iodide was added (0.287 mL, 4.61 mmol) and the reaction allowed to warm to room temperature overnight. The reaction was quenched with water, extracted with ethyl acetate, and the organic phase washed with saturated brine. The crude product was chromatographed on silica gel with 6:4 ethyl acetate: methylene
30 chloride. The title compound was obtained as a golden oil. ¹H NMR (CDCl₃, 400 MHz) δ 7.78 (1H, s), 7.61 (2H, d, J=8Hz), 7.15 (2H, d, J=8Hz), 7.02, (1H, s), 4.00 (2H, s), 1.99 (6H, s), 0.79 (9H, s), -0.74 (6H, s).

- 157 -

Step D: 1-[2-(4-Cyanophenyl)-2-propyl]-5-hydroxymethyl-
imidazole

Tetra-N-butylammonium fluoride (2.99 mL, 1M in THF) was added to a solution of 5-*tert*-butoxydimethylsilyloxymethyl-1-[2-(4-cyanophenyl)-2-propyl]imidazole (0.750 g, 2.72 mmol) in THF (10 mL). After 2 h at room temperature, the reaction was poured into ethyl acetate and extracted with saturated sodium bicarbonate solution. The organic phase was extracted with saturated brine, and dried over magnesium sulfate. The crude product was chromatographed on silica gel with 3% methanol in ethyl acetate. The title compound was obtained as a semi-solid. ¹H NMR (CDCl₃, 400 MHz) δ 7.77 (1H, s), 7.62 (2H, d, J=8Hz), 7.19 (2H, d, J=8Hz), 7.02, (1H, s), 4.01 (2H, s), 2.57 (1H, br s), 2.01 (6H, s).

Step E: 1-[2-(4-Cyanophenyl)-2-propyl]imidazole-5-
carboxaldehyde

A solution of 1-[2-(4-cyanophenyl)-2-propyl]-5-hydroxymethylimidazole (0.450 g, 1.87 mmol) was refluxed in dioxane (20 mL) with manganese dioxide (1.62 g, 18.7 mmol) for 5h. The reaction was cooled, filtered through celite and concentrated. The crude product was purified by chromatography on silica gel; the title compound was isolated as a semi-solid. ¹H NMR (CDCl₃, 400 MHz) δ 9.37 (1H, s), 8.11 (1H, s), 7.92 (1H, s), 7.58 (2H, d, J=8Hz), 7.15 (2H, d, J=8Hz), 2.01 (6H, s).

Step F: 4-[1-(2-(4-Cyanophenyl)-2-propyl)-5-imidazolylmethyl]-1-(3-chlorophenyl)-5(S)-(2-methylsulfonyl)ethyl)piperazin-2-one dihydrochloride

The title compound was prepared according to the procedure described in Example 39, Step E, except using 1-(3-chlorophenyl)-5(S)-(2-methylsulfonyl)ethyl)piperazin-2-one hydrochloride (0.297 g, 0.84 mmol), 1-[2-(4-cyanophenyl)-2-propyl]imidazole-5-carboxaldehyde (0.200 g, 0.84 mmol), N-

- 158 -

methyldmorpholine (0.092 mL, 0.84 mmol) and sodium triacetoxymorohydride (0.267 g, 1.26 mmol). The crude product was purified by preparative HPLC with an 80 to 58% Solvent A gradient. The pure fractions were combined and converted to the hydrochloride salt, yielding the title compound. FAB ms: 540 (M+1). Anal. Calc for $C_{27}H_{30}ClN_5O_3S \cdot 2 HCl \cdot 3 H_2O$, C, 48.62; H, 5.74; N, 10.50. Found: C, 48.62; H, 5.73; N, 9.89.

EXAMPLE 50

10

5(S)-*n*-Butyl-4-[1-(4-cyanobenzyl)-5-imidazolymethyl]-1-(2-methylphenyl)piperazin-2-one dihydrochloride

15

Step A: N-(2-Methylphenyl)-2(S)-(tert-butoxycarbonylamino)-hexanamine

20

The title compound was prepared according to the procedure described in Example 39, Step C, except using *o*-toluidine (0.32 mL, 3.00 mmol), 2(S)-(tert-butoxycarbonylamino)hexanal (0.538, 2.50 mmol), sodium triacetoxymorohydride (0.795 g, 3.75 mmol) in dichloroethane (10 mL). The crude product was purified by column chromatography to yield the title compound.

25

Step B: 4-tert-Butoxycarbonyl-5(S)-*n*-butyl-1-(2-methylphenyl)piperazin-2-one

30

The title compound was prepared essentially according to the procedure described in Example 39, Step D, except using N-(2-methylphenyl)-2(S)-(tert-butoxycarbonylamino)hexanamine (0.506 g, 1.65 mmol), chloroacetyl chloride (0.158 mL, 1.98 mmol) in ethyl acetate-saturated sodium bicarbonate at 0°C. The crude product thus obtained was dissolved in DMF (15 mL), cooled to 0°C under nitrogen, and treated with cesium carbonate (1.61 g, 4.95 mmol). The reaction was stirred at 0°C for 2h, and at room temperature for 2h. The reaction was quenched with saturated ammonium chloride, and

extracted with ethyl acetate. The extracts were dried and evaporated to give the title compound.

5 Step C: 5(S)-*n*-Butyl-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-1-(2-methylphenyl)piperazin-2-one dihydrochloride

The product from Step B (0.534 g, 1.50 mmol) was deprotected with trifluoroacetic acid (4 mL) in methylene chloride (10 mL). The title compound was prepared according to the procedure described in Example 39, Step E, except using 5(S)-*n*-butyl-1-(2-methylphenyl)piperazin-2-one ditrifluoroacetic acid salt, 1-(4-cyanobenzylimidazole)-5-carboxaldehyde (0.317 g, 1.50 mmol), and sodium triacetoxyborohydride (0.477 g, 2.25 mmol) in dichloroethane (15 mL). The crude product was injected onto a preparative HPLC column and purified with a mixed gradient of acetonitrile/0.1% TFA and 0.1% aqueous TFA. The pure fractions were combined and converted to the HCl salt. The title compound was obtained as a white solid. FAB ms ($m+1$) 442. Anal. Calc. for $C_{27}H_{31}N_5O \cdot 2.5 HCl \cdot 2.05 H_2O$: C, 56.95; H, 6.66; N, 12.30. Found: C, 56.93; H, 5.75; N, 11.55.

EXAMPLE 51

25 4-[1-(4-Cyanobenzyl)-5-imidazolylmethyl]-5(S)-(2-fluoroethyl)-1-(3-
chlorophenyl)piperazin-2-one dihydrochloride

Step A: 4-*tert*-Butoxycarbonyl-1-(3-chlorophenyl)-5(S)-(2-fluoroethyl)-piperazin-2-one

30 4-*tert*-Butoxycarbonyl-1-(3-chlorophenyl)-5(S)-(2-methylsulfonyloxyethyl)-piperazin-2-one (0.433 g, 1.00 mmol) and *tert*-butylammonium fluoride (3.0 mL, 1M in THF) were stirred at room temperature in acetonitrile (5 mL) for 72 h. The reaction was quenched with saturated sodium bicarbonate and extracted with ethyl

- 160 -

acetate. The organic extracts were dried, concentrated and purified by column chromatography using 20% ethyl acetate in hexane. The title compound was obtained as a thick oil.

5

Step B: 4-[1-(4-Cyanobenzyl)-5-imidazolylmethyl]-5(S)-(2-fluoroethyl)-1-(3-chlorophenyl)piperazin-2-one dihydrochloride

4-*tert*-Butoxycarbonyl-1-(3-chlorophenyl)-5(S)-(2-fluoroethyl)-piperazin-2-one (0.191 g, 0.54 mmol) was deprotected with trifluoroacetic acid (4 mL) in methylene chloride (10 mL). The title compound was prepared according to the procedure described in Example 39, Step E, except using 1-(3-chlorophenyl)-5(S)-(2-fluoroethyl)piperazin-2-one ditrifluoroacetic acid salt, 1-(4-cyanobenzylimidazole)-5-carboxaldehyde (0.114 g, 0.54 mmol), and sodium triacetoxymethylborohydride (0.172 g, 2.25 mmol) in dichloroethane (5 mL). The crude product was injected onto a preparative HPLC column and purified with a mixed gradient of acetonitrile/0.1% TFA and 0.1% aqueous TFA. The pure fractions were combined and converted to the HCl salt. The title compound was obtained as a white solid. FAB ms (m+1) 452. Anal. Calc. for C₂₄H₂₃ClFN₅O · 2. HCl · 1.7 H₂O: C, 51.90; H, 5.15; N, 12.61. Found: C, 52.22; H, 5.10; N, 12.22.

25

EXAMPLE 52

4-[3-(4-Cyanobenzyl)pyridin-4-yl]-1-(3-chlorophenyl)-5(S)-(2-methylsulfonyl)ethyl)-piperazin-2-one

30 Step A: 3-(4-Cyanobenzyl)pyridin-4-carboxylic acid methyl ester
A solution of 4-cyanobenzyl bromide (625 mg, 3.27 mmol) in dry THF (4mL) was added slowly over ~3 min. to a suspension of activated Zn (dust; 250 mg) in dry THF (2 mL) at 0° under an argon atmosphere.

- 161 -

The ice-bath was removed and the slurry was stirred at room temperature for a further 30 min. Then 3-bromopyridin-4-carboxylic acid methyl ester (540 mg, 2.5 mmol) followed by dichlorobis(triphenylphosphine)nickel (II) (50 mg). The resultant reddish-brown mixture was stirred for 3h at ~40-45°C. The mixture was cooled and distributed between EtOAc (100 ml) and 5% aqueous citric acid (50 mL). The organic layer was washed with H₂O (2X50 mL), dried with Na₂SO₄. After evaporation of the solvent the residue was purified on silica gel, eluting with 35% EtOAc in hexane to give 420 mg as a clear gum. FAB ms (M+1) 253.

Step B: 3-(4-Cyanobenzyl)-4-(hydroxymethyl)pyridine

The title compound was obtained by sodium borohydride (300 mg) reduction of the ester from Step A (415 mg) in methanol (5 mL) at room temperature. After stirring for 4 h the solution was evaporated and the product was purified on silica gel, eluting with 2% methanol in chloroform to give the title compound. FAB ms (M+1) 225.

Step C: 3-(4-Cyanobenzyl)-4-pyridinal

The title compound was obtained by activated manganese dioxide (1.0g) oxidation of the alcohol from Step B (240 mg, 1.07 mmol) in dioxane (10 mL) at reflux for 30 min. Filtration and evaporation of the solvent provided title compound, mp 80-83°C.

Step D: 4-[3-(4-Cyanobenzyl)pyridine-4-yl]-1-(3-chlorophenyl)-5(S)-(2-methylsulfonylethyl)-piperazin-2-one.

This compound was prepared essentially by the same methods as described in Example 39, Step E, except that the imidazolyl carboxaldehyde was replaced by an equal amount of product from Step C (vide supra) and the piperazinone was replaced by an equal amount of 1-(3-chlorophenyl)-5(S)-(2-methylsulfonylethyl)-piperazin-2-one, there was obtained the title compound. FAB ms 523. Anal. Calc. for

- 162 -

$C_{27}H_{27}ClN_4O_3S \cdot 0.15 CHCl_3$: C, 60.28; H, 5.06; N, 10.36 Found: C, 60.37; H, 5.03; N, 10.64.

EXAMPLE 53

5

4-[5-(4-Cyanobenzyl)-1-imidazolylethyl]-1-(3-chlorophenyl)piperazin-2-one.

Step A: 1-Trityl-4-(4-cyanobenzyl)-imidazole.

10

To a suspension of activated zinc dust (3.57g, 54.98mmol) in THF (50ml) was added dibromoethane (0.315ml, 3.60mmol) and the reaction stirred under argon at 20°C. The suspension was cooled to 0°C and α -bromo-p-tolunitrile (9.33g, 47.6mmol) in THF (100ml) was added dropwise over a period of 10 min. The reaction was then allowed to stir at 20°C for 6hr and bis(triphenylphosphine)Nickel II chloride (2.4g, 3.64mmol) and 5-iodotrityl imidazole (15.95g, 36.6mmol) was added in one portion. The resulting mixture was stirred 16hr at 20°C and then quenched by addition of saturated NH_4Cl solution (100ml) and the mixture stirred for 2 hours. Saturated $NaHCO_3$ solution was added to give a pH of 8 and the solution was extracted with EtOAc (2x250ml), dried $MgSO_4$ and the solvent evaporated in vacuo. The residue was chromatographed (SiO_2 , 0-20% EtOAc/ CH_2Cl_2) to afford the title compound as a white solid.

15

20

25

1H NMR δ $CDCl_3$ (7.54 (2H,d, $J=7.9Hz$), 7.38(1H,s), 7.36-7.29 (11H,m), 7.15-7.09(6H,m), 6.58(1H,s), and 3.93(2H,s)ppm.

Step B: 1-Methyl acetyl-5-(4-cyanobenzyl)-imidazole.

To a solution of 1-trityl-4-(4-cyanobenzyl)-imidazole (3.01 g, 6.91 mmol) in acetonitrile (50ml) was added methyl bromoacetate (0.687ml, 7.26mmol) and the mixture heated at 55°C for 16 hours, and then the solvent was evaporated in vacuo. The solids were triturated with EtOAc, collected by filtration and dissolved in methanol (60ml). This suspension was heated at reflux for 20min, cooled and evaporated

30

- 163 -

to dryness. The residue was triturated with EtOAc and the title compound was obtained as a white solid by filtration

¹H NMR δ CDCl₃ (7.61 (2H,d, J=7.9Hz), 7.53(1H,s), 7.27(2H,d,J=7.9Hz), 6.89(1H,s), 4.47(2H,s), 3.98(2H,s) and 3.66(3H,s)ppm.

Step C: 1-Hydroxyethyl-5-(4-cyanobenzyl)-imidazole.

A solution of 1-methyl acetyl-5-(4-cyanobenzyl)-imidazole (0.113 g, 0.472 mmol) in methanol (2ml) at 0°C was treated with sodium borohydride (80.7mg, 2.1mmol). After 1 hour the reaction was quenched by addition of saturated NH₄Cl solution (2ml). Saturated NaHCO₃ was added and the mixture extracted with ethyl acetate (3x25ml), dried MgSO₄ and evaporated in vacuo. The title compound was isolated by chromatography (SiO₂, 10%MeOH in CH₂Cl₂) as a white solid. ¹H NMR δ CDCl₃(7.61 (2H,d, J=7.9Hz), 7.55(1H,s), 7.27(2H,d,J=7.9Hz), 6.83(1H,s), 4.05(2H,s), 3.87(2H,t, J=5.1Hz) and 3.74(2H,t, J=5.1Hz)ppm.

Step D: 2-(-5-(4-Cyanobenzyl)-imidazolyl)ethyl methanesulfonate

A solution of 1-hydroxyethyl-5-(4-cyanobenzyl)-imidazole (0.532 g, 2.34 mmol) in methylene chloride (70ml) at 0°C was treated with Hunigs base (0.489ml, 2.81mmol) and methane sulfonyl chloride (0.219ml, 2.81mmol). After 2 hour the reaction was quenched by addition of saturated NaHCO₃ solution (50ml) and the mixture extracted with methylene chloride (50ml), dried MgSO₄ and the solvent evaporated in vacuo. The title compound was used without further purification. ¹H NMR δ CDCl₃(7.62 (2H,d, J=7.9Hz), 7.54(1H,s), 7.29(2H,d,J=7.9Hz), 6.87(1H,s), 4.25(2H,t), 4.10-4.00(4H,m), 3.74(2H,t, J=5.1Hz) and 2.90(3H,s)ppm.

Step E: 4-[5-(4-Cyanobenzyl)-1-imidazolylethyl]-1-(3-chlorophenyl)piperazin-2-one.

- 164 -

A solution of 2-(-5-(4-cyanobenzyl)-imidazolyl)ethyl methanesulfonate (24 mg, 0.079 mmol) in DMF (0.2ml) was added to 3-(chlorophenyl)piperazin-2-one (17.7mg, 0.084mmol), sodium iodide (50mg, 0.336mmol) and Hunigs base (0.0146ml, 0.084mmol). The mixture was stirred at 55°C for 12 hours, and the solvent evaporated in vacuo. The residue was purified by preparative tlc eluting with 10% saturated ammonia/acetonitrile to afford the title compound.

¹H NMR δ CDCl₃ (7.61 (2H,d, J=8.4Hz), 7.56(1H,s), 7.35-7.20 (7H,m), 7.16(1H,d, J=8Hz), 6.85(1H,s), 4.03(2H,s), , 3.83(2H,t, J=6.5Hz), 3.61(2H,t, J=5.4Hz), 3.27(2H,s), 2.68(2H,t, J=5.4Hz) and 2.57(2H,t, J=6.5Hz)ppm.

EXAMPLE 54

15 *In vitro* inhibition of ras farnesyl transferase

Assays of farnesyl-protein transferase. Partially purified bovine FPTase and Ras peptides (Ras-CVLS, Ras-CVIM and Ras-CAIL) were prepared as described by Schaber *et al.*, *J. Biol. Chem.* 265:14701-14704 (1990), Pompliano, *et al.*, *Biochemistry* 31:3800 (1992) and Gibbs *et al.*, *PNAS* U.S.A. 86:6630-6634 (1989), respectively. Bovine FPTase was assayed in a volume of 100 µl containing 100 mM N-(2-hydroxy ethyl) piperazine-*N'*-(2-ethane sulfonic acid) (HEPES), pH 7.4, 5 mM MgCl₂, 5 mM dithiothreitol (DTT), 100 mM [³H]-farnesyl diphosphate ([³H]-FPP; 740 CBq/mmol, New England Nuclear), 650 nM Ras-CVLS and 10 µg/ml FPTase at 31°C for 60 min. Reactions were initiated with FPTase and stopped with 1 ml of 1.0 M HCL in ethanol. Precipitates were collected onto filter-mats using a TomTec Mach II cell harvester, washed with 100% ethanol, dried and counted in an LKB β-plate counter. The assay was linear with respect to both substrates, FPTase levels and time; less than 10% of the [³H]-FPP was utilized during the reaction period. Purified compounds were dissolved in 100% dimethyl sulfoxide (DMSO) and were diluted 20-fold into the assay. Percentage inhibition is measured by the amount of incorporation of radioactivity in the presence of the test compound

- 165 -

when compared to the amount of incorporation in the absence of the test compound.

Human FPTase was prepared as described by Omer *et al.*, Biochemistry 32:5167-5176 (1993). Human FPTase activity was assayed as described above with the exception that 0.1% (w/v) polyethylene glycol 20,000, 10 μM ZnCl_2 and 100 nM Ras-CVIM were added to the reaction mixture. Reactions were performed for 30 min., stopped with 100 μl of 30% (v/v) trichloroacetic acid (TCA) in ethanol and processed as described above for the bovine enzyme.

The compounds of the instant invention described in the above Examples and in the Tables hereinafter were tested for inhibitory activity against human FPTase by the assay described above and were found to have IC_{50} of $\leq 50 \mu\text{M}$.

EXAMPLE 55

In vivo ras farnesylation assay

The cell line used in this assay is a v-ras line derived from either Rat1 or NIH3T3 cells, which expressed viral Ha-ras p21. The assay is performed essentially as described in DeClue, J.E. *et al.*, Cancer Research 51:712-717, (1991). Cells in 10 cm dishes at 50-75% confluency are treated with the test compound (final concentration of solvent, methanol or dimethyl sulfoxide, is 0.1%). After 4 hours at 37°C, the cells are labelled in 3 ml methionine-free DMEM supplemented with 10% regular DMEM, 2% fetal bovine serum and 400 mCi [^{35}S]methionine (1000 Ci/mmol). After an additional 20 hours, the cells are lysed in 1 ml lysis buffer (1% NP40/20 mM HEPES, pH 7.5/5 mM MgCl_2 /1mM DTT/10 mg/ml aprotinin/2 mg/ml leupeptin/2 mg/ml antipain/0.5 mM PMSF) and the lysates cleared by centrifugation at 100,000 x g for 45 min. Aliquots of lysates containing equal numbers of acid-precipitable counts are brought to 1 ml with IP buffer (lysis buffer lacking DTT) and immunoprecipitated with the ras-specific monoclonal antibody Y13-259 (Furth, M.E. *et al.*, J. Virol. 43:294-304, (1982)). Following a 2 hour antibody incubation at 4°C, 200 ml of a

- 166 -

25% suspension of protein A-Sepharose coated with rabbit anti rat IgG is added for 45 min. The immunoprecipitates are washed four times with IP buffer (20 nM HEPES, pH 7.5/1 mM EDTA/1% Triton X-100.0.5% deoxycholate/0.1%/SDS/0.1 M NaCl) boiled in SDS-PAGE sample buffer and loaded on 13% acrylamide gels. When the dye front reached the bottom, the gel is fixed, soaked in Enlightening, dried and autoradiographed. The intensities of the bands corresponding to farnesylated and nonfarnesylated ras proteins are compared to determine the percent inhibition of farnesyl transfer to protein.

10

EXAMPLE 56

In vivo growth inhibition assay

To determine the biological consequences of FPTase inhibition, the effect of the compounds of the instant invention on the anchorage-independent growth of Rat1 cells transformed with either a *v-ras*, *v-raf*, or *v-mos* oncogene is tested. Cells transformed by v-Raf and v-Mos maybe included in the analysis to evaluate the specificity of instant compounds for Ras-induced cell transformation.

20 Rat 1 cells transformed with either *v-ras*, *v-raf*, or *v-mos* are seeded at a density of 1×10^4 cells per plate (35 mm in diameter) in a 0.3% top agarose layer in medium A (Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum) over a bottom agarose layer (0.6%). Both layers contain 0.1% methanol or an appropriate concentration of the instant compound (dissolved in methanol at 1000 times the final concentration used in the assay). The cells are fed twice weekly with 0.5 ml of medium A containing 0.1% methanol or the concentration of the instant compound.

25 Photomicrographs are taken 16 days after the cultures are seeded and
30 comparisons are made.

- 167 -


EXAMPLE 57

Tables 1-18 show other compounds of the instant invention that were prepared by the procedures described in Examples 1-53.

- 5 These compounds are meant to be illustrative and are not meant to be limiting. In Table 2, the stereochemistry of the amino-containing carbon atom is as indicated (R or S) or, if unknown, the two separated stereoisomers are delineated by an "A" or "B" designation.

- 168 -


Table 1



R	FAB ms (m+1)	Formula Analysis (Calc, Found)
	526	$C_{33}H_{39}N_3OS \cdot 2.3 \text{ TFA} \cdot 0.9 \text{ H}_2O$ C, 56.16; H, 5.40; N, 7.22. C, 56.12; H, 5.37; N, 7.39.
	506	$C_{30}H_{39}N_3O_2S \cdot 2.55 \text{ TFA} \cdot 0.55 \text{ H}_2O$ C, 53.19; H, 5.46; N, 5.36. C, 53.17; H, 5.46; N, 5.60.
	494	$C_{29}H_{36}N_3OS \cdot 2.25 \text{ TFA} \cdot 0.35 \text{ H}_2O$ C, 53.19; H, 5.18; N, 5.55. C, 53.16; H, 5.18; N, 5.68.
	490	$C_{30}H_{39}N_3OS \cdot 2 \text{ TFA} \cdot 0.9 \text{ H}_2O$ C, 55.64; H, 5.88; N, 5.73. C, 55.57; H, 5.81; N, 5.83.
	512	$C_{29}H_{35}F_2N_3OS \cdot 2.4 \text{ TFA} \cdot 0.6 \text{ H}_2O$ C, 50.99; H, 4.89; N, 5.28. C, 50.98; H, 4.86; N, 5.56.

- 169 -

Table 2



5

				FAB mass spectrum	
amine					
10	stereo.	X	Y	(M+1)	Analysis (calcd., found)
	A	n-Bu	2-O(CH ₂) ₁₃ CH ₃	642.5	C ₄₂ H ₆₃ N ₃ O ₂ •2.55HCl C, 68.46; H, 8.97; N, 5.70. C, 68.40; H, 8.70; N, 5.61.
15	B	n-Bu	2-O(CH ₂) ₁₃ CH ₃	642.6	C ₄₂ H ₆₃ N ₃ O ₂ •2.65HCl C, 68.63; H, 8.99; N, 5.72. C, 68.66; H, 8.90; N, 5.70.
	A	n-Bu	2-O(CH ₂) ₁₁ CH ₃	614.4	C ₄₀ H ₅₉ N ₃ O ₂ •2.8HCl C, 67.09; H, 8.70; N, 5.87. C, 67.03; H, 8.66; N, 5.76.
20					
	B	n-Bu	2-O(CH ₂) ₁₁ CH ₃	614.4	C ₄₀ H ₅₉ N ₃ O ₂ •2.85HCl•0.05H ₂ O C, 66.84; H, 8.69; N, 5.85. C, 67.20; H, 8.70; N, 5.45.
25					

- 170 -


				FAB mass	
amine				spectrum	
stereo.	X	Y	(M+1)	Analysis (calcd., found)	
5	A	n-Bu	2-O(CH ₂) ₉ CH ₃	586.5	C ₃₈ H ₅₅ N ₃ O ₂ •3HCl•0.25hexane C, 66.18; H, 8.65; N, 5.86. C, 66.14; H, 8.44; N, 5.50.
10	B	n-Bu	2-O(CH ₂) ₉ CH ₃	586.5	C ₃₈ H ₅₅ N ₃ O ₂ •3HCl•0.15hexane C, 65.97; H, 8.55; N, 5.93. C, 66.04; H, 8.41; N, 5.57.
15	A	n-Bu	2-O(CH ₂) ₇ CH ₃	558	C ₃₆ H ₅₁ N ₃ O ₂ •2.15HCl•0.65H ₂ O C, 66.73; H, 8.47; N, 6.49. C, 66.74; H, 8.46; N, 6.52.
20	B	n-Bu	2-O(CH ₂) ₇ CH ₃	558	C ₃₆ H ₅₁ N ₃ O ₂ •2.15HCl•0.65H ₂ O C, 66.82; H, 8.42; N, 6.49. C, 66.85; H, 8.43; N, 6.52.
25	A	n-Bu	2-O(CH ₂) ₃ Ph	564	C ₃₇ H ₄₅ N ₃ O ₂ •2HCl•1.05H ₂ O C, 67.78; H, 7.55; N, 6.41. C, 67.78; H, 7.52; N, 6.26.
25	B	n-Bu	2-O(CH ₂) ₃ Ph	564	C ₃₇ H ₄₅ N ₃ O ₂ •2HCl•1.05H ₂ O C, 67.78; H, 7.55; N, 6.41. C, 67.78; H, 7.44; N, 6.33.
30	A	n-Bu	3-OCH ₂ Ph	536	C ₃₅ H ₄₁ N ₃ O ₂ •3.55HCl C, 63.19; H, 6.75; N, 6.32. C, 63.12; H, 6.61; N, 6.47.
30	B	n-Bu	3-OCH ₂ Ph	536	C ₃₅ H ₄₁ N ₃ O ₂ •3.55HCl•0.4H ₂ O C, 62.52; H, 6.80; N, 6.25. C, 62.14; H, 6.69; N, 6.65.

- 171 -

	amine stereo.	X	Y	FAB mass spectrum (M+1)	Analysis (calcd., found)
5	A	n-Bu	3-O(CH ₂) ₁₃ CH ₃	642	C ₄₂ H ₆₃ N ₃ O ₂ •2.4HCl C, 69.15; H, 9.04; N, 5.76. C, 69.08; H, 9.08; N, 5.73.
10	B	n-Bu	3-O(CH ₂) ₁₃ CH ₃	642.5	C ₄₂ H ₆₃ N ₃ O ₂ •2.35HCl•0.25H ₂ O C, 69.15; H, 9.04; N, 5.76. C, 69.08; H, 9.08; N, 5.73.
15	R	n-Bu	4-O(CH ₂) ₁₃ CH ₃	642	C ₄₂ H ₆₃ N ₃ O ₂ •2.2HCl•1H ₂ O C, 68.14; H, 9.15; N, 5.68. C, 68.13; H, 9.03; N, 5.68.
20	S	n-Bu	4-O(CH ₂) ₁₃ CH ₃	642	C ₄₂ H ₆₃ N ₃ O ₂ •2.8HCl C, 67.79; H, 8.91; N, 5.65. C, 67.77; H, 9.80; N, 5.64.
25	S	n-Bu	4-OCH ₂ Ph	536	C ₃₅ H ₄₁ N ₃ O ₂ •2.15HCl•1H ₂ O C, 66.49; H, 7.20; N, 6.65. C, 66.48; H, 7.15; N, 6.62.
	R/S	H	2-O(CH ₂) ₁₃ CH ₃	586	C ₃₈ H ₅₅ N ₃ O ₂ •1.5HCl•0.3H ₂ O C, 66.88; H, 8.58; N, 6.16. C, 66.82; H, 8.64; N, 6.15.

- 172 -


Table 3



5	Y	FAB mass spectrum (M+1)	Analysis (calcd., found)
10	3-SO ₂ Me	506	C ₂₇ H ₃₁ N ₅ O ₃ S•3.0 HCl•0.10 H ₂ O C, 52.61; H, 5.59; N, 11.36. C, 52.63; H, 5.41; N, 10.72.
15	3-OCF ₃	512.23	C ₂₇ H ₂₈ N ₅ O ₂ F ₃ •2.30 HCl•0.50 H ₂ O C, 53.66; H, 5.22; N, 11.59. C, 53.73; H, 5.23; N, 10.86.


- 173 -

Table 4



5	FAB mass spectrum			Analysis (calcd., found)
	X	Y	(M+1)	
10	CH ₂ OCH ₂ Ph	CF ₃	574	C ₃₂ H ₃₀ N ₅ O ₂ F ₃ •1.40HCl•0.30H ₂ O C, 61.02; H, 5.12; N, 11.12. C, 61.01; H, 5.10; N, 10.78.
15	CH ₂ SO ₂ Et	CF ₃	560	C ₂₇ H ₂₈ N ₅ O ₃ F ₃ S •3.20HCl•0.10H ₂ O C, 47.86; H, 4.67; N, 10.34. C, 47.89; H, 4.49; N, 9.92.
20	CH ₂ SO ₂ Ph	CF ₃	608	C ₃₁ H ₂₈ N ₅ O ₃ F ₃ S•2.10HCl C, 54.42; H, 4.43; N, 10.24. C, 54.43; H, 4.42; N, 9.98.
25	CH ₂ SO ₂ Me	Cl	512	C ₂₅ H ₂₆ N ₅ O ₃ ClS •2.0HCl•0.25CHCl ₃ •0.25H ₂ O C, 48.97; H, 4.68; N, 11.31. C, 48.95; H, 4.64; N, 11.51.


- 174 -

Table 4 (continued)

5	FAB mass spectrum			Analysis (calcd., found)
	X	Y	(M+1)	
10	CH ₂ SO ₂ Et	Cl	526	C ₂₆ H ₂₈ N ₅ O ₃ ClS•2HCl•0.25CHCl ₃ C, 50.14; H, 4.85; N, 11.14. C, 50.20; H, 4.91; N, 10.93.
15	CONHMe	CF ₃	511	C ₂₆ H ₂₅ N ₆ O ₂ F ₃ •1.90HCl•1.70H ₂ O C, 51.20; H, 5.01; N, 13.78. C, 51.17; H, 4.99; N, 13.36.
20	CONHEt	CF ₃	525	C ₂₇ H ₂₇ N ₆ O ₂ F ₃ •1.60HCl•0.10H ₂ O C, 55.54; H, 4.97; N, 14.39. C, 55.57; H, 4.87; N, 15.28.
25	CH ₂ SO ₂ Ph	Cl	574	C ₃₀ H ₂₈ N ₅ O ₃ ClS•2HCl•0.30H ₂ O C, 55.23; H, 4.73; N, 10.73. C, 55.25; H, 4.57; N, 10.61.
30				

- 175 -


Table 4 (continued)



			FAB mass spectrum (M+1)	Analysis (calcd., found)
5	X	Y		
10	CONHMe	Cl	477	C ₂₅ H ₂₅ N ₆ O ₂ Cl•2HCl •0.105CHCl ₃ •0.85H ₂ O C, 52.24; H, 5.03; N, 14.56. C, 52.21; H, 5.07; N, 14.98.
15	CONHEt	Cl	491	C ₂₆ H ₂₇ N ₆ O ₂ Cl•2HCl•0.25CHCl ₃ C, 53.10; H, 4.97; N, 14.15 C, 53.32; H, 5.21; N, 13.77.
20	CONHc-Pr	Cl	503	C ₂₇ H ₂₇ N ₆ O ₂ Cl•2HCl•0.40CHCl ₃ C, 52.77; H, 4.75; N, 13.47. C, 53.01; H, 4.99; N, 13.32.
25	CONHc-Pr	CF ₃	537	C ₂₈ H ₂₇ N ₆ O ₂ F ₃ •2.45HCl•0.55dioxane C, 53.81; H, 5.06; N, 12.47. C, 53.76; H, 5.09; N, 12.44.
30				


- 176 -

Table 4 (continued)



5	FAB mass spectrum			Analysis (calcd., found)
	X	Y	(M+1)	
10	NHCOMe	Cl	477	C ₂₅ H ₂₅ N ₆ O ₂ Cl •2HCl•0.60CHCl ₃ •2.0H ₂ O C, 46.76; H, 4.84; N, 12.78. C, 46.79; H, 4.46; N, 12.37.
15	CONMe ₂	CF ₃	525	C ₂₇ H ₂₇ N ₆ O ₂ F ₃ •2.0HCl•0.10H ₂ O C, 54.12; H, 4.91; N, 14.02. C, 54.10; H, 4.96; N, 13.79.
20	SO ₂ Et	Cl	512	C ₂₅ H ₂₆ N ₅ O ₃ ClS•2.0HCl•0.10H ₂ O C, 51.18; H, 4.84; N, 11.94. C, 51.24; H, 5.19; N, 11.10.
25	CH ₂ SMe	Cl	480	C ₂₅ H ₂₆ N ₅ OClS •0.15CHCl ₃ •0.05H ₂ O C, 52.83; H, 4.98; N, 12.25. C, 52.82; H, 5.36; N, 11.85.
30				

- 177 -

Table 4 (continued)FAB mass
spectrum

5


<u>X</u>	<u>Y</u>	<u>(M+1)</u>	<u>Analysis (calcd., found)</u>
(±) C≡CMe	Cl	458	C ₂₆ H ₂₄ N ₅ OCl•2.0HCl•1.00H ₂ O

10

C, 56.89; H, 5.14; N, 12.76.
C, 56.99; H, 5.20; N, 12.42.


- 178 -

Table 5




5	FAB mass spectrum		
	X	Y	(M+1) Analysis (calcd., found)
10	CH ₂ CH ₂	H	386 C ₂₃ H ₂₃ N ₅ O•1.40HCl•0.40H ₂ O C, 62.29; H, 5.73; N, 15.79. C, 26.26; H, 5.71; N, 15.43.
15	CH ₂ CO	H	400 C ₂₃ H ₂₁ N ₅ O ₂ •2.6HCl•1.70H ₂ O C, 52.71; H, 5.19; N, 13.36. C, 52.82; H, 5.21; N, 13.04.
20	CH ₂	H	372 C ₂₂ H ₂₁ N ₅ O•2.0HCl•2.60H ₂ O C, 53.80; H, 5.79; N, 14.26. C, 52.86; H, 5.98; N, 13.92.
25	CH ₂	3-Cl	406 C ₂₂ H ₂₀ N ₅ OCl•2.50HCl•0.90H ₂ O C, 51.48; H, 4.77; N, 13.65. C, 51.55; H, 4.75; N, 13.34.
30			

- 179 -

Table 5 (continued)


5	FAB mass spectrum			Analysis (calcd., found)
	X	Y	(M+1)	
10	NHSO ₂	H		C ₂₁ H ₂₀ N ₆ O ₃ S•1.0HCl•2.60H ₂ O C, 48.53; H, 5.08; N, 16.17 C, 48.60; H, 5.19; N, 15.80.
15	CH ₂ CH ₂ CO	3-Cl	448	C ₂₄ H ₂₂ N ₅ O ₂ Cl•1.0HCl•1.10H ₂ O C, 57.17; H, 5.04; N, 13.89. C, 57.22; H, 4.94; N, 13.47.
20	CH ₂	2,3-Cl ₂	440	C ₂₂ H ₁₉ N ₅ OCl ₂ •2.0HCl•0.60H ₂ O C, 50.42; H, 4.27; N, 13.36. C, 50.51; H, 4.56; N, 12.18.
25	CH ₂	2-Br	450	C ₂₂ H ₂₀ N ₅ OBr•2.0HCl•0.30H ₂ O C, 49.98; H, 4.31; N, 13.25. C, 49.94; H, 4.47; N, 12.53.
30				

- 180 -

Table 5 (continued)

		FAB mass spectrum	
		(M+1)	Analysis (calcd., found)
5	X	Y	
	CH ₂	3-CF ₃	440 C ₂₃ H ₂₀ N ₅ OF ₃ •1.70HCl•0.40H ₂ O
10			C, 54.40; H, 4.47; N, 13.79. C, 54.45; H, 4.49; N, 13.82.
	CH ₂	4-Cl	C ₂₂ H ₂₀ N ₅ OCl•1.50HCl•0.80H ₂ O
15			C, 55.71; H, 4.91; N, 14.77. C, 55.81; H, 4.94; N, 14.11.
20	CH ₂ CO	3-Cl	C ₂₃ H ₂₀ N ₅ O ₂ Cl•1.40HCl•1.10H ₂ O
			C, 54.73; H, 4.71; N, 13.87. C, 54.80; H, 4.73; N, 13.46.
25	CH ₂	3-F	C ₂₂ H ₂₀ N ₅ OF•2HCl •0.35CHCl ₃ •0.95H ₂ O
			C, 51.50; H, 4.69; N, 13.44. C, 51.56; H, 4.73; N, 13.30.
30			


- 181 -

Table 5 (continued)

5	FAB mass spectrum		
	X	Y	(M+1) Analysis (calcd., found)
	CH ₂	3-Br	C ₂₂ H ₂₀ N ₅ OBr•1.40HCl•1.30H ₂ O
10			C, 50.35; H, 4.61; N, 13.34. C, 50.36; H, 4.63; N, 12.84.


- 182 -

Table 6



5	FAB mass spectrum				
	X	Y	Z	(M+1)	Analysis (calcd., found)
10	4-CN	CH ₂	5-CH ₂ CH ₂ CO	554	C ₂₇ H ₂₈ N ₅ O ₄ ClS•1.30HCl•1.20H ₂ O C, 52.05; H, 5.13; N, 11.24. C, 52.09; H, 5.15; N, 10.95.
15	3-CN	CH ₂	5-CH ₂	512	C ₂₅ H ₂₆ N ₅ O ₃ ClS• 2.0HCl•0.10CHCl ₃ •0.60H ₂ O C, 49.61; H, 4.86; N, 11.52. C, 49.63; H, 5.01; N, 11.14.
20	2-CN	CH ₂	5-CH ₂	512	C ₂₅ H ₂₆ N ₅ O ₃ ClS•2HCl•0.35H ₂ O C, 50.79; H, 4.89; N, 11.84. C, 50.82; H, 5.29; N, 11.90.
25	4-CN	CH ₂	4-CH ₂ CH ₂ CO	554	C ₂₇ H ₂₈ N ₅ O ₄ ClS •2.0HCl•0.50CHCl ₃ •0.60H ₂ O C, 47.36; H, 4.58; N, 10.04. C, 47.35; H, 4.60; N, 9.66.
30					

- 183 -

Table 6 (continued)

5

FAB mass
spectrum

X	Y	Z	(M+1)	Analysis (calcd., found)
---	---	---	-------	--------------------------

4-CN	(CH ₂) ₂	4-CH ₂ CH ₂ CO		
------	---------------------------------	--------------------------------------	--	--


10

C₂₈H₃₀N₅O₄ClS•1.0HCl•0.40CHCl₃•0.95H₂O

C, 50.96; H, 5.01; N, 10.46.


C, 50.91; H, 5.02; N, 10.13.

- 184 -

Table 7

	X	Y	FAB mass spectrum (M+1)	Analysis (calcd., found)
5				
10	4-Me	CH ₂ CH ₂ SO ₂ Me	526	C ₂₆ H ₂₈ N ₅ O ₃ ClS •2HCl•0.35CHCl ₃ •0.85H ₂ O C, 48.24; H, 4.92; N, 10.67. C, 48.25; H, 4.93; N, 10.36.
15	4-Me	H	420	C ₂₃ H ₂₂ N ₅ OCl•2.0HCl•0.90H ₂ O C, 54.27; H, 5.11; N, 13.76. C, 54.21; H, 5.37; N, 12.97.
20				
	2-Me	H	420	C ₂₃ H ₂₂ N ₅ OCl•3.20HCl•0.10H ₂ O C, 51.31; H, 4.76; N, 13.01. C, 51.30; H, 4.74; N, 12.87.
25				


- 185 -

TABLE 8

Y	FAB ms (m+1)	Formula	
		Analysis	Calc Found
3-CF ₃	496	C ₂₇ H ₂₈ F ₃ N ₅ O · 2 HCl · 1.55 H ₂ C	
		C, 54.36; H, 5.59; N, 11.74	
		C, 54.40; H, 5.29; N, 11.26	
3-CH ₃	442	C ₂₇ H ₃₁ N ₅ O · 2 HCl	
		C, 63.01; H, 6.47; N, 13.61	
		C, 63.45; H, 6.71; N, 13.53	
2,3-(CH ₂) ₄	482	C ₃₀ H ₃₅ N ₅ O · 5.2 HCl · 0.1 H ₂ C	
		C, 53.66; H, 6.06; N, 10.43	
		C, 53.62; H, 5.30; N, 9.35	
3-OCH ₃	458	C ₂₇ H ₃₁ N ₅ O ₂ · 2 HCl	
		C, 61.13; H, 6.27; N, 13.20	
		C, 62.43; H, 6.53; N, 13.24	


- 186 -

TABLE 8 (continued)



Y	FAB ms (m+1)	Formula	
		Analysis	Calc Found
3-Cl	463	$C_{26}H_{28}ClN_5O \cdot 2 HCl \cdot 0.75 H_2O$ C, 56.94; H, 5.79; N, 12.77 C, 56.92; H, 5.59; N, 12.30	
2-CH ₃ , 3-Cl	477	$C_{27}H_{30}ClN_5O_2 \cdot 2 HCl \cdot 3.1 H_2O$ C, 53.62; H, 6.37; N, 11.58 C, 53.44; H, 5.37; N, 10.76	
H	428	$C_{26}H_{29}N_5O \cdot 3.9 HCl \cdot 2 H_2O$ C, 51.63; H, 6.51; N, 11.58 C, 51.59; H, 5.10; N, 10.35	


- 187 -

TABLE 8 (continued)


Y	FAB ms (m+1)	Formula	
		Analysis	Calc Found
3-F	446	$C_{26}H_{28}FN_5O \cdot 2 HCl \cdot 2.55 H_2O$ C, 55.87; H, 6.22; N, 12.53 C, 55.83; H, 5.84; N, 11.71	

- 188 -


TABLE 9



Formula

R	FAB ms (m+1)	Analysis	Calc Found
$\text{CH}_2\text{CH}_2\text{OCH}_2\text{CF}_3$	566	$\text{C}_{27}\text{H}_{25}\text{F}_6\text{N}_5\text{O}_2 \cdot 2 \text{HCl} \cdot 0.67 \text{H}_2\text{O}$ C, 49.85; H, 4.39; N, 10.77 C, 49.86; H, 4.26; N, 10.33	
 OCH_2CH_2	524	$\text{C}_{28}\text{H}_{28}\text{F}_3\text{N}_5\text{O}_2 \cdot 2 \text{HCl}$ C, 56.38; H, 5.07; N, 11.74 C, 56.21; H, 5.27; N, 11.46	
$\text{CH}_2\text{CH}_2\text{N}_3$	509	$\text{C}_{25}\text{H}_{23}\text{F}_3\text{N}_8\text{O}_2 \cdot 2 \text{HCl} \cdot 2.35 \text{H}_2\text{O}$ C, 48.14; H, 4.80; N, 17.96 C, 48.79; H, 4.38; N, 16.99	
$\text{CH}_2\text{CH}_2\text{NHCOCH}_3$	525	$\text{C}_{27}\text{H}_{27}\text{F}_3\text{N}_6\text{O}_2 \cdot 2 \text{HCl} \cdot 0.7 \text{H}_2\text{O}$ C, 53.16; H, 5.02; N, 13.78 C, 53.30; H, 5.07; N, 13.39	
$\text{CH}_2\text{CH}_2\text{NHCOC}_2\text{H}_5$	539	$\text{C}_{28}\text{H}_{29}\text{F}_3\text{N}_6\text{O}_2 \cdot 2 \text{HCl} \cdot 3.66 \text{H}_2\text{O}$ C, 52.43; H, 5.39; N, 13.10 C, 52.44; H, 5.20; N, 12.48	








- 189 -

TABLE 9 (continued)


R	FAB ms (m+1)	Formula	
		Analysis	Calc Found
CH ₂ CH ₂ SO ₂ CH(CH ₃) ₂	574	C ₂₈ H ₃₀ F ₃ N ₅ O ₃ S · 2 HCl · 1.35 H ₂ O C, 50.13; H, 5.21; N, 10.44 C, 50.10; H, 4.88; N, 10.08	
CH ₂ CH ₂ SCH(CH ₃) ₂	542	C ₂₈ H ₃₀ F ₃ N ₅ OS · 2 HCl · 0.75 H ₂ O C, 53.55; H, 5.38; N, 11.51 C, 53.57; H, 5.33; N, 11.04	
CH ₂ CH ₂ S(O)CH(CH ₃) ₂	558	C ₂₈ H ₃₀ F ₃ N ₅ O ₂ S · 2 HCl · 0.7 H ₂ O C, 52.29; H, 5.23; N, 10.89 C, 52.27; H, 5.08; N, 10.35	

- 190 -

TABLE 10

R	FAB ms (m+1)	Formula	
		Analysis	Calc Found
			
	462	$C_{26}H_{31}N_5O_3 \cdot 2 CF_2CO_2H \cdot 2.8 H_2C$ C, 48.69; H, 5.26; N, 9.46 C, 48.70; H, 4.42; N, 9.12	
	470	$C_{29}H_{35}N_5O \cdot 2 HCl \cdot 0.3 CH_2Cl_2$ $\cdot 0.6 C_6H_{14}$ C, 63.76; H, 7.48; N, 11.30 C, 64.77; H, 7.07; N, 11.26	
	431	$C_{27}H_{34}N_4O \cdot 2 HCl$ C, 64.41; H, 7.21; N, 11.13 C, 66.35; H, 7.28; N, 11.25	
	445	$C_{28}H_{36}N_4O \cdot 2 HCl \cdot 1.3 H_2O$ C, 62.17; H, 7.56; N, 10.36 C, 62.15; H, 7.09; N, 9.83	
	470	$C_{29}H_{35}N_5O \cdot 2 HCl \cdot 3.05 H_2O$ C, 58.30; H, 7.27; N, 11.72 C, 58.30; H, 6.28; N, 10.90	
	548, 550	$C_{29}H_{34}BrN_5O \cdot 2 HCl \cdot 0.1 H_2O$ C, 55.89; H, 5.85; N, 11.24 C, 55.91; H, 5.96; N, 10.73	


- 191 -

TABLE 11

R	Y	FAB ms (m+1)	Formula	
			Analysis	Calc Found
	Cl	511	$C_{26}H_{27}ClN_4O_3S$ $\cdot 2 HCl \cdot 2.85 H_2O$	C, 49.16; H, 5.51; N, 8.82 C, 49.17; H, 4.74; N, 8.87
	F	496	$C_{25}H_{26}FN_5O_3S$ $\cdot 2 HCl \cdot 1.65 H_2O$	C, 50.20; H, 5.27; N, 11.71 C, 50.19; H, 4.90; N, 11.44
	Cl	527	$C_{26}H_{28}ClN_5O_3S$ $\cdot 2 HCl \cdot 0.55 H_2O$	C, 51.29; H, 5.15; N, 11.50 C, 51.32; H, 5.39; N, 11.31
	Cl	530	$C_{26}H_{32}ClN_5O_3S$ $\cdot 3 HCl \cdot 0.6 HCl$	C, 48.02; H, 5.61; N, 10.77 C, 48.30; H, 5.91; N, 9.78


- 192 -

TABLE 12





R	FAB ms (m+1)	Formula	
		Analysis	Calc Found
	545	$C_{26}H_{30}ClN_5O_4S$ $\cdot 2 HCl \cdot 0.75 H_2O$	C, 49.53; H, 5.36; N, 11.11 C, 49.56; H, 5.58; N, 10.65
	552	$C_{28}H_{29}ClN_5O_3S$ $\cdot 2 HCl$	C, 53.90; H, 5.01; N, 11.22 C, 54.06; H, 5.78; N, 10.67
	396	$C_{17}H_{20}ClN_4O_3S$ $\cdot 2 HCl \cdot 0.7 H_2O$	C, 42.41; H, 4.90; N, 11.64 C, 42.98; H, 5.49; N, 11.06
	544	$C_{26}H_{30}ClN_5O_4S$ $\cdot 2 HCl \cdot 0.75 H_2O$	C, 49.51; H, 5.36; N, 11.11 C, 49.74; H, 5.66; N, 10.11

- 193 -







TABLE 12 (continued)

Formula

FAB ms
(m+1)Analysis Calc
Found524 $C_{27}H_{27}ClN_4O_3S \cdot 0.55 CHCl_3$ C, 58.67; H, 4.92; N, 10.03
C, 58.71; H, 4.94; N, 9.95523 $C_{28}H_{28}ClN_3O_3S \cdot 0.1 CHCl_3$
 $\cdot 0.2 CH_3OH$ C, 62.89; H, 5.39; N, 7.78
C, 62.87; H, 5.37; N, 7.84


- 194 -

TABLE 13

		Formula
R	FAB ms (m+1)	Analysis Calc Found
	563	$C_{30}H_{31}ClN_4O_3S$ $\cdot 2 HCl \cdot 1.8 H_2O$ C, 53.90; H, 5.52; N, 8.38 C, 53.90; H, 5.38; N, 7.82
	540	$C_{27}H_{30}ClN_5O_3S$ $\cdot 2 HCl \cdot 1.35 H_2O$ C, 50.88; H, 5.49; N, 10.99 C, 51.37; H, 5.51; N, 10.20
	512	$C_{25}H_{26}ClN_5O_3S$ $\cdot 2 HCl \cdot 0.3 H_2O$ C, 50.86; H, 4.88; N, 11.86 C, 51.06; H, 5.33; N, 10.87
	415	$C_{28}H_{27}ClN_4O_3S$ $\cdot 3 HCl \cdot 4 H_2O$ C, 36.25; H, 6.42; N, 9.39 C, 38.12; H, 5.91; N, 7.41
	580	$C_{25}H_{30}ClN_5O_5S_2$ $\cdot 2 HCl \cdot 0.85 H_2O$ C, 44.93; H, 5.08; N, 10.48 C, 44.96; H, 5.08; N, 9.96

- 195 -

TABLE 14




Formula

R ¹	R ²	FAB ms (m+1)	Analysis	Calc Found
	H	355	C ₁₉ H ₁₉ ClN ₄ O · 2 HCl · 0.5 H ₂ O C, 52.25; H, 5.08; N, 12.83 C, 52.31; H, 5.14; N, 12.23	
	CH ₂ CH ₂ F	452	C ₂₄ H ₂₃ ClFN ₅ O · 2 HCl · 1.70 H ₂ O C, 51.90; H, 5.15; N, 12.61 C, 52.22; H, 5.10; N, 12.22	
	H	448	C ₂₄ H ₂₂ ClN ₅ O ₂ · 1 HCl 0.30 CHCl ₃ · 0.85 H ₂ O C, 54.50; H, 4.71; N, 13.08 C, 54.51; H, 4.69; N, 12.87	

- 196 -


TABLE 15



R	Y	FAB ms (m+1)	Formula	
			Analysis	Calc Found
	2,3-(CH ₂) ₄	426	C ₂₆ H ₂₇ N ₅ O · 2 HCl · 0.15 H ₂ O	C, 62.31; H, 5.89; N, 13.97 C, 62.38; H, 6.18; N, 13.27
	2-CH ₃ , 3-Cl	420	C ₂₃ H ₂₂ ClN ₅ O · 2 HCl · 1.0 H ₂ O	C, 54.08; H, 5.13; N, 13.71 C, 54.57; H, 5.77; N, 12.92
	3-Cl	420	C ₂₂ H ₁₈ ClN ₅ O ₂ · 2 HCl · 0.5 H ₂ O	C, 56.78; H, 4.33; N, 15.05 C, 56.79; H, 4.69; N, 13.47

- 197 -


TABLE 16



Formula

R	FAB ms (m+1)	Analysis	Calc Found
	506	$C_{32}H_{35}N_5O \cdot 3.2 CF_3CO_2H$ $\cdot 2.65 H_2O$	C, 54.14; H, 5.75; N, 9.15 C, 54.15; H, 5.74; N, 9.18
	484	$C_{29}H_{33}N_5O_2 \cdot 4 HCl \cdot 3 H_2O$	C, 51.03; H, 6.35; N, 10.26 C, 51.02; H, 5.66; N, 9.67
	506	$C_{32}H_{35}N_5O \cdot 2 HCl \cdot 1.7 H_2C$	C, 63.09; H, 6.68; N, 10.26 C, 51.02; H, 5.66; N, 9.67

- 198 -

Table 17

Formula

R	FAB ms (m+1)	Analysis	Calc Found
	347	$C_{21}H_{22}N_4O \cdot 0.2 CHCl_3$ $\cdot 0.4 CH_3OH$ C, 67.71; H, 6.26; N, 14.62 C, 67.71; H, 6.26; N, 14.53	
	358	$C_{23}H_{23}N_3O \cdot 0.55 CHCl_3$ C, 66.85; H, 5.61; N, 9.93 C, 66.87; H, 5.70; N, 10.03	


- 199 -

Table 18

5	FAB mass spectrum (M+1)	Analysis (calcd., found)
		$C_{29}H_{33}N_5O_2 \cdot 1.0HCl \cdot 0.25CHCl_3 \cdot 0.70H_2O$ C, 62.45; H, 6.39; N, 12.45. C, 62.45; H, 6.46; N, 12.37.
10		$C_{22}H_{20}N_5OCl \cdot 2.60HCl$ C, 53.97; H, 4.78; N, 14.30 C, 53.97; H, 5.21; N, 13.10.
15		$C_{26}H_{28}N_5O_3ClS \cdot 2.0HCl \cdot 1.20H_2O$ C, 50.32; H, 5.26; N, 11.28 C, 50.31; H, 5.31; N, 10.72.
20		

- 200 -


Table 18 (continued)

	FAB mass spectrum (M+1)	<u>Analysis (calcd., found)</u>
5		
		
10	420	$C_{23}H_{22}N_5OCl \cdot 2.0HCl \cdot 2.10H_2O$ C, 52.06; H, 5.36; N, 13.20 C, 51.99; H, 5.36; N, 12.57.

- 201 -

WHAT IS CLAIMED IS:

1. A compound which inhibits farnesyl-protein transferase of the formula A:



wherein:

R^{1a} and R^{1b} are independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂N-C(O)-, CN, NO₂, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-,
- c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocyclic, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂N-C(O)-, CN, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, and R¹¹OC(O)-NR¹⁰-;

R² and R³ are independently selected from: H; unsubstituted or substituted C₁-8 alkyl, unsubstituted or substituted C₂-8 alkenyl, unsubstituted or substituted C₂-8 alkynyl, unsubstituted or substituted aryl,

unsubstituted or substituted heterocycle, $\begin{array}{c} \text{NR}^6\text{R}^7 \\ | \\ \text{O} \end{array}$ or $\begin{array}{c} \text{OR}^6 \\ | \\ \text{O} \end{array}$,









wherein the substituted group is substituted with one or more of:

- 1) aryl or heterocycle, unsubstituted or substituted with:

- 202 -



5

- a) C₁₋₄ alkyl,
 b) (CH₂)_pOR⁶,
 c) (CH₂)_pNR⁶R⁷,
 d) halogen,
 e) CN,
 2) C₃₋₆ cycloalkyl,
 3) OR⁶,
 4) SR^{6a}, S(O)R^{6a}, SO₂R^{6a},



10

- 203 -





R^2 and R^3 are attached to the same C atom and are combined to form
 - $(CH_2)_u$ - wherein one of the carbon atoms is optionally replaced by a
 5 moiety selected from: O, $S(O)_m$, $-NC(O)-$, and $-N(COR^{10})-$;

R^4 and R^5 are independently selected from H and CH_3 ;

10 and any two of R^2 , R^3 , R^4 and R^5 are optionally attached to the
 same carbon atom;

R^6 , R^7 and R^{7a} are independently selected from: H; C₁₋₄ alkyl, C₃₋₆
 cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl,
 heteroarylsulfonyl, unsubstituted or substituted with:


- 15 a) C₁₋₄ alkoxy,
 b) aryl or heterocycle,
 c) halogen,
 d) HO,



R^6 and R^7 may be joined in a ring;
 R^7 and R^{7a} may be joined in a ring;

- 204 -

R^{6a} is selected from: C₁₋₄ alkyl, C₃₋₆ cycloalkyl, heterocycle, aryl, unsubstituted or substituted with:

- 5
- a) C₁₋₄ alkoxy,
 - b) aryl or heterocycle,
 - c) halogen,
 - d) HO,
 - e) 

$$\begin{array}{c} \text{R}^{11} \\ | \\ \text{C} \\ || \\ \text{O} \end{array}$$
 - f) $-\text{SO}_2\text{R}^{11}$, or
 - g) $\text{N}(\text{R}^{10})_2$;
- 10

R⁸ is independently selected from:

- a) hydrogen,
 - b) aryl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and
 - c) C₁₋₆ alkyl unsubstituted or substituted by aryl, cyanophenyl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NH-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹⁰OC(O)NH-;
- 15
- 20

25 R⁹ is selected from:

- a) hydrogen,
 - b) C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and
- 30

- 205 -

- c) C₁-C₆ alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

5

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

10

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR¹⁰-, -NR¹⁰C(O)-, O, -N(R¹⁰)-, -S(O)₂N(R¹⁰)-, -N(R¹⁰)S(O)₂-, or S(O)_m;

15 V is selected from:

- a) hydrogen,
- b) heterocycle,
- c) aryl,
- d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C₂-C₂₀ alkenyl,

20

provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

25 W is a heterocycle;

X is -CH₂-, -C(=O)-, or -S(=O)_m-;

30 Y is unsubstituted or substituted aryl or unsubstituted or substituted heterocycle, wherein the substituted aryl or substituted heterocycle is substituted with one or more of:

- 1) C₁-4 alkyl, unsubstituted or substituted with:
 - a) C₁-4 alkoxy,


- 206 -

- 5 b) NR^6R^7 ,
 c) C₃₋₆ cycloalkyl,
 d) aryl or heterocycle,
 e) HO,
 f) $-\text{S}(\text{O})_m\text{R}^{6a}$, or
 g) $-\text{C}(\text{O})\text{NR}^6\text{R}^7$,
 10 2) aryl or heterocycle,
 3) halogen,
 4) OR^6 ,
 5) NR^6R^7 ,
 6) CN,
 7) NO_2 ,
 8) CF_3 ;
 15 9) $-\text{S}(\text{O})_m\text{R}^{6a}$,
 10) $-\text{C}(\text{O})\text{NR}^6\text{R}^7$, or
 11) C_{3-C6} cycloalkyl;
- 20 m is 0, 1 or 2;
 n is 0, 1, 2, 3 or 4;
 p is 0, 1, 2, 3 or 4;
 q is 1 or 2;
 r is 0 to 5, provided that r is 0 when V is hydrogen;
 s is 0 or 1;
 t is 0 or 1; and
 25 u is 4 or 5;

or an optical isomer or pharmaceutically acceptable salt thereof.

30 2. A compound which inhibits farnesyl-protein
 transferase of the formula B:

- 207 -



wherein:

R^{1a} and R^{1b} are independently selected from:

- 5 a) hydrogen,
- b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, CN(R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
- 10 R¹¹OC(O)NR¹⁰,
- c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocyclic, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-,
- 15 R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, and R¹¹OC(O)-NR¹⁰;

20 R² and R³ are independently selected from: H; unsubstituted or substituted C₁-8 alkyl, unsubstituted or substituted C₂-8 alkenyl, unsubstituted or substituted C₂-8 alkynyl, unsubstituted or substituted aryl,

unsubstituted or substituted heterocycle, $\begin{array}{c} \text{NR}^6\text{R}^7 \\ | \\ \text{O} \end{array}$ or $\begin{array}{c} \text{OR}^6 \\ | \\ \text{O} \end{array}$,

wherein the substituted group is substituted with one or more of:

- 25 1) aryl or heterocycle, unsubstituted or substituted with:
 - a) C₁-4 alkyl,
 - b) (CH₂)_pOR⁶,

- 208 -

c) $(\text{CH}_2)_p\text{NR}^6\text{R}^7$,

d) halogen,

e) CN,

2) C₃₋₆ cycloalkyl,3) OR^6 ,4) SR^{6a} , $\text{S(O)}\text{R}^{6a}$, SO_2R^{6a} ,5) $-\text{NR}^6\text{R}^7$,6)
$$\begin{array}{c} \text{R}^6 \\ | \\ -\text{N}-\text{C}(=\text{O})-\text{R}^7 \end{array}$$
,7)
$$\begin{array}{c} \text{R}^6 \\ | \\ -\text{N}-\text{C}(=\text{O})-\text{NR}^7\text{R}^{7a} \end{array}$$
,8)
$$\begin{array}{c} -\text{O}-\text{C}(=\text{O})-\text{NR}^6\text{R}^7 \end{array}$$
,9)
$$\begin{array}{c} -\text{O}-\text{C}(=\text{O})-\text{OR}^6 \end{array}$$
,10)
$$\begin{array}{c} \text{C}(=\text{O})-\text{NR}^6\text{R}^7 \end{array}$$
,11) $-\text{SO}_2-\text{NR}^6\text{R}^7$,12)
$$\begin{array}{c} \text{R}^6 \\ | \\ -\text{N}-\text{SO}_2-\text{R}^{6a} \end{array}$$
,

- 209 -





R^2 and R^3 are attached to the same C atom and are combined to form
 - $(CH_2)_u$ - wherein one of the carbon atoms is optionally replaced by a
 5 moiety selected from: O, $S(O)_m$, $-NC(O)-$, and $-N(COR^{10})-$;

R^4 is selected from H and CH_3 ;

10 and any two of R^2 , R^3 and R^4 are optionally attached to the same
 carbon atom;

R^6 , R^7 and R^{7a} are independently selected from: H; C₁₋₄ alkyl, C₃₋₆
 cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl,
 heteroarylsulfonyl, unsubstituted or substituted with:

- 15 a) C₁₋₄ alkoxy,
 b) aryl or heterocycle,
 c) halogen,
 d) HO,






R^6 and R^7 may be joined in a ring;
 R^7 and R^{7a} may be joined in a ring;

- 210 -

R_{6a} is selected from: C₁₋₄ alkyl, C₃₋₆ cycloalkyl, heterocycle, aryl, unsubstituted or substituted with:

- a) C₁₋₄ alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,



R₈ is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and
- c) C₁₋₆ alkyl unsubstituted or substituted by aryl, cyanophenyl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NH-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹⁰OC(O)NH-;

R₉ is selected from:

- a) hydrogen,
- b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and

- 211 -

- c) C₁-C₆ alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

5

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

10

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR¹⁰-, -NR¹⁰C(O)-, O, -N(R¹⁰)-, -S(O)₂N(R¹⁰)-, -N(R¹⁰)S(O)₂-, or S(O)_m;

15

G is H₂ or O;

V is selected from:

- a) hydrogen,
- b) heterocycle,
- 20 c) aryl,
- d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C₂-C₂₀ alkenyl,

25

provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

W is a heterocycle;

30

X is -CH₂-, -C(=O)-, or -S(=O)_m-;

Z is a unsubstituted or substituted group selected from aryl, heteroaryl, arylmethyl, heteroarylmethyl, arylsulfonyl,

- 212 -

heteroarylsulfonyl, wherein the substituted group is substituted with one or more of the following:

- 1) C₁₋₄ alkyl, unsubstituted or substituted with:
 - a) C₁₋₄ alkoxy,
 - b) NR⁶R⁷,
 - c) C₃₋₆ cycloalkyl,
 - d) aryl or heterocycle,
 - e) HO,
 - f) -S(O)_mR^{6a}, or
 - g) -C(O)NR⁶R⁷,
- 2) aryl or heterocycle,
- 3) halogen,
- 4) OR⁶,
- 5) NR⁶R⁷,
- 6) CN,
- 7) NO₂,
- 8) CF₃;
- 9) -S(O)_mR^{6a},
- 10) -C(O)NR⁶R⁷, or
- 11) C₃₋₆ cycloalkyl;

- m is 0, 1 or 2;
 n is 0, 1, 2, 3 or 4;
 p is 0, 1, 2, 3 or 4;
 q is 1 or 2;
 r is 0 to 5, provided that r is 0 when V is hydrogen;
 s is 0 or 1;
 t is 0 or 1; and
 u is 4 or 5;

30

provided that when G is H₂ and W is imidazolyl, then the substituent (R⁸)_r-V-A¹(CR^{1a}₂)_nA²(CR^{1a}₂)_n- is not H and


- 213 -

provided that when X is -C(=O)-, or -S(=O)_m-, then t is 1 and the substituent (R⁸)_r V - A¹(CR^{1a}₂)_nA²(CR^{1a}₂)_n - is not H;

or an optical isomer or pharmaceutically acceptable salt thereof.

5

3. A compound which inhibits farnesyl-protein transferase of the formula C:



wherein:

10

R^{1a} and R^{1b} are independently selected from:



- a) hydrogen,
- b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-,
- c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocyclic, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, and R¹¹OC(O)-NR¹⁰-;

20

25 R² and R³ are independently selected from: H; unsubstituted or substituted C₁-8 alkyl, unsubstituted or substituted C₂-8 alkenyl,




- 214 -

unsubstituted or substituted C₂-8 alkynyl, unsubstituted or substituted aryl,






unsubstituted or substituted heterocycle,  or ,

wherein the substituted group is substituted with one or more of:

- 1) aryl or heterocycle, unsubstituted or substituted with:
 - a) C₁-4 alkyl,
 - b) (CH₂)_pOR⁶,
 - c) (CH₂)_pNR⁶R⁷,
 - d) halogen,
 - e) CN,
- 2) C₃-6 cycloalkyl,
- 3) OR⁶,
- 4) SR^{6a}, S(O)R^{6a}, SO₂R^{6a},



- 215 -



- 5 R^2 and R^3 are attached to the same C atom and are combined to form $\text{—(CH}_2)_u\text{—}$ wherein one of the carbon atoms is optionally replaced by a moiety selected from: O, S(O)_m , —NC(O)— , and $\text{—N(COR}^{10})\text{—}$;

R^4 is selected from H and CH_3 ;

10


and any two of R^2 , R^3 and R^4 are optionally attached to the same carbon atom;

- 216 -

R⁶, R⁷ and R^{7a} are independently selected from: H; C₁₋₄ alkyl, C₃₋₆ cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, unsubstituted or substituted with:

5

- a) C₁₋₄ alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,




- f) $-\text{SO}_2\text{R}^{11}$, or
- g) $\text{N}(\text{R}^{10})_2$; or

10

R⁶ and R⁷ may be joined in a ring;
R⁷ and R^{7a} may be joined in a ring;

15 R^{6a} is selected from: C₁₋₄ alkyl, C₃₋₆ cycloalkyl, heterocycle, aryl, unsubstituted or substituted with:

- a) C₁₋₄ alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,



20

- f) $-\text{SO}_2\text{R}^{11}$, or
- g) $\text{N}(\text{R}^{10})_2$;

R⁸ is independently selected from:

25

- a) hydrogen,
- b) aryl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN,

- 217 -

NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
R¹¹OC(O)NR¹⁰-, and

- c) C₁-C₆ alkyl unsubstituted or substituted by aryl,
cyanophenyl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆
alkenyl, C₂-C₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-,
R¹¹S(O)_m-, R¹⁰C(O)NH-, (R¹⁰)₂NC(O)-, R¹⁰₂N-
C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
R¹⁰OC(O)NH-;

10 R⁹ is selected from:

- a) hydrogen,
b) C₂-C₆ alkenyl, C₂-C₆ alkynyl, perfluoroalkyl, F, Cl, Br,
R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-,
R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃,
-N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and
c) C₁-C₆ alkyl unsubstituted or substituted by perfluoroalkyl,
F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-,
(R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-,
R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

20

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and
aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

25

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-,
-C(O)-, -C(O)NR¹⁰-, -NR¹⁰C(O)-, O, -N(R¹⁰)-,
-S(O)₂N(R¹⁰)-, -N(R¹⁰)S(O)₂-, or S(O)_m;

30

G is O;

V is selected from:

- a) hydrogen,
b) heterocycle,

- 218 -

- c) aryl,
 d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
 5 e) C₂-C₂₀ alkenyl,
 provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

W is a heterocycle;

10

X is -CH₂-, -C(=O)-, or -S(=O)_m-;

15

Z is a unsubstituted or substituted group selected from aryl, heteroaryl, arylmethyl, heteroarylmethyl, arylsulfonyl, heteroarylsulfonyl, wherein the substituted group is substituted with one or more of the following:

20

1) C₁₋₄ alkyl, unsubstituted or substituted with:

- a) C₁₋₄ alkoxy,
 b) NR⁶R⁷,
 c) C₃₋₆ cycloalkyl,
 d) aryl or heterocycle,
 e) HO,
 f) -S(O)_mR^{6a}, or
 g) -C(O)NR⁶R⁷,

25

2) aryl or heterocycle,

3) halogen,

4) OR⁶,

5) NR⁶R⁷,

6) CN,

30

7) NO₂,

8) CF₃;

9) -S(O)_mR^{6a},

10) -C(O)NR⁶R⁷, or

11) C₃-C₆ cycloalkyl;

- 219 -


- m is 0, 1 or 2;
 n is 0, 1, 2, 3 or 4;
 p is 0, 1, 2, 3 or 4;
 5 q is 1 or 2;
 r is 0 to 5, provided that r is 0 when V is hydrogen;
 s is 1;
 t is 0 or 1; and
 u is 4 or 5;

10

or an optical isomer or pharmaceutically acceptable salt thereof.

4. The compound according to Claim 1 of the formula

A:



15

wherein:

R^{1a} is independently selected from: hydrogen or C₁-C₆ alkyl;


20


R^{1b} is independently selected from:

- a) hydrogen,
 b) aryl, heterocycle, cycloalkyl, R¹⁰O-, -N(R¹⁰)₂ or C₂-C₆ alkenyl,
 25 c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocycle, cycloalkyl, alkenyl, R¹⁰O- and -N(R¹⁰)₂;

- 220 -

R³, R⁴ and R⁵ are independently selected from H and CH₃;

R² is H; ; or C₁-5 alkyl, unbranched or branched, unsubstituted or substituted with one or more of:

- 1) aryl,
- 2) heterocycle,
- 3) OR⁶,
- 4) SR^{6a}, SO₂R^{6a}, or
- 5) ;

and any two of R², R³, R⁴, and R⁵ are optionally attached to the same carbon atom;

R⁶, R⁷ and R^{7a} are independently selected from:

- H; C₁-4 alkyl, C₃-6 cycloalkyl, aryl, heterocycle, unsubstituted or substituted with:
- a) C₁-4 alkoxy,
 - b) halogen, or
 - c) aryl or heterocycle;

R^{6a} is selected from:

- C₁-4 alkyl or C₃-6 cycloalkyl, unsubstituted or substituted with:
- a) C₁-4 alkoxy,
 - b) halogen, or
 - c) aryl or heterocycle;

R⁸ is independently selected from:

- a) hydrogen,
- b) C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ perfluoroalkyl, F, Cl, R¹⁰O-, R¹⁰C(O)NR¹⁰-, CN, NO₂,

- 221 -

$(R^{10})_2N-C(NR^{10})-$, $R^{10}C(O)-$, $R^{10}OC(O)-$, $-N(R^{10})_2$, or $R^{11}OC(O)NR^{10}-$, and

- c) C_1-C_6 alkyl substituted by C_1-C_6 perfluoroalkyl, $R^{10}O-$, $R^{10}C(O)NR^{10}-$, $(R^{10})_2N-C(NR^{10})-$, $R^{10}C(O)-$, $R^{10}OC(O)-$, $-N(R^{10})_2$, or $R^{11}OC(O)NR^{10}-$;

R^9 is selected from:

- a) hydrogen,
 b) C_2-C_6 alkenyl, C_2-C_6 alkynyl, C_1-C_6 perfluoroalkyl, F, Cl, $R^{10}O-$, $R^{11}S(O)_m-$, $R^{10}C(O)NR^{10}-$, CN, NO_2 , $(R^{10})_2N-C(NR^{10})-$, $R^{10}C(O)-$, $R^{10}OC(O)-$, $-N(R^{10})_2$, or $R^{11}OC(O)NR^{10}-$, and
 c) C_1-C_6 alkyl unsubstituted or substituted by C_1-C_6 perfluoroalkyl, F, Cl, $R^{10}O-$, $R^{11}S(O)_m-$, $R^{10}C(O)NR^{10}-$, CN, $(R^{10})_2N-C(NR^{10})-$, $R^{10}C(O)-$, $R^{10}OC(O)-$, $-N(R^{10})_2$, or $R^{11}OC(O)NR^{10}-$;

R^{10} is independently selected from hydrogen, C_1-C_6 alkyl, benzyl and aryl;

R^{11} is independently selected from C_1-C_6 alkyl and aryl;

A^1 and A^2 are independently selected from: a bond, $-CH=CH-$, $-C\equiv C-$, $-C(O)-$, $-C(O)NR^{10}-$, O, $-N(R^{10})-$, or $S(O)_m$;

V is selected from:

- a) hydrogen,
 b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,
 c) aryl,
 d) C_1-C_{20} alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and

- 222 -

e) C₂-C₂₀ alkenyl, and
 provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen
 if A¹ is a bond, n is 0 and A² is S(O)_m;

5 W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or isoquinolinyl;

X is -CH₂- or -C(=O)-;

10

Y is mono- or bicyclic aryl, or mono- or bicyclic heterocycle, unsubstituted or substituted with one or more of:

15

- a) C₁₋₄ alkyl,
- b) C₁₋₄ alkoxy,
- c) halogen, or
- d) NR⁶R⁷;

m is 0, 1 or 2;

n is 0, 1, 2, 3 or 4;

20

p is 0, 1, 2, 3 or 4;

r is 0 to 5, provided that r is 0 when V is hydrogen;

s is 0 or 1; and


t is 0 or 1;

25 or an optical isomer or pharmaceutically acceptable salt thereof.

5. The compound according to Claim 2 of the formula

B:

- 223 -



wherein:

R^{1a} is independently selected from: hydrogen or C₁-C₆ alkyl;

R^{1b} is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, cycloalkyl, R¹⁰O-, -N(R¹⁰)₂ or C₂-C₆ alkenyl,
- c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocycle, cycloalkyl, alkenyl, R¹⁰O- and -N(R¹⁰)₂;

R³ and R⁴ are independently selected from H and CH₃;

R² is H; $\begin{array}{c} \text{NR}^6\text{R}^7 \\ | \\ \text{O} \end{array}$; or C₁-5 alkyl, unbranched or branched, unsubstituted or substituted with one or more of:

- 1) aryl,
- 2) heterocycle,
- 3) OR⁶,
- 4) SR^{6a}, SO₂R^{6a}, or
- 5) $\begin{array}{c} \text{NR}^6\text{R}^7 \\ | \\ \text{O} \end{array}$;

and any two of R², R³, R⁴, and R⁵ are optionally attached to the same carbon atom;

- 224 -

R⁶, R⁷ and R^{7a} are independently selected from:

H; C₁₋₄ alkyl, C₃₋₆ cycloalkyl, aryl, heterocycle,
unsubstituted or substituted with:

- 5 a) C₁₋₄ alkoxy,
 b) halogen, or
 c) aryl or heterocycle;

R^{6a} is selected from:

- 10 C₁₋₄ alkyl or C₃₋₆ cycloalkyl,
 unsubstituted or substituted with:
 a) C₁₋₄ alkoxy,
 b) halogen, or
 c) aryl or heterocycle;

15 R⁸ is independently selected from:

- a) hydrogen,
 b) C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆
 perfluoroalkyl, F, Cl, R¹⁰O-, R¹⁰C(O)NR¹⁰-, CN, NO₂,
 (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, -N(R¹⁰)₂, or
20 R¹¹OC(O)NR¹⁰-, and
 c) C₁₋₆ alkyl substituted by C₁₋₆ perfluoroalkyl, R¹⁰O-,
 R¹⁰C(O)NR¹⁰-, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-,
 R¹⁰OC(O)-, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

25 R⁹ is selected from:

- a) hydrogen,
 b) C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ perfluoroalkyl, F,
 Cl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, CN, NO₂,
 (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-, -N(R¹⁰)₂, or
30 R¹¹OC(O)NR¹⁰-, and
 c) C₁₋₆ alkyl unsubstituted or substituted by C₁₋₆
 perfluoroalkyl, F, Cl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-,
 CN, (R¹⁰)₂N-C(NR¹⁰)-, R¹⁰C(O)-, R¹⁰OC(O)-,
 -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

- 225 -

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and aryl;

5 R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR¹⁰-, O, -N(R¹⁰)-, or S(O)_m;

10 V is selected from:

- a) hydrogen,
- b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,
- 15 c) aryl,
- d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C₂-C₂₀ alkenyl, and

20 provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

G is H₂ or O;

25 W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or isoquinolinyl;

X is -CH₂- or -C(=O)-;

30

Z is mono- or bicyclic aryl, mono- or bicyclic heteroaryl, mono- or bicyclic arylmethyl, mono- or bicyclic heteroarylmethyl, mono- or bicyclic arylsulfonyl, mono- or bicyclic heteroarylsulfonyl, unsubstituted or

- 226 -

substituted with one or two of the following:

- 1) C₁₋₄ alkyl, unsubstituted or substituted with:
 - a) C₁₋₄ alkoxy,
 - b) NR⁶R⁷,
 - c) C₃₋₆ cycloalkyl,
 - d) aryl or heterocycle,
 - e) HO,
 - f) -S(O)_mR⁶, or
 - g) -C(O)NR⁶R⁷,
- 2) aryl or heterocycle,
- 3) halogen,
- 4) OR⁶,
- 5) NR⁶R⁷,
- 6) CN,
- 7) NO₂,
- 8) CF₃;
- 9) -S(O)_mR⁶,
- 10) -C(O)NR⁶R⁷, or
- 11) C₃₋₆ cycloalkyl;

- 20
- m is 0, 1 or 2;
- n is 0, 1, 2, 3 or 4;
- p is 0, 1, 2, 3 or 4;
- r is 0 to 5, provided that r is 0 when V is hydrogen;
- 25 s is 0 or 1;
- t is 0 or 1; and
- u is 4 or 5;

30 provided that when G is H₂ and W is imidazolyl, then the substituent (R⁸)_r-V-A¹(CR^{1a2})_nA²(CR^{1a2})_n- is not H and

provided that when X is -C(=O)-, or -S(=O)_m-, then t is 1 and the substituent (R⁸)_r-V-A¹(CR^{1a2})_nA²(CR^{1a2})_n- is not H;

- 227 -

or pharmaceutically acceptable salt thereof.

6. A compound which inhibits farnesyl-protein transferase which is:

- 5 2(S)-Butyl-1-(2,3-diaminoprop-1-yl)-4-(1-naphthoyl)-piperazine
- 10 1-(3-Amino-2-(2-naphthylmethylamino)prop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine
- 2(S)-Butyl-1-{5-[1-(2-naphthylmethyl)]-4,5-dihydroimidazol }methyl-4-(1-naphthoyl)piperazine
- 15 1-[5-(1-Benzylimidazol)methyl]-2(S)-butyl-4-(1-naphthoyl)piperazine
- 1-{5-[1-(4-Nitrobenzyl)imidazolyl]methyl}-2(S)-butyl-4-(1-naphthoyl)piperazine
- 20 1-(3-Acetamidomethylthio-2(R)-aminoprop-1-yl)-2(S)-butyl-4-(1-naphthoyl)piperazine
- 2(S)-Butyl-1-[2-(1-imidazolyl)ethyl]sulfonyl-4-(1-naphthoyl)piperazine
- 25 2(R)-Butyl-1-imidazolyl-4-methyl-4-(1-naphthoyl)piperazine
- 2(S)-Butyl-4-(1-naphthoyl)-1-(3-pyridylmethyl)piperazine
- 1-2(S)-Butyl-(2(R)-(4-nitrobenzyl)amino-3-hydroxypropyl)-4-(1-naphthoyl)piperazine
- 30 1-(2(R)-Amino-3-hydroxyheptadecyl)-2(S)-butyl-4-(1-naphthoyl)piperazine
- 35 2(S)-Benzyl-1-imidazolyl-4-methyl-4-(1-naphthoyl)piperazine

- 228 -

1-(2(R)-Amino-3-(3-benzylthio)propyl)-2(S)-butyl-4-(1-naphthoyl)-
piperazine

5 1-(2(R)-Amino-3-[3-(4-nitrobenzylthio)propyl]))-2(S)-butyl-4-(1-
naphthoyl)piperazine

2(S)-Butyl-1-[(4-imidazolyl)ethyl]-4-(1-naphthoyl)piperazine

10 2(S)-Butyl-1-[(4-imidazolyl)methyl]-4-(1-naphthoyl)piperazine

2(S)-Butyl-1-[(1-naphth-2-ylmethyl)-1H-imidazol-5-yl)acetyl]-4-(1-
naphthoyl)piperazine

15 2(S)-Butyl-1-[(1-naphth-2-ylmethyl)-1H-imidazol-5-yl)ethyl]-4-(1-
naphthoyl)piperazine

1-(2(R)-Amino-3-hydroxypropyl)-2(S)-butyl-4-(1-
naphthoyl)piperazine

20 1-(2(R)-Amino-4-hydroxybutyl)-2(S)-butyl-4-(1-naphthoyl)piperazine

1-(2-Amino-3-(2-benzyloxyphenyl)propyl)-2(S)-butyl-4-(1-
naphthoyl)piperazine

25 1-(2-Amino-3-(2-hydroxyphenyl)propyl)-2(S)-butyl-4-(1-
naphthoyl)piperazine

30 1-[3-(4-Imidazolyl)propyl]-2(S)-butyl-4-(1-
naphthoyl)piperazine

2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(1-naphthylmethyl)imidazol-5-
ylmethyl]-piperazine

35 2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(2-naphthylmethyl)imidazol-5-
ylmethyl]-piperazine

- 229 -

2(S)-*n*-Butyl-1-[1-(4-cyanobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine

5 2(S)-*n*-Butyl-1-[1-(4-methoxybenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine

2(S)-*n*-Butyl-1-[1-(3-methyl-2-butenyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine

10 2(S)-*n*-Butyl-1-[1-(4-fluorobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine

2(S)-*n*-Butyl-1-[1-(4-chlorobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine

15 1-[1-(4-Bromobenzyl)imidazol-5-ylmethyl]-2(S)-*n*-butyl-4-(1-naphthoyl)piperazine

20 1-[1-(4-Bromobenzyl)imidazol-5-ylmethyl]-2(S)-*n*-butyl-4-(1-naphthoyl)piperazine

2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(4-trifluoromethylbenzyl)imidazol-5-ylmethyl]-piperazine

25 2(S)-*n*-Butyl-1-[1-(4-methylbenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)-piperazine

2(S)-*n*-Butyl-1-[1-(3-methylbenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)-piperazine

30 1-[1-(4-Phenylbenzyl)imidazol-5-ylmethyl]-2(S)-*n*-butyl-4-(1-naphthoyl)-piperazine

35 2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(2-phenylethyl)imidazol-5-ylmethyl]-piperazine

- 230 -

2(S)-*n*-Butyl-4-(1-naphthoyl)-1-[1-(4-trifluoromethoxy)-imidazol-5-ylmethyl]piperazine

5 1- {[1-(4-cyanobenzyl)-1H-imidazol-5-yl]acetyl}-2(S)-*n*-butyl-4-(1-naphthoyl)piperazine

5(S)-*n*-Butyl-1-(2,3-dimethylphenyl)-4-(4-imidazolylmethyl)-piperazin-2-one

10 5(S)-*n*-Butyl-4-[1-(4-cyanobenzyl)imidazol-5-ylmethyl]-1-(2,3-dimethylphenyl)piperazin-2-one

15 4-[1-(4-Cyanobenzyl)imidazol-5-ylmethyl]-1-(2,3-dimethylphenyl)-5(S)-(2-methoxyethyl)piperazin-2-one

(S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(methanesulfonyl)ethyl]-2-piperazinone

20 (S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)ethyl]-2-piperazinone

(S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)methyl]-2-piperazinone

25 (S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[*N*-ethyl-2-acetamido]-2-piperazinone

30 (±)-5-(2-Butynyl)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone

1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone

35 5(S)-Butyl-4-[1-(4-cyanobenzyl-2-methyl)-5-imidazolylmethyl]-1-(2,3-dimethylphenyl)-piperazin-2-one

- 231 -

4-[1-(2-(4-Cyanophenyl)-2-propyl)-5-imidazolylmethyl]-1-(3-chlorophenyl)-5(S)-(2-methylsulfonylethyl)piperazin-2-one

5 5(S)-*n*-Butyl-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-1-(2-methylphenyl)piperazin-2-one

4-[1-(4-Cyanobenzyl)-5-imidazolylmethyl]-5(S)-(2-fluoroethyl)-1-(3-chlorophenyl)piperazin-2-one

10 4-[3-(4-Cyanobenzyl)pyridin-4-yl]-1-(3-chlorophenyl)-5(S)-(2-methylsulfonylethyl)-piperazin-2-one


15 4-[5-(4-Cyanobenzyl)-1-imidazolylethyl]-1-(3-chlorophenyl)piperazin-2-one.






or a pharmaceutically acceptable salt or optical isomer thereof.

20 7. A compound which inhibits farnesyl-protein transferase which is:


- 232 -

Table 1




R

- 233 -

Table 2

5		<u>stereo.</u>	<u>X</u>	<u>Y</u>
	R	n-Bu	2-O(CH ₂) ₁₃ CH ₃	
10	S	n-Bu	2-O(CH ₂) ₁₃ CH ₃	
	R	n-Bu	2-O(CH ₂) ₁₁ CH ₃	
	S	n-Bu	2-O(CH ₂) ₁₁ CH ₃	
15	R	n-Bu	2-O(CH ₂) ₉ CH ₃	
	S	n-Bu	2-O(CH ₂) ₉ CH ₃	
20	R	n-Bu	2-O(CH ₂) ₇ CH ₃	
	S	n-Bu	2-O(CH ₂) ₇ CH ₃	
	R	n-Bu	2-O(CH ₂) ₃ Ph	
25	S	n-Bu	2-O(CH ₂) ₃ Ph	
	R	n-Bu	3-OCH ₂ Ph	


- 234 -

Table 2 (continued)

5	<u>stereo.</u>	<u>X</u>	<u>Y</u>
	S	n-Bu	3-OCH ₂ Ph
	R	n-Bu	3-O(CH ₂) ₁₃ CH ₃
10	S	n-Bu	3-O(CH ₂) ₁₃ CH ₃
	R	n-Bu	4-O(CH ₂) ₁₃ CH ₃
	S	n-Bu	4-O(CH ₂) ₁₃ CH ₃
15	S	n-Bu	4-OCH ₂ Ph
	R/S	H	2-O(CH ₂) ₁₃ CH ₃

20

- 235 -


Table 3

5

Y3-SO₂Me3-OCF₃


10

- 236 -

Table 4

	X	Y
5	CH ₂ OCH ₂ Ph	CF ₃
	CH ₂ SO ₂ Et	CF ₃
10	CH ₂ SO ₂ Ph	CF ₃
	CH ₂ SO ₂ Me	Cl
	CH ₂ SO ₂ Et	Cl
15	CONHMe	CF ₃
	CONHEt	CF ₃
20	CH ₂ SO ₂ Ph	Cl
	CONHMe	Cl
	CONHEt	Cl
25	CONHc-Pr	Cl
	CONHc-Pr	CF ₃


- 237 -

Table 4 (continued)

	<u>X</u>	<u>Y</u>
5	NHCOMe	Cl
	CONMe ₂	CF ₃
10	SO ₂ Et	Cl
	CH ₂ SMe	Cl
15	(±) C≡CMe	Cl


- 238 -

Table 5




5	X	Y
	CH ₂ CH ₂	H
	CH ₂ CO	H
10	CH ₂	H
	CH ₂	3-Cl
15	NHSO ₂	H
	CH ₂ CH ₂ CO	3-Cl
	CH ₂	2,3-Cl ₂
20	CH ₂	2-Br
	CH ₂	3-CF ₃
25	CH ₂	4-Cl
	CH ₂ CO	3-Cl
	CH ₂	3-F
30	CH ₂	3-Br

- 239 -

Table 6

5	X	Y	Z
	4-CN	CH ₂	5-CH ₂ CH ₂ CO
	3-CN	CH ₂	5-CH ₂
10	2-CN	CH ₂	5-CH ₂
	4-CN	CH ₂	4-CH ₂ CH ₂ CO
15	4-CN	(CH ₂) ₂	4-CH ₂ CH ₂ CO

- 240 -

Table 7

5


X	Y
---	---

4-Me	CH ₂ CH ₂ SO ₂ Me
------	--

10	4-Me	H
----	------	---

	2-Me	H
--	------	---

- 241 -

TABLE 8

Y

3- CF_3 3- CH_3 2,3- $(\text{CH}_2)_4$ 3- OCH_3


3-Cl

2- CH_3 , 3-Cl






H

3-F


- 242 -

TABLE 9


R








- 243 -

TABLE 10




- 244 -

TABLE 11







- 245 -

TABLE 12R


- 246 -

TABLE 12 (continued)R





- 247 -

TABLE 13R

- 248 -





TABLE 14

- 249 -




TABLE 15RY2,3-(CH₂)₄2-CH₃, 3-Cl

3-Cl

- 250 -


TABLE 16R

- 251 -



Table 17R

- 252 -


Table 18



5



or



10


or a pharmaceutically acceptable salt or optical isomer thereof.

15

8. The compound according to Claim 6 which is:

- 253 -

1-{5-[1-(4-nitrobenzyl)]imidazolymethyl}-2(S)-butyl-4-(1-naphthoyl)piperazine




5 or a pharmaceutically acceptable salt thereof.

9. The compound according to Claim 6 which is:

1-[5-(1-Benzylimidazol)methyl]-2(S)-butyl-4-(1-naphthoyl)piperazine

10




or a pharmaceutically acceptable salt or optical isomer thereof.

10. The compound according to Claim 6 which is:

15

1-(2(R)-Amino-3-(3-benzylthio)propyl)-2(S)-butyl-4-(1-naphthoyl)piperazine

- 254 -




or a pharmaceutically acceptable salt or optical isomer thereof.

5

11. The compound according to Claim 6 which is:

1-(2(R)-Amino-3-[3-(4-nitrobenzylthio)propyl])-2(S)-butyl-4-(1-naphthoyl)piperazine




10

or a pharmaceutically acceptable salt or optical isomer thereof.

12. The compound according to Claim 6 which is:

15 2(S)-*n*-Butyl-1-[1-(4-cyanobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine


- 255 -



or a pharmaceutically acceptable salt or optical isomer thereof.

5 13. The compound according to Claim 6 which is:

2(S)-*n*-Butyl-1-[1-(4-cyanobenzyl)imidazol-5-ylmethyl]-4-(2,3-dimethylphenyl)piperazin-5-one




10

or a pharmaceutically acceptable salt or optical isomer thereof.

14. The compound according to Claim 6 which is:

15 2(S)-*n*-Butyl-1-[1-(4-chlorobenzyl)imidazol-5-ylmethyl]-4-(1-naphthoyl)piperazine

- 256 -




or a pharmaceutically acceptable salt or optical isomer thereof.

5

15. The compound according to Claim 6 which is:

1-[[1-(4-cyanobenzyl)-1H-imidazol-5-yl]acetyl]-2(S)-n-butyl-4-(1-naphthoyl)piperazine

10




or a pharmaceutically acceptable salt or optical isomer thereof.

15

16. The compound according to Claim 6 which is:

1-[1-(4-Cyanobenzyl)imidazol-5-ylmethyl]-4-(2,3-dimethylphenyl)-2(S)-(2-methoxyethyl)piperazin-5-one


- 257 -



or a pharmaceutically acceptable salt or optical isomer thereof.

17. The compound according to Claim 6 which is:

5 5(S)-*n*-Butyl-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-1-(2-methylphenyl)piperazin-2-one




or a pharmaceutically acceptable salt or optical isomer thereof.

18. The compound according to Claim 6 which is:

15 (S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(methanesulfonyl)ethyl]-2-piperazinone

- 258 -




or a pharmaceutically acceptable salt or optical isomer thereof.

5

19. The compound according to Claim 6 which is:

(S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)ethyl]-2-piperazinone



10


or a pharmaceutically acceptable salt or optical isomer thereof.

15

20. The compound according to Claim 6 which is:

(S)-1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)methyl]-2-piperazinone

- 259 -




or a pharmaceutically acceptable salt or optical isomer thereof.

5

21. The compound according to Claim 6 which is:

1-(3-Chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolyl-methyl]-2-piperazinone



10

or a pharmaceutically acceptable salt or optical isomer thereof.

22. A pharmaceutical composition comprising a
15 pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 1.

23. A pharmaceutical composition comprising a
20 pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 2.

- 260 -

24. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 3.

5 25. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 6.

10 26. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 22.

15 27. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 23.


20 28. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 24.

 29. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 25.

25 30. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of the formula B:

30

- 261 -



wherein:

R^{1a} and R^{1b} are independently selected from:

- 5 a) hydrogen,
- b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, CN(R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
- 10 R¹¹OC(O)NR¹⁰-,
- c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocyclic, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-,
- 15 R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, and R¹¹OC(O)-NR¹⁰-;

20 R² and R³ are independently selected from: H; unsubstituted or substituted C₁-8 alkyl, unsubstituted or substituted C₂-8 alkenyl, unsubstituted or substituted C₂-8 alkynyl, unsubstituted or substituted aryl,

unsubstituted or substituted heterocycle,

$$\begin{array}{c} \text{NR}^6\text{R}^7 \\ | \\ \text{O} \end{array} \text{ or } \begin{array}{c} \text{OR}^6 \\ | \\ \text{O} \end{array}$$









wherein the substituted group is substituted with one or more of:

- 25 1) aryl or heterocycle, unsubstituted or substituted with:
- a) C₁-4 alkyl,
- b) (CH₂)_pOR⁶,

- 262 -

5

- c) $(\text{CH}_2)_p\text{NR}^6\text{R}^7$,
 d) halogen,
 e) CN,
 2) C3-6 cycloalkyl,
 3) OR^6 ,
 4) SR^{6a} , S(O)R^{6a} , SO_2R^{6a} ,



10

- 263 -





R^2 and R^3 are attached to the same C atom and are combined to form
 - $(\text{CH}_2)_u$ - wherein one of the carbon atoms is optionally replaced by a
 5 moiety selected from: O, $\text{S}(\text{O})_m$, $-\text{NC}(\text{O})-$, and $-\text{N}(\text{COR}^{10})-$;

R^4 is selected from H and CH_3 ;

10 and any two of R^2 , R^3 and R^4 are optionally attached to the same
 carbon atom;

R^6 , R^7 and R^{7a} are independently selected from: H; C₁₋₄ alkyl, C₃₋₆
 cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl,
 heteroarylsulfonyl, unsubstituted or substituted with:

- 15 a) C₁₋₄ alkoxy,
 b) aryl or heterocycle,
 c) halogen,
 d) HO,





20 g) $\text{N}(\text{R}^{10})_2$; or

R^6 and R^7 may be joined in a ring;
 R^7 and R^{7a} may be joined in a ring;

- 264 -

R^{6a} is selected from: C₁₋₄ alkyl, C₃₋₆ cycloalkyl, heterocycle, aryl, unsubstituted or substituted with:

- 5 a) C₁₋₄ alkoxy,
 b) aryl or heterocycle,
 c) halogen,
 d) HO,



10

R⁸ is independently selected from:

- a) hydrogen,
b) aryl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-,
15 R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN,
 NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
 R¹¹OC(O)NR¹⁰-, and
c) C₁₋₆ alkyl unsubstituted or substituted by aryl,
cyanophenyl, heterocycle, C₃₋₁₀ cycloalkyl, C₂₋₆
20 alkenyl, C₂₋₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-,
 R¹¹S(O)_m-, R¹⁰C(O)NH-, (R¹⁰)₂NC(O)-, R¹⁰₂N-
 C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
 R¹⁰OC(O)NH-;

25 R⁹ is selected from:

- a) hydrogen,
b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-,
R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-
C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃,
30 -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and

- 265 -

- c) C₁-C₆ alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

5

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

10

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR¹⁰-, -NR¹⁰C(O)-, O, -N(R¹⁰)-, -S(O)₂N(R¹⁰)-, -N(R¹⁰)S(O)₂-, or S(O)_m;

15

G is H₂;

V is selected from:

- a) hydrogen,
 b) heterocycle,
 20 c) aryl,
 d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
 e) C₂-C₂₀ alkenyl,

25 provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

W is imidazolyl;

30

X is -CH₂-, -C(=O)-, or -S(=O)_m-;

Z is

a unsubstituted or substituted group selected from aryl, heteroaryl, arylmethyl, heteroarylmethyl, arylsulfonyl,

- 266 -

heteroarylsulfonyl, wherein the substituted group is substituted with one or more of the following:

- 1) C₁₋₄ alkyl, unsubstituted or substituted with:
 - a) C₁₋₄ alkoxy,
 - b) NR⁶R⁷,
 - c) C₃₋₆ cycloalkyl,
 - d) aryl or heterocycle,
 - e) HO,
 - f) -S(O)_mR^{6a}, or
 - g) -C(O)NR⁶R⁷,
- 2) aryl or heterocycle,
- 3) halogen,
- 4) OR⁶,
- 5) NR⁶R⁷,
- 6) CN,
- 7) NO₂,
- 8) CF₃;
- 9) -S(O)_mR^{6a},
- 10) -C(O)NR⁶R⁷, or
- 11) C₃₋₆ cycloalkyl;


- | | |
|------|--|
| m is | 0, 1 or 2; |
| n is | 0, 1, 2, 3 or 4; |
| p is | 0, 1, 2, 3 or 4; |
| q is | 1 or 2; |
| r is | 0 to 5, provided that r is 0 when V is hydrogen; |
| s is | 0 or 1; |
| t is | 1; and |
| u is | 4 or 5; |

provided that the substituent (R⁸)_r- V - A¹(CR^{1a}₂)_nA²(CR^{1a}₂)_n - is H;

or a pharmaceutically acceptable salt thereof.

- 267 -

31. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of the formula:



wherein:

10

R^{1a} and R^{1b} are independently selected from:

15

- a) hydrogen,
- b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, CN(R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;
- c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocyclic, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, and R¹¹OC(O)-NR¹⁰-;


25

R² and R³ are independently selected from: H; unsubstituted or substituted C₁-8 alkyl, unsubstituted or substituted C₂-8 alkenyl,

- 268 -




unsubstituted or substituted C2-8 alkynyl, unsubstituted or substituted aryl,

unsubstituted or substituted heterocycle,








wherein the substituted group is substituted with one or more of:

- 1) aryl or heterocycle, unsubstituted or substituted with:
 - a) C1-4 alkyl,
 - b) $(\text{CH}_2)_p\text{OR}^6$,
 - c) $(\text{CH}_2)_p\text{NR}^6\text{R}^7$,
 - d) halogen,
 - e) CN,
- 2) C3-6 cycloalkyl,
- 3) OR^6 ,
- 4) SR^{6a} , $\text{S}(\text{O})\text{R}^{6a}$, SO_2R^{6a} ,



- 269 -



- 5 R^2 and R^3 are attached to the same C atom and are combined to form $\text{—(CH}_2\text{)}_u\text{—}$ wherein one of the carbon atoms is optionally replaced by a moiety selected from: O, S(O)_m , —NC(O)— , and $\text{—N(COR}^{10}\text{)—}$;


R^4 is selected from H and CH_3 ;

10

and any two of R^2 , R^3 and R^4 are optionally attached to the same carbon atom;

- 270 -


R⁶, R⁷ and R^{7a} are independently selected from: H; C₁-4 alkyl, C₃-6 cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, unsubstituted or substituted with:

- 5
- a) C₁-4 alkoxy,
 - b) aryl or heterocycle,
 - c) halogen,
 - d) HO,
 - e) ,
 - f) $-\text{SO}_2\text{R}^{11}$, or
 - g) N(R¹⁰)₂; or

10

R⁶ and R⁷ may be joined in a ring;
 R⁷ and R^{7a} may be joined in a ring;

15 R^{6a} is selected from: C₁-4 alkyl, C₃-6 cycloalkyl, heterocycle, aryl, unsubstituted or substituted with:

- a) C₁-4 alkoxy,
 - b) aryl or heterocycle,
 - c) halogen,
 - d) HO,
 - e) ,
 - f) $-\text{SO}_2\text{R}^{11}$, or
 - g) N(R¹⁰)₂;
- 20

R⁸ is independently selected from:

- 25
- a) hydrogen,
 - b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN,

- 271 -

NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
R¹¹OC(O)NR¹⁰-, and

- c) C₁-C₆ alkyl unsubstituted or substituted by aryl,
cyanophenyl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆
alkenyl, C₂-C₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-,
R¹¹S(O)_m-, R¹⁰C(O)NH-, (R¹⁰)₂NC(O)-, R¹⁰₂N-
C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
R¹⁰OC(O)NH-;

10 R⁹ is selected from:

- a) hydrogen,
b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-,
R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-
C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃,
-N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and
c) C₁-C₆ alkyl unsubstituted or substituted by perfluoroalkyl,
F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-,
(R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-,
R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

20

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and
aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

25

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-,
-C(O)-, -C(O)NR¹⁰-, -NR¹⁰C(O)-, O, -N(R¹⁰)-,
-S(O)₂N(R¹⁰)-, -N(R¹⁰)S(O)₂-, or S(O)_m;

30

G is H₂ or O;

V is selected from:

- a) hydrogen,
b) heterocycle,

- 272 -

- c) aryl,
 d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
 5 e) C₂-C₂₀ alkenyl,
 provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

W is a heterocycle;

10

X is -C(=O)- or -S(=O)_m-;

15

Z is a unsubstituted or substituted group selected from aryl, heteroaryl, arylmethyl, heteroarylmethyl, arylsulfonyl, heteroarylsulfonyl, wherein the substituted group is substituted with one or more of the following:

20

1) C₁-4 alkyl, unsubstituted or substituted with:

- a) C₁-4 alkoxy,
 b) NR⁶R⁷,
 c) C₃-6 cycloalkyl,
 d) aryl or heterocycle,
 e) HO,
 f) -S(O)_mR^{6a}, or
 g) -C(O)NR⁶R⁷,

25

2) aryl or heterocycle,

3) halogen,

4) OR⁶,

5) NR⁶R⁷,

6) CN,

30

7) NO₂,

8) CF₃;

9) -S(O)_mR^{6a},

10) -C(O)NR⁶R⁷, or

11) C₃-C₆ cycloalkyl;

- 273 -

m is 0, 1 or 2;
n is 0, 1, 2, 3 or 4;
p is 0, 1, 2, 3 or 4;
5 q is 1 or 2;
r is 0 to 5, provided that r is 0 when V is hydrogen;
s is 0 or 1;
t is 0 or 1; and
u is 4 or 5;

10 provided that if t is 1, then the substituent
(R⁸)_r-V-A¹(CR^{1a}₂)_nA²(CR^{1a}₂)_n- is H;

or a pharmaceutically acceptable salt thereof.

15 32. A method for treating cancer which comprises
administering to a mammal in need thereof a therapeutically effective
amount of a composition of Claim 22.

20 33. A method for treating cancer which comprises
administering to a mammal in need thereof a therapeutically effective
amount of a composition of Claim 23.

25 34. A method for treating cancer which comprises
administering to a mammal in need thereof a therapeutically effective
amount of a composition of Claim 24.

30 35. A method for treating cancer which comprises
administering to a mammal in need thereof a therapeutically effective
amount of a composition of Claim 25.

36. A method for treating neurofibromin benign
proliferative disorder which comprises administering to a mammal in

- 274 -

need thereof a therapeutically effective amount of a composition of Claim 23.

5 37. A method for treating blindness related to retinal vascularization which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 23.


10 38. A method for treating infections from hepatitis delta and related viruses which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 23.

15 39. A method for preventing restenosis which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 23.

 40. A method for treating polycystic kidney disease which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 23.

20 41. A method for treating or preventing a disease selected from cancer, neurofibromin benign proliferative disorder, blindness related to retinal vascularization, infections from hepatitis delta and related viruses, restenosis and polycystic kidney disease, which
25 comprises administering to a mammal in need thereof a therapeutically effective amount of a pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of the formula B:

- 275 -





wherein:

R^{1a} and R^{1b} are independently selected from:

- 5 a) hydrogen,
- b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, CN(R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-,
- 10 c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocyclic, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, and R¹¹OC(O)-NR¹⁰-;
- 15

R² and R³ are independently selected from: H; unsubstituted or substituted C₁-8 alkyl, unsubstituted or substituted C₂-8 alkenyl, unsubstituted or substituted C₂-8 alkynyl, unsubstituted or substituted aryl,

unsubstituted or substituted heterocycle,  or ,

wherein the substituted group is substituted with one or more of:

- 25 1) aryl or heterocycle, unsubstituted or substituted with:
 - a) C₁-4 alkyl,
 - b) (CH₂)_pOR⁶,

- 276 -

c) $(\text{CH}_2)_p\text{NR}^6\text{R}^7$,

d) halogen,

e) CN,

2) C₃₋₆ cycloalkyl,3) OR^6 ,4) SR^{6a} , S(O)R^{6a} , SO_2R^{6a} ,5) $-\text{NR}^6\text{R}^7$,6)
$$\begin{array}{c} \text{R}^6 \\ | \\ -\text{N}-\text{C}(=\text{O})-\text{R}^7 \end{array}$$
,7)
$$\begin{array}{c} \text{R}^6 \\ | \\ -\text{N}-\text{C}(=\text{O})-\text{NR}^7\text{R}^{7a} \end{array}$$
,8)
$$\begin{array}{c} -\text{O}-\text{C}(=\text{O})-\text{NR}^6\text{R}^7 \end{array}$$
,9)
$$\begin{array}{c} -\text{O}-\text{C}(=\text{O})-\text{OR}^6 \end{array}$$
,10)
$$\begin{array}{c} \text{CH}_3 \\ | \\ \text{C}(=\text{O})-\text{NR}^6\text{R}^7 \end{array}$$
,11) $-\text{SO}_2-\text{NR}^6\text{R}^7$,12)
$$\begin{array}{c} \text{R}^6 \\ | \\ -\text{N}-\text{SO}_2-\text{R}^{6a} \end{array}$$
,

- 277 -





R^2 and R^3 are attached to the same C atom and are combined to form
 - $(\text{CH}_2)_u$ - wherein one of the carbon atoms is optionally replaced by a
 5 moiety selected from: O, $\text{S}(\text{O})_m$, $-\text{NC}(\text{O})-$, and $-\text{N}(\text{COR}^{10})-$;

R^4 is selected from H and CH_3 ;

10 and any two of R^2 , R^3 and R^4 are optionally attached to the same
 carbon atom;

R^6 , R^7 and R^{7a} are independently selected from: H; C₁₋₄ alkyl, C₃₋₆
 cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl,
 heteroarylsulfonyl, unsubstituted or substituted with:

- 15 a) C₁₋₄ alkoxy,
 b) aryl or heterocycle,
 c) halogen,
 d) HO,




20 g) $\text{N}(\text{R}^{10})_2$; or

R^6 and R^7 may be joined in a ring;
 R^7 and R^{7a} may be joined in a ring;

- 278 -

R^{6a} is selected from: C₁₋₄ alkyl, C₃₋₆ cycloalkyl, heterocycle, aryl, unsubstituted or substituted with:

- a) C₁₋₄ alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,
- e) 
- f) —SO₂R¹¹, or
- g) N(R¹⁰)₂;

10

R⁸ is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, C_{3-C10} cycloalkyl, C_{2-C6} alkenyl, C_{2-C6} alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and
- c) C_{1-C6} alkyl unsubstituted or substituted by aryl, cyanophenyl, heterocycle, C_{3-C10} cycloalkyl, C_{2-C6} alkenyl, C_{2-C6} alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NH-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹⁰OC(O)NH-;

15

20

25 R⁹ is selected from:

- a) hydrogen,
- b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and

30

- 279 -

- c) C₁-C₆ alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

5

R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

10

A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR¹⁰-, -NR¹⁰C(O)-, O, -N(R¹⁰)-, -S(O)₂N(R¹⁰)-, -N(R¹⁰)S(O)₂-, or S(O)_m;

15

G is H₂;

V is selected from:

- a) hydrogen,
 b) heterocycle,
 20 c) aryl,
 d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
 e) C₂-C₂₀ alkenyl,

25

provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

W is imidazolyl;

30

X is -CH₂-, -C(=O)-, or -S(=O)_m-;

Z is

a unsubstituted or substituted group selected from aryl, heteroaryl, arylmethyl, heteroarylmethyl, arylsulfonyl,

- 280 -

heteroarylsulfonyl, wherein the substituted group is substituted with one or more of the following:

- 1) C₁₋₄ alkyl, unsubstituted or substituted with:
 - a) C₁₋₄ alkoxy,
 - b) NR⁶R⁷,
 - c) C₃₋₆ cycloalkyl,
 - d) aryl or heterocycle,
 - e) HO,
 - f) -S(O)_mR^{6a}, or
 - g) -C(O)NR⁶R⁷,
- 2) aryl or heterocycle,
- 3) halogen,
- 4) OR⁶,
- 5) NR⁶R⁷,
- 6) CN,
- 7) NO₂,
- 8) CF₃;
- 9) -S(O)_mR^{6a},
- 10) -C(O)NR⁶R⁷, or
- 11) C₃₋₆ cycloalkyl;


- m is 0, 1 or 2;
 n is 0, 1, 2, 3 or 4;
 p is 0, 1, 2, 3 or 4;
 q is 1 or 2;
 r is 0 to 5, provided that r is 0 when V is hydrogen;
 s is 0 or 1;
 t is 1; and
 u is 4 or 5;

provided that the substituent (R⁸)_r - V - A¹(CR^{1a}₂)_nA²(CR^{1a}₂)_n - is H;

or a pharmaceutically acceptable salt thereof.

- 281 -

42. A method for treating or preventing a disease selected from cancer, neurofibromin benign proliferative disorder, blindness related to retinal vascularization, infections from hepatitis delta and related viruses, restenosis and polycystic kidney disease, which comprises administering to a mammal in need thereof a therapeutically effective amount of a pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of the formula:



wherein:

R^{1a} and R^{1b} are independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, CN(R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-,
- c) unsubstituted or substituted C₁-C₆ alkyl wherein the substituent on the substituted C₁-C₆ alkyl is selected from unsubstituted or substituted aryl, heterocyclic, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, and R¹¹OC(O)-NR¹⁰-;

- 282 -




R^2 and R^3 are independently selected from: H; unsubstituted or substituted C₁-8 alkyl, unsubstituted or substituted C₂-8 alkenyl, unsubstituted or substituted C₂-8 alkynyl, unsubstituted or substituted aryl,

unsubstituted or substituted heterocycle, $\text{C}(=\text{O})\text{NR}^6\text{R}^7$ or $\text{C}(=\text{O})\text{OR}^6$,








5 wherein the substituted group is substituted with one or more of:

- 1) aryl or heterocycle, unsubstituted or substituted with:
 - a) C₁-4 alkyl,
 - b) $(\text{CH}_2)_p\text{OR}^6$,
 - c) $(\text{CH}_2)_p\text{NR}^6\text{R}^7$,
 - d) halogen,
 - e) CN,
- 2) C₃-6 cycloalkyl,
- 3) OR^6 ,
- 4) SR^{6a} , $\text{S}(\text{O})\text{R}^{6a}$, SO_2R^{6a} ,

15



- 283 -



- 5 R^2 and R^3 are attached to the same C atom and are combined to form $\text{—(CH}_2)_u\text{—}$ wherein one of the carbon atoms is optionally replaced by a moiety selected from: O, S(O)_m , —NC(O)— , and $\text{—N(COR}^{10})\text{—}$;


R^4 is selected from H and CH_3 ;

10

and any two of R^2 , R^3 and R^4 are optionally attached to the same carbon atom;

- 284 -


R⁶, R⁷ and R^{7a} are independently selected from: H; C₁-4 alkyl, C₃-6 cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, unsubstituted or substituted with:

- 5
- a) C₁-4 alkoxy,
 - b) aryl or heterocycle,
 - c) halogen,
 - d) HO,
 - e) 
 - f) $-\text{SO}_2\text{R}^{11}$, or
 - g) N(R¹⁰)₂; or

10

R⁶ and R⁷ may be joined in a ring;
R⁷ and R^{7a} may be joined in a ring;

15 R^{6a} is selected from: C₁-4 alkyl, C₃-6 cycloalkyl, heterocycle, aryl, unsubstituted or substituted with:

- a) C₁-4 alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,
- e) 
- f) $-\text{SO}_2\text{R}^{11}$, or
- g) N(R¹⁰)₂;

20

R⁸ is independently selected from:

- 25
- a) hydrogen,
 - b) aryl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN,

- 285 -

NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
R¹¹OC(O)NR¹⁰-, and

- c) C₁-C₆ alkyl unsubstituted or substituted by aryl,
cyanophenyl, heterocycle, C₃-C₁₀ cycloalkyl, C₂-C₆
alkenyl, C₂-C₆ alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-,
R¹¹S(O)_m-, R¹⁰C(O)NH-, (R¹⁰)₂NC(O)-, R¹⁰₂N-
C(NR¹⁰)-, CN, R¹⁰C(O)-, R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or
R¹⁰OC(O)NH-;

10 R⁹ is selected from:

- a) hydrogen,
b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R¹⁰O-,
R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-, (R¹⁰)₂NC(O)-, R¹⁰₂N-
C(NR¹⁰)-, CN, NO₂, R¹⁰C(O)-, R¹⁰OC(O)-, N₃,
-N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-, and
c) C₁-C₆ alkyl unsubstituted or substituted by perfluoroalkyl,
F, Cl, Br, R¹⁰O-, R¹¹S(O)_m-, R¹⁰C(O)NR¹⁰-,
(R¹⁰)₂NC(O)-, R¹⁰₂N-C(NR¹⁰)-, CN, R¹⁰C(O)-,
R¹⁰OC(O)-, N₃, -N(R¹⁰)₂, or R¹¹OC(O)NR¹⁰-;

20 R¹⁰ is independently selected from hydrogen, C₁-C₆ alkyl, benzyl and
aryl;

R¹¹ is independently selected from C₁-C₆ alkyl and aryl;

25 A¹ and A² are independently selected from: a bond, -CH=CH-, -C≡C-,
-C(O)-, -C(O)NR¹⁰-, -NR¹⁰C(O)-, O, -N(R¹⁰)-,
-S(O)₂N(R¹⁰)-, -N(R¹⁰)S(O)₂-, or S(O)_m;

30 G is H₂ or O;

V is selected from:

- a) hydrogen,
b) heterocycle,

- 286 -

- c) aryl,
d) C₁-C₂₀ alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
5 e) C₂-C₂₀ alkenyl,
provided that V is not hydrogen if A¹ is S(O)_m and V is not hydrogen if A¹ is a bond, n is 0 and A² is S(O)_m;

W is a heterocycle;

10

X is --C(=O)- or -S(=O)_m-;

Z is a unsubstituted or substituted group selected from aryl, heteroaryl, arylmethyl, heteroarylmethyl, arylsulfonyl, heteroarylsulfonyl, wherein the substituted group is substituted with one or more of the following:

15

1) C₁-4 alkyl, unsubstituted or substituted with:

20

- a) C₁-4 alkoxy,
b) NR⁶R⁷,
c) C₃-6 cycloalkyl,
d) aryl or heterocycle,
e) HO,
f) -S(O)_mR^{6a}, or
g) -C(O)NR⁶R⁷,

25

2) aryl or heterocycle,

3) halogen,

4) OR⁶,

5) NR⁶R⁷,

6) CN,

30

7) NO₂,

8) CF₃;

9) -S(O)_mR^{6a},

10) -C(O)NR⁶R⁷, or

11) C₃-C₆ cycloalkyl;

- 287 -

- m is 0, 1 or 2;
n is 0, 1, 2, 3 or 4;
p is 0, 1, 2, 3 or 4;
5 q is 1 or 2;
r is 0 to 5, provided that r is 0 when V is hydrogen;
s is 0 or 1;
t is 0 or 1; and
u is 4 or 5;

10

provided that if t is 1, then the substituent
(R⁸)_r-V-A¹(CR^{1a}₂)_nA²(CR^{1a}₂)_n- is H;

or a pharmaceutically acceptable salt thereof.

15

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 96/04019

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D233/54 A61K31/495 A61K31/415 C07D241/08 C07D295/18
C07D401/06 C07D401/14 C07D403/06 C07D403/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BE,A,820 242 (MALESCI S.A.S ISTITUTO FARMACOBIOLOGICO) 16 January 1975 see page 3 - page 6; table 1 see page 12 - page 24; examples ---	2,5
X	WO,A,92 12134 (NEUROGEN CORP) 23 July 1992 see page 26 - page 27; example VIII ---	2,5
X	EP,A,0 064 705 (MERRELL DOW PHARMA) 17 November 1982 see page 13 - page 16; examples 6-9 --- -/--	2,5

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

19 July 1996

Date of mailing of the international search report

01.08.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (-31-70) 340-2040, Tx. 31 651 epo nl,
Fax (-31-70) 340-3016

Authorized officer

Fink, D

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 96/04019

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 35, no. 22, 30 October 1992, WASHINGTON US, pages 4118-4134, XP002008877 E. CARCELLER ET AL.: "(Pyridylcyanomethyl)piperazines as Orally Active PAF Antagonists" see table I; compounds no. 30-32 and 34 see table IV; compounds no. 68-70 ---	1,4
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 7, no. 2, March 1964, WASHINGTON US, pages 154-158, XP002008878 O.E. FANCHER ET AL.: "4-Alkyl- (or Aralkyl) 1-Aryl-2-piperazones" see page 156; table I ---	2,3,5
A	WO,A,95 00497 (MERCK & CO INC ;GRAHAM SAMUEL L (US); WILLIAMS THERESA M (US)) 5 January 1995 see the whole document ---	1-42
P,X	US,A,5 478 934 (J. YUAN ET AL.) 26 December 1995 see column 13 - column 14; example II ---	2,5
P,X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 38, no. 12, 9 June 1995, WASHINGTON US, pages 2251-2255, XP002008879 A. THURKAUF ET AL.: "2-Phenyl-4-(aminomethyl)imidazoles as Potential Antipsychotic Agents. Synthesis as Dopamine D2 Receptor Binding" see page 2263; 2nd column, compound no. 29 -----	2,5

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/04019

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 26-42 are directed to a method of treatment of (diagnostic method practised on) the human/animal body the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claims not searched: 10, 11, 15
see next page
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US96/04019

FURTHER INFORMATION CONTINUED FROM PCT/ISA/

The novelty search on the final compounds of general formulae A-C revealed a vast amount of novelty-destroying compounds with respect to claims 1-3 of the present application.

Therefore, the search and the search report - as far as the novelty of the compounds of formulae A-C is concerned - has been limited (for economical reasons; cf. WIPO: PCT Search Guidelines, 1992; Chapter III, item 2) to the

1-({opt. subst.}imidazol-5(or 4)-ylmethyl)-
4-({opt. subst.}aryl (or {opt. subst.}arylcarbonyl))-
(opt. subst)piperazines/piperazinones

according to the present application.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 96/04019

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
BE-A-820242	16-01-75	NONE	
WO-A-9212134	23-07-92	US-A- 5159083	27-10-92
		CA-A- 2098301	29-06-92
		EP-A- 0640075	01-03-95
		JP-T- 6504054	12-05-94
EP-A-0064705	17-11-82	US-A- 4371737	01-02-83
		AU-B- 547243	10-10-85
		AU-B- 8317982	11-11-82
		CA-A- 1175827	09-10-84
		GB-A, B 2098605	24-11-82
		JP-C- 1577512	13-09-90
		JP-B- 2004221	26-01-90
		JP-A- 57185267	15-11-82
WO-A-9500497	05-01-95	AU-B- 7041294	17-01-95
		CA-A- 2165176	05-01-95
		EP-A- 0703905	03-04-96
US-A-5478934	26-12-95	WO-A- 9616057	30-05-96